

Publication 7

Josué Dejinnin Georges Avakoudjo, Fouad Kolawalé Yde Soumanou, Fred Dètonджи Hodonou, Euodias Djengue .

Sertoli Cell-Only Syndrome with Normal Karyotype.

Journal of Urology and Renal Diseases 2018; 02 : 178

Case Report

Sertoli Cell-Only Syndrome with Normal Karyotype

Josué Dejinnin Georges Avakoudjo¹, Fouad Kolawalé Yde Soumanou^{1*}, Fred Dètonndji Hodonou¹, Euodias Djengue²

¹Department of urology at university hospital of Cotonou, Benin

²Department of gynecology-obstetric at hospital of calavi, Benin

*Corresponding author: Fouad Kolawalé Yde Soumanou, Urologist and hospital practitioner at university hospital of Cotonou, 06 BP 2538, Cotonou, Benin. Tel: 229-96880788/95406504; Email: soumfou@yahoo.fr

Citation: Avakoudjo JDG, Soumanou FKY, Hodonou FD, Djengue E (2018) Sertoli Cell-Only Syndrome with Normal Karyotype. J Urol Ren Dis: JURD-178. DOI: 10.29011/2575-7903.000078

Received Date: 20 February, 2018; Accepted Date: 26 February, 2018; Published Date: 06 March, 2018

Abstract

Sertoli cell-only syndrome is usually associated with micro deletion Y chromosome. We reported rare case in the 37years-old-man married 10 years ago. A left varicocele was associated with this syndrome. No micro deletion Y chromosome was noted. Left varicocelectomy has been performed.

Keywords: Azoospermia; Sertoli Cell-Only Syndrome; Testicular Biopsy.

Introduction

The seminiferous epithelium contains both somatic cells (Sertoli cells) and germ cells (from the early stem cells to spermatozoa). Sertoli cells perform crucial functions that initiate and sustain spermatogenesis. Consequently, an abnormality in their function may disrupt the full progression of spermatogenesis [1,2]. Azoospermia due to spermatogenic failure, or non-obstructive azoospermia, affects approximately 1% of the general population and 10%-15% of men seeking an infertility evaluation [3,4].

Case Report

A 37 year-old-man patient presented with primary infertility. He was married 10 years ago. He was not decreased libido or erectile dysfunction. His medical (chlamydia, Mycoplasma, syphilis, tuberculosis) and surgical history were negative. On examination, height was 175 cm, weight 74 kg. Body hair was normal. Sexual maturing rating was normal with a testicular volume bilaterally. He had not gynecomastia. Biological screening shows creatinine level was 6 mg/L, hemoglobin was 14 g/dL, and white blood cell count was 6 G/L. Semen analysis showed azoospermia on three consecutive samples even after centrifugation. Doppler color ultrasound of the scrotum showed left varicocele. The testicular, the epididymis and the vas deferens was normal.

Serum Follicle-Stimulating Hormone (FSH) level was 45,93 mIU/ml [1-5], Serum Luteinizing Hormone (LH) level -13,14 mIU/ml [1-8] and total serum testosterone - 4,26 ng/ml [2,5-8]. Serum

Prolactin level - 15,4 ng/ml [3-25]. Patient has a normal Karyotype (Normal 46, XY karyotype, as shown Giemsa-trypsin [G-banding]). Left varicocelectomy has been performed and testicular biopsy. The postoperative course was uneventful. His testicular biopsy revealed Sertoli Cell-Only Syndrome (SCOS). However, the patient done an Intracytoplasmic Sperm Injection (ICSI) after Testicular Sperm Extraction (TESE) successfully to be father.

Discussion

Primary testicular failure occurs in approximately 1% of all males, and is present in 10% of those obtaining medical consultation for infertility [5]. Sertoli cell-only syndrome (SCOS), [5,6] also known as germinal cell aplasia, can be found in a few testicular biopsies that are performed in these patients. The pathognomonic feature of SCOS is the absence of germ cells. The seminiferous tubules are lined by Sertoli cells that may be either immature (prepubertal) or mature or have other changes that may correlate with specific etiologies and clinical findings [7]. For the prognostic purpose, Anniballo et al. [5] divided SCOS into two categories: Pure (congenital) and mixed (secondary). The pure form is caused by failure of migration of germ cells, [8] while the mixed form is related to the postnatal damage to previously healthy testicular tissue. Retrieving germ cells in cases of pure SCOS is the impossible [9] and proper identification of the type of SCOS is essential to spare unnecessary medical expense and inconvenience to patients. The combination of increased inhibin B and normal serum FSH levels also indicates the presence of spermatids [10]. However, the role of inhibin B in predicting the presence of sperm within the testis in patients with NOA is still controversial [11].

Although some cases of azoospermia due to SCOS (approximately 10%-15%) are due to microdeletions of variable portions of the AZF region of the Y chromosome, the etiology of most is unknown. The correct identification of infertile patients with true SCOS by using selective molecular markers for germ cells might help in establishing the true incidence of this problem and in directing research toward understanding the causes of this defect [12]. Many investigators have suggested that as more genes are deleted, spermatogenesis defects may progress from hypospermatogenesis to SCOS. Cases with extensive deletions involving AZFa, AZFb, and AZFc are almost invariably associated with SCOS [13]. Ying Hui Lin et al. said [14], this case had deletions of the whole long arm of the Y chromosome, and only Sertoli cells were found in the testicular specimen, a finding in concordance with previous reports [15,16].

However, many of these patients have a normal XY karyotype with normal secondary male sexual characteristics, yet are infertile and azoospermic or severe oligozoospermic. karyotype. (Normal 46, XY karyotype, as shown Giemsa-trypsin [G-banding], but on polymerase chain reaction amplification or fluorescence in situ hybridisation microdeletions of the Y chromosome are seen, rarely macroscopic deletions of the Y-chromosome long arm that are detectable by karyotyping [i.e. 46, X, Yq-]) [7]. Predictive factors for the presence of spermatozoa within the testis have been actively studied, including serum FSH and inhibin-B levels, testicular volume, and testicular histopathology [17-18].

The analysed patients with Sertoli cell only syndrome have significantly elevated serum FSH level. The distinctly elevated basal serum LH levels and serum testosterone levels in normal range, demonstrated that there was compensated dysfunction of the Leydig cells in patients with Sertoli cell only syndrome. The elevation of gonadotropins in patients with this condition was not so high and levels of serum testosterone not lowered, as those seen in the patients with Klinefelter's syndrome [19] Suominen et al. [20] concluded that there is not any relation between testicular morphology (i.e. spermatogenesis) and serum prolactin, that increased prolactin values are usually associated with low testosterone values and that not correlation exists between serum gonadotrophin and prolactin levels. Intracytoplasmic Sperm Injection (ICSI) and with testicular sperm extraction (TESE) can be useful in retrieving sperm from these men with mixed SCOS [21,22].

Conclusion

The sertoli cell-only syndrome is often of genetic origin with micro deletion of the Y chromosome. It's often associated by the serum FSH, LH and serum testosterone level rising. This micro deletion may not exist but exceptionally.

References

1. Russell LD (1993) Morphological and functional evidence for Sertoli-germ cells relationships. In: Russell LD, Griswold MD, eds. *The Sertoli cell*. Clearwater, FL: Cache River Press 1993: 365-390.
2. de Kretser DM, Loveland KL, Meinhardt A, Simorangkir D, Wreford N (1998) Spermatogenesis. *Hum Reprod* 13: 1-8.
3. Willott GM (1982) Frequency of azoospermia. *Forensic Sci Int* 20: 9-10.
4. Jarow JP, Espeland MA, Lipshultz LI (1989) Evaluation of the azoospermic patient. *J Urol* 142: 62-65.
5. Anniballo R, Ubaldi F, Cobellis L, Sorrentino M, Rienzi L, et al. (2000) Criteria predicting the absence of spermatozoa in the Sertoli cell-only syndrome can be used to improve success rates of sperm retrieval. *Hum Reprod* 15: 2269-2277.
6. Nistal M and Paniagua R (1997) Non-neoplastic diseases of the testis. In: Bostwick DG, Eble JN, editors. *Urologic Surgical Pathology*. St. Louis: Mosby 1997: 498-501.
7. Babureddy Hanmayyagari, Mounika Guntaka, Srinagesh (2015) A rare case of male infertility: Sertoli only syndrome. *Chrismed Journal of Health and Research* 2: 64-67.
8. Wartenberg H (1989) Differentiation and development of the testes. In: Burger A, de Kretser, DM, editors, *The Testis*. New York, USA: Raven Press 1989: 67-118.
9. Su LM, Palermo GD, Goldstein M, Veeck LL, Rosenwaks Z, et al. (1999) Testicular sperm extraction with intracytoplasmic sperm injection for nonobstructive azoospermia: Testicular histology can predict success of sperm retrieval. *J Urol* 161: 112-116.
10. von Eckardstein S, Simoni M, Bergmann M, Weinbauer GF, Gassner P, et al. (1999) Serum inhibin B in combination with serum follicle-stimulating hormone (FSH) is a more sensitive marker than serum FSH alone for impaired spermatogenesis in men, but cannot predict the presence of sperm in testicular tissue samples. *J Clin Endocrinol Metab* 84: 2496-2501.
11. Orna Weller, Leah Yogev, Haim Yavetz (2005) Differentiating between primary and secondary Sertoli-cell-only syndrome by histologic and hormonal parameters *Fertil Steril* 83: 1856-1858.
12. Patrizio P, Ricci SM, Tomaszewski JE, Hecht NB (2000) Identification of meiotic and postmeiotic gene expression in testicular tissue of patients histologically classified as Sertoli cell only. *Fertility and Sterility* 74: 785-790.
13. Foresta C, Ferlin A, Garolla A, Moro E, Pistorello M, et al. (1998) High frequency of well-defined Y-chromosome deletions in idiopathic. Sertoli-cell syndrome. *Hum Reprod* 13: 302-307.
14. Ying Hui Lin, Louise Chuang, Yung Ming Lin, Ying Hung Lin, M. Yeng Ni Teng, et al. (2005) Isochromosome of Yp in a man with Sertoli-cell-only syndrome. *Fertility and Sterility* 83: 764-766.
15. Bablok L, Janczewski Z, Kwiatkowska Z, St. FRACKI. (1978) The Relationship between Plasma FSH, Testosterone Levels and Testicular Histology in Males with Azoospermia. *Andrologia* 10: 502-505.

16. Oxana Blagosklonova, Christiane Joanne, Christophe Roux, Christophe Tripogney, Florence Fellmann, et al. (2003) Sertoli cell and Leydig cell immunophenotypes inpatients with or without AZF microdeletions. *Fertility and sterility* 79: 1146-1148.
17. Bohring C, Schroeder-Printzen I, Weidner W, Krause W (2002) Serum levels of inhibin B and follicle-stimulating hormone may predict successful sperm retrieval in men with azoospermia who are undergoing testicular sperm extraction. *Fertil Steril* 78: 1195-1198.
18. Boitrelle F, Robin G, Marcelli F, Albert M, Leroy-Martin B, et al. (2011) A predictive score for testicular sperm extraction quality and surgical ICSI outcome in non-obstructive azoospermia: a retrospective study. *Hum Reprod* 26: 3215-3221.
19. Micic S, Ilic V, Micic M, Cenbacev O, Dotlic R (1983) Endocrine Profile of 45 Patients with Sertoli Cell Only Syndrome. *Andrologia* 15: 228-232.
20. Suominen JJO, Nikkanen V, Multamxki S, Hyyppa M (1979) Prolactin in Azoospermic Men and its Relation to Testicular Morphology, Serum Testosterone and Gonadotrophin Levels. *andrologia* 11: 15-17.
21. Schwarzer JU, Fiedler K, v Hertwig I, Krüsmann G, Würfel W, et al. (2003) Sperm retrieval procedures and intracytoplasmatic spermatozoa injection with epididymal and testicular sperms. *Urol Int* 70: 119-123.
22. Seo JT and Ko WJ (2001) Predictive factors of successful testicular sperm recovery in non-obstructive azoospermia patients. *Int J Androl* 24: 306-310.