




Journal of Mahani Mathematical Research



Shahid Bahonar
University of Kerman

Print ISSN: 2251-7952 Online ISSN: 2645-4505

Home Browse ▾ Journal Info ▾ Guide for Authors Submit Manuscript Reviewers Contact Us Login Register



Journal of Mahani Mathematical Research (JMMR) was approved by the Ministry of Science, Research and Technology of Iran with rank (A).

<https://journals.msrt.ir/home/detail/15002/>

ISC Citation Impact (2021): 0.077

ISC Prompt Impact (2020): 0.143

=====

Call for Papers for Vol. 13, Issue 3 (2024)

Special Issue

Dedicated to Prof. Esfandiar Eslami

Journal of Mahani Mathematical Research (JMMR)

Journal of Mahani Mathematical Research is an international semiannual journal published by the Shahid Bahonar University of Kerman.

This journal has decided to publish Vol. 13, Issue 3 (2024) in honor of Prof. Esfandiar Eslami

Publication Information

Publisher

Shahid Bahonar University of Kerman

Editor-in-Chief

Arsham Borumand Saeid

Director-in-Charge

Nosratollah Shajareh
Poursalavati

Frequency

Semiannual

– Volume 13 (2024)

📄 Issue 1

+ Volume 12 (2023)

+ Volume 11 (2022)

+ Volume 10 (2021)

+ Volume 9 (2020)

+ Volume 8 (2019)

+ Volume 7 (2018)

+ Volume 6 (2017)

+ Volume 5 (2016)

+ Volume 4 (2015)

+ Volume 3 (2014)

+ Volume 2 (2013)

+ Volume 1 (2012)

A. Borumand Saeid (Lead Guest Editor), Department of Pure Mathematics, Faculty of Mathematics and Computer, Shahid Bahonar University of Kerman, Kerman, Iran. Email: arsham@uk.ac.ir.

R. A. Borzooei, Department of Mathematics, Shahid Beheshti University, Tehran, Iran. Email: borzooei@sbu.ac.ir.

B. Davvaz, Department of Mathematics Yazd University, Yazd, Iran. Email: davvaz@yazd.ac.ir.
M. Golshani School of Mathematics, Institute for Research in Fundamental Sciences (IPM) Iran. Email: golshani.m@gmail.com.

M. Mashinchi, Department of Statistics, Faculty of Mathematics and Computer, Shahid Bahonar University of Kerman, Kerman, Iran. Email: mashinchi@uk.ac.ir

M. Moniri, Department of Mathematics, Shahid Beheshti University, Tehran, Iran. E-mail: m-moniri@sbu.ac.ir

=====

Call for Papers for Vol. 13, Issue 4 (2024)

Special Issue

Dedicated to memory of Prof. Mahbanoo Tata

Journal of Mahani Mathematical Research (JMMR)

Journal of Mahani Mathematical Research is an international semiannual journal published by the Shahid Bahonar University of Kerman. The editorial board of JMMR has decided to publish Vol. 13, Issue 4 (2024) in honor of Prof. Mahbanoo Tata for her valuable services, training many of students and outstanding the fields Statistics, Fuzzy statistics, Information theory and applications, and other related fields showcasing her research.

Manuscripts should be written in English, under the LaTeX style of the journal. Submissions should be sent online until May 15, 2024 in the website of the journal (<https://jmmrc.uk.ac.ir/>).

Note: Please add the following comment in the comment section when submitting the paper to

Advanced Search

Indexing and Abstracting

zbMATH

fidmath
Fachinformationsdienst Mathematik

EBSCO

ArabWorld Research Source: Al Masdar
Database Coverage List

EBSCO
HOST

DICTIONARY OF

sn:2251-7952



Fields ▾

Operators ▾

Help ▾

Journal of Mahani Mathematical Research Center

Short Title: J. Mahani Math. Res. Cent.

Publisher: Mahani Mathematical Research Center c/o Shahid Bahonar University of Kerman, Faculty of Mathematics & Computer, Kerman

ISSN: 2251-7952; 2645-4505/e

Online: http://jmmrc.uk.ac.ir/browse?_action=issue 

Comments: Journal; Indexed cover-to-cover; This journal is available open access.

Documents Indexed: [181 Publications](#) (since 2012)

References Indexed: [127 Publications](#) with 2,813 References.

Latest Issues

all ▾

12, No. 2 (2023)

12, No. 1 (2023)

Authors

4 Arabpour, Alireza

4 Najafzadeh, Shahram

all ▾

Fields

21 Statistics (62-XX)

19 Numerical analysis (65-XX)

all ▾

10.22103/jmmr.2023.19918.1303



Fields ▾

Operators ▾

Help ▾

Degla, G.; Zinsou, P. H.**Quantitative and stability study of the evolution of a viscoelastic body. (English)** Zbl 07708025**J. Mahani Math. Res. Cent. 12, No. 2, 443-458 (2023).**

Summary: This paper deals with a generalization of the model describing the evolution of a linear viscoelastic body studied by *M. Kirane* and *B. Said-Houari* in [Z. Angew. Math. Phys. 62, No. 6, 1065–1082 (2011; Zbl 1242.35163)]. We prove the existence and uniqueness of the solution of the model using a C_0 -semi-group contraction method with a linear operator parameter. Moreover the strong stability of the solution is shown in a particular case.

MSC:

- 74D05 Linear constitutive equations for materials with memory
- 74H20 Existence of solutions of dynamical problems in solid mechanics
- 74H25 Uniqueness of solutions of dynamical problems in solid mechanics
- 74H55 Stability of dynamical problems in solid mechanics
- 35Q74 PDEs in connection with mechanics of deformable solids
- 35L20 Initial-boundary value problems for second-order hyperbolic equations

QUANTITATIVE AND STABILITY STUDY OF THE EVOLUTION OF A VISCOELASTIC BODY

G. DEGLA  AND P.H. ZINSOU  ✉

Article type: Research Article

(Received: 18 July 2022, Received in revised form 11 December 2022)

(Accepted: 28 January 2023, Published Online: 29 January 2023)

ABSTRACT. This paper deals with a generalization of the model describing the evolution of a linear viscoelastic body studied by Kirane M. and B.S. Houari in 2011. We prove the existence and uniqueness of the solution of the model using a C_0 -semi-group contraction method with a linear operator parameter. Moreover the strong stability of the solution is shown in a particular case.

Keywords: Viscoelasticity; Homogeneous evolution equation; Monotone maximal operator

2020 MSC: Primary 47D06, 92D15, 65J08, 47H05.

1. Introduction

Several mathematical models that come from physics (viscoelasticity motion) lead to the study of partial differential equations (PDEs) and sometimes evolution equations allowing mathematicians to describe the behavior of a quantity that depends on several variables [6–8, 15]. We consider the existence, uniqueness and strong stability of the solutions of the equations of linear viscoelasticity at large time. The simplest model is provided by the one-dimensional inhomogeneous case

$$(1) \quad \begin{aligned} \rho u_{tt}(t, x) &= cu_{xx}(t, x) - \int_{-\infty}^t g(t - \tau) u_{xx}(\tau, x) d\tau \\ &+ \mu_1 u_t(t, x) + \mu_2 u_t(t - s, x), \end{aligned}$$

where ρ, c are positive constants, μ_1, μ_2 are real numbers, g is independent of x , and integrable, $u(t, x)$ has been assigned to (t, x) in $[0, +\infty[\times [0, \pi]$ and $u_{xx} = \frac{\partial^2}{\partial x^2} u$.

Kirane and Houari [9] studied this equation with suitable initial boundary value conditions with $\rho = 1$ and $c = 1$. Nicaise and Pignotti [11] studied this equation with conditions $\rho = 1$, $c = 1$ and $g = 0$.

In this paper, we prove the existence and uniqueness of a solution to a system with conditions $\rho > 0$, $c > 0$ and g is a positive nonincreasing C^1 and integrable function. Moreover, the stability of the solution is shown in particular case.

✉ hinpascal51@gmail.com, ORCID: 0000-0002-7026-8630

DOI: 10.22103/jmmr.2023.19918.1303

Publisher: Shahid Bahonar University of Kerman

How to cite: G. Degla, P.H. Zinsou, *Quantitative and stability study of the evolution of a viscoelastic body*, J. Mahani Math. Res. 2023; 12(2): 443-458.



© the Authors

The mathematical model of viscoelasticity aims to take into the behaviour of those materials whose mechanical behaviour is determined not only by the present but also on its past history. Since then a wide variety of results and applications have been obtained in many different fields. In [14] synthetic tissues which mimic human bones are investigated, while in [13] cardiological tissues are considered. Besides a model of apples regarded as viscoelastic bodies is studied in [1, 3–5, 10].

The present investigation concerns viscolastic bodies and their mechanical behaviour aiming to widen the range of applicative cases the theory can be applied. This work is presented in four parts. In the first part, we have presented the mathematical model of the system describing the evolution of a body in viscoelastic motion. In the second part, we have shown the existence and uniqueness of the solution of the system through the methods of semi groups. In the third part, we have shown that the system is strongly stable in a particular case.

2. Mathematical model

Let us consider a one-dimensional homogeneous body of density $\rho > 0$, and having the constitutive equation

$$(2) \quad \sigma(t, x) = cu_x(t, x) - \int_{-\infty}^t g(t - \tau)u_x(\tau, x)d\tau,$$

where $g : \mathbb{R}_+ \rightarrow \mathbb{R}$ is a positive, nonincreasing C^1 function and integrable satisfying

$$(3) \quad a = \left(c - \int_0^{+\infty} g(s)ds \right) > 0.$$

The state of the body at the time $t \in \mathbb{R}$ is characterized by the displacement $u(t, x)$, the momentum $v(t, x)$ and the history of the displacement $w(t, s, x)$, which is defined by

$$w(t, s, x) = u(t - s, x) \quad \text{for } s \in \mathbb{R}_+.$$

The equation of motion is

$$(4) \quad \rho u_{tt}(t, x) = \sigma_x(t, x) + \mu_1 u_t(t, x) + \mu_2 u_t(t - s, x).$$

We have

$$(5) \quad \begin{aligned} \rho u_{tt}(t, x) &= cu_{xx}(t, x) - \int_{-\infty}^t g(t - \tau)u_{xx}(\tau, x)d\tau \\ &+ \mu_1 u_t(t, x) + \mu_2 u_t(t - s, x). \end{aligned}$$

We shall study the existence and uniqueness of the equation (5) firstly in the case $\mu_1 = 0 = \mu_2$ and secondly in the case $\mu_1 \neq 0$ and $\mu_2 \neq 0$. We assume that

$u(t, x) \in H_0^1(0, \pi) \cap H^2(0, \pi)$ and we set $v = \rho u_t$. Therefore

$$\begin{aligned} w_t(t, s, x) &= \lim_{h \rightarrow 0} \frac{w(t+h, s, x) - w(t, s, x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{u(t+h-s, x) - u(t-s, x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{u(t-s+h, x) - u(t-s, x)}{h}. \end{aligned}$$

$$\begin{aligned} w_s(t, s, x) &= \lim_{h \rightarrow 0} \frac{w(t, s+h, x) - w(t, s, x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{u(t-s-h, x) - u(t-s, x)}{h} \\ &= - \lim_{\lambda \rightarrow 0} \frac{u(t-s+\lambda, x) - u(t-s, x)}{\lambda} \\ &= -w_t(t, s, x). \end{aligned}$$

For $\mu_1 = 0 = \mu_2$, we obtain the system

$$(6) \quad \begin{cases} u_t = \rho^{-1}v & (t, x) \in Q_\infty, \\ v_t = cu_{xx} - \int_0^{+\infty} g(s)w_{xx}(t, s, x)ds & (t, x) \in Q_\infty, \\ w_t = -w_s & (t, s, x) \in R_\infty, \\ u(t, 0) = u(t, \pi) = 0 & t \in \mathbb{R}_+, \\ u(0, x) = u_0(x); v_0(x) = v(0, x) & x \in [0, \pi], \\ w(0, s, x) = w_0(s, x) & (s, x) \in Q_\infty, \end{cases}$$

where $Q_\infty = \mathbb{R}_+ \times [0, \pi]$ and $R_\infty = \mathbb{R}_+^2 \times [0, \pi]$.

Definition 2.1. Let g be an integrable function on the interval $[0, \pi]$.

The subspace $L_g^2(\mathbb{R}_+, H_0^1(0, \pi))$ is defined by

$$L_g^2(\mathbb{R}_+, H_0^1(0, \pi)) = \left\{ u \in L^2(\mathbb{R}_+, H_0^1(0, \pi)) \mid \int_0^\infty \int_0^\pi g(s)u_x^2(s, x)dxds < \infty \right\}.$$

The norm on L_g^2 is defined by

$$\|u\|_{L_g^2(\mathbb{R}_+, H_0^1(0, \pi))} = \left(\int_0^\infty g(s) \|u\|_{H_0^1(0, \pi)}^2 ds \right)^{\frac{1}{2}},$$

where

$$\|u\|_{H_0^1(0, \pi)}^2 = \|u_x\|_{L^2(0, \pi)}^2 \quad \text{with} \quad \|u\|_{L^2(0, \pi)} = \left(\int_0^\pi |u|^2 dx \right)^{\frac{1}{2}}.$$

$$\begin{aligned} H^m(\Omega) &= \{u \in L^2(\Omega), D^\alpha u \in L^2(\Omega), \alpha \in \mathbb{N}^n, |\alpha| \leq m\} \\ H_0^1(\Omega) &= \{u \in H^1(\Omega), u = 0 \text{ on } \partial\Omega\} \end{aligned}$$

Proposition 2.1. Let the subspaces $H_0^1(0, \pi)$, $H^2(0, \pi)$ and $L_g^2(\mathbb{R}_+; H_0^1(0, \pi))$ be defined as before. Let $L^2(0, \pi)$ be the set of square integrable functions on the interval $[0, \pi]$. Then

$$(7) \quad \begin{aligned} \langle (u, v, w), (\tilde{u}, \tilde{v}, \tilde{w}) \rangle &= \int_0^\pi \left(au_x \tilde{u}_x + \rho^{-1} v \tilde{v} \right. \\ &\quad \left. + \int_0^\infty g(s)[u_x - w_x][\tilde{u}_x - \tilde{w}_x] ds \right) dx, \end{aligned}$$

is an inner product on the subspace

$$H = \left(H_0^1(0, \pi) \cap H^2(0, \pi) \right) \times L^2(0, \pi) \times L_g^2(\mathbb{R}_+; H_0^1(0, \pi)),$$

with

$$\|(u, v, w)\|_H = (\langle (u, v, w), (u, v, w) \rangle)^{\frac{1}{2}},$$

where

$$\langle (u, v, w), (u, v, w) \rangle = \int_0^\pi \left(au_x^2 + \rho^{-1} v^2 + \int_0^\infty g(s)(u_x - w_x)^2 ds \right) dx.$$

Proof. Let $(u, v, w), (u', v', w'), (\tilde{u}, \tilde{v}, \tilde{w}) \in H$ and $\lambda \in \mathbb{R}$. From linearity property of the integral operator, we obtain

$$\begin{aligned} \langle \lambda(u, v, w) + (u', v', w'), (\tilde{u}, \tilde{v}, \tilde{w}) \rangle &= \lambda \langle (u, v, w), (\tilde{u}, \tilde{v}, \tilde{w}) \rangle \\ &\quad + \langle (u', v', w'), (\tilde{u}, \tilde{v}, \tilde{w}) \rangle . \\ \langle (u, v, w), \lambda(\tilde{u}, \tilde{v}, \tilde{w}) + (u', v', w') \rangle &= \lambda \langle (u, v, w), (\tilde{u}, \tilde{v}, \tilde{w}) \rangle \\ &\quad + \langle (u', v', w'), (\tilde{u}, \tilde{v}, \tilde{w}) \rangle . \end{aligned}$$

Then $\langle \dots \rangle$ is bilinear.

$$\begin{aligned} \langle (u, v, w), (\tilde{u}, \tilde{v}, \tilde{w}) \rangle &= \int_0^\pi \left(au_x \tilde{u}_x + \rho^{-1} v \tilde{v} \right) dx \\ &\quad + \int_0^\pi \left(\int_0^\infty g(s)[u_x - w_x][\tilde{u}_x - \tilde{w}_x] ds \right) dx. \\ &= \int_0^\pi \left(a \tilde{u}_x u_x + \rho^{-1} \tilde{v} v \right) dx \\ &\quad + \int_0^\pi \left(\int_0^\infty g(s)[\tilde{u}_x - \tilde{w}_x][u_x - w_x] ds \right) dx. \\ &= \langle (\tilde{u}, \tilde{v}, \tilde{w}), (u, v, w) \rangle . \end{aligned}$$

Then $\langle \dots \rangle$ is symmetric. We show that $\langle \dots \rangle$ is positive definite.

$$\langle (u, v, w), (u, v, w) \rangle = \int_0^\pi \left(au_x^2 + \rho^{-1} v^2 + \int_0^\infty g(s)(u_x - w_x)^2 ds \right) dx.$$

$au_x^2 \geq 0, \rho^{-1}v^2 \geq 0, \int_0^\infty g(s)(u_x - w_x)^2 ds \geq 0$, then

$$(au_x^2 + \rho^{-1}v^2 + \int_0^\infty g(s)(u_x - w_x)^2 ds) \geq 0,$$

and

$$\int_0^\pi (au_x^2 + \rho^{-1}v^2 + \int_0^\infty g(s)(u_x - w_x)^2 ds) dx \geq 0.$$

Then $\langle (u, v, w), (u, v, w) \rangle \geq 0$ and $\langle \cdot, \cdot \rangle$ is positive.

$$\begin{aligned} \langle (u, v, w), (u, v, w) \rangle = 0 &\iff \int_0^\pi (au_x^2 + \rho^{-1}v^2 \\ &\quad + \int_0^\infty g(s)(u_x - w_x)^2 ds) dx = 0. \\ &\iff \begin{cases} u_x = 0 & \forall (t, x) \in \mathbb{R}_+ \times [0, \pi], \\ v = 0 & \forall (t, x) \in \mathbb{R}_+ \times [0, \pi], \\ u_x - w_x = 0 & \forall (t, x) \in \mathbb{R}_+ \times [0, \pi]. \end{cases} \\ &\iff (u, v, w) = (0, 0, 0). \end{aligned}$$

Conclusion : $\langle \cdot, \cdot \rangle$ is an inner product on H . □

Proposition 2.2. H equipped with the inner product $\langle \cdot, \cdot \rangle$ defined by (7) is a Hilbert space.

Proof. H equipped with the inner product $\langle \cdot, \cdot \rangle$ defined by (7) is a prehilbertian space. We will show that H is complete.

Let $(u^n, v^n, w^n)_{n \geq 0}$ be a Cauchy sequence in H . Then

$$\forall \epsilon > 0, \exists n_0 : \forall n, m \geq n_0; \|(u^n, v^n, w^n) - (u^m, v^m, w^m)\|_H \leq \epsilon.$$

$$\begin{aligned} \|(u^n, v^n, w^n) - (u^m, v^m, w^m)\|_H^2 &= \|(u^n - u^m, v^n - v^m, w^n - w^m)\|_H^2 \\ &= \int_0^\pi (a(u_x^n - u_x^m)^2 + \rho^{-1}(v^n - v^m)^2) dx \\ &\quad + \int_0^\pi \left(\int_0^\infty g(s)(u_x^n - u_x^m - w_x^n + w_x^m)^2 ds \right) dx. \\ &= a \int_0^\pi (u_x^n - u_x^m)^2 dx + \rho^{-1} \int_0^\pi (v^n - v^m)^2 dx \\ &\quad + \int_0^\pi \int_0^\infty g(s)(u_x^n - u_x^m - w_x^n + w_x^m)^2 ds dx. \end{aligned}$$

Since $\|(u^n, v^n, w^n) - (u^m, v^m, w^m)\|_H \leq \epsilon$, we have

$$a \|(u_x^n - u_x^m)\|_{L^2(0, \pi)}^2 + \rho^{-1} \|v^n - v^m\|_{L^2(0, \pi)}^2 + \int_0^\pi g(s) \|w_x^n - w_x^m\|_{L^2(0, \pi)}^2 ds \leq \epsilon.$$

So

$$\begin{cases} a \| (u_x^n - u_x^m) \|_{L^2(0, \pi)}^2 \leq \epsilon, \\ \rho^{-1} \| v^n - v^m \|_{L^2(0, \pi)}^2 \leq \epsilon, \\ \int_0^\infty g(s) \| w_x^n - w_x^m \|_{L^2(0, \pi)}^2 ds \leq \epsilon, \end{cases}$$

and

$$\begin{cases} \| (u_x^n - u_x^m) \|_{L^2(0, \pi)}^2 \leq \epsilon_1 & \forall \epsilon_1 > 0, \\ \| v^n - v^m \|_{L^2(0, \pi)}^2 \leq \epsilon_2 & \forall \epsilon_2 > 0, \\ \| w^n - w^m \|_{L_g^2(\mathbb{R}_+, H_0^1(0, \pi))}^2 \leq \epsilon_3 & \forall \epsilon_3 > 0. \end{cases}$$

Then u^n , v^n and w^n are Cauchy sequences in $H_0^1(0, \pi)$, $L^2(0, \pi)$ and $L_g^2(\mathbb{R}_+, H_0^1(0, \pi))$ respectively.

u^n is a Cauchy sequence in $H_0^1(0, \pi)$ and since $H_0^1(0, \pi)$ is complete, there exists a rank n_1 from which u^n converges to an element l_1 in $H_0^1(0, \pi)$.

v^n is a Cauchy sequence in $L^2(0, \pi)$ and since $L^2(0, \pi)$ is complete, there exists a rank n_2 from which v^n converges to an element l_2 in $L^2(0, \pi)$.

$L_g^2(\mathbb{R}_+, H_0^1(0, \pi))$ is a closed subspace of $L^2(\mathbb{R}_+, H_0^1(0, \pi))$ and $L^2(\mathbb{R}_+, H_0^1(0, \pi))$ complete. Thus $L_g^2(\mathbb{R}_+, H_0^1(0, \pi))$ is complete.

w^n is a Cauchy sequence in $L_g^2(\mathbb{R}_+, H_0^1(0, \pi))$ and since $L_g^2(\mathbb{R}_+, H_0^1(0, \pi))$ is complete, there exists a rank n_3 from which w^n converges to an element l_3 in $L_g^2(\mathbb{R}_+, H_0^1(0, \pi))$.

Taking $n' = \max(n_1, n_2, n_3)$, $(u^n, v^n, w^n)_n$ converges to the triplet (l_1, l_2, l_3) belonging to H from rank n' .

Conclusion : H is a Hilbert space. \square

Let us define the operator A as follows

$$A : D(A) \subseteq H \longrightarrow H$$

$$A \begin{pmatrix} u \\ v \\ w \end{pmatrix} = \begin{pmatrix} cu_{xx}(t, x) - \int_0^\infty g(s) w_{xx}(t, s, x) ds \\ \rho^{-1} v \\ w_s \end{pmatrix},$$

in which $D(A) = \{(u, v, w) \in H; (u, v, w) \text{ satisfy (8)}\}$, and the meaning of condition (8) is that

$$(8) \quad \begin{cases} v \in H_0^1(0, \pi), w_s \in L_g^2(\mathbb{R}_+; H_0^1(0; \pi)), w(t, 0, x) = u(t, x) \\ cu_{xx}(t, x) - \int_0^\infty g(s) w_{xx}(t, s, x) ds \in L^2(0, \pi) \end{cases}.$$

At this point, let us observe that (6) can be equivalently rewritten under the abstract form

$$(9) \quad \begin{cases} Z'(t) = AZ(t), & t \geq 0, \\ Z(0) = \xi, \end{cases}$$

where $Z(t)(x) = (u(t, x), v(t, x), w(t, x))$ and $\xi = (u_0, v_0, w_0)$.

3. Main results

3.1. Existence and uniqueness of the solution of problem (9).

The existence and uniqueness of the system (9) depends on the operator A . If A is an infinitesimal generator of a C_0 -semi group of contraction, then the system admits a unique solution. In this section, we will show that $-A$ is maximal monotone which means that A is infinitesimal generator. [12]

Proposition 3.1. The operator $-A$ is maximal monotone on H .

Proof. We will show that the operator $I - A$ is surjective.

Let $(\tilde{u}, \tilde{v}, \tilde{w}) \in H_0^1(0, \pi) \times L^2(0, \pi) \times L_g^2(\mathbb{R}_+; H_0^1(0; \pi))$ and $(u, v, w) \in D(A)$.

$(I - A)(u, v, w) = (\tilde{u}, \tilde{v}, \tilde{w}) \implies$

$$(10) \quad \begin{cases} u - \rho^{-1}v = \tilde{u}, \\ v - cu_{xx} + \int_0^\infty g(s)w_{xx}ds = \tilde{v}, \\ w + w_s = \tilde{w}, \\ w(t, 0, x) = u(t, x). \end{cases}$$

$w_s + w = \tilde{w}$, then

$$(11) \quad w(s, x) = w(0, x)e^{-s} + \int_0^s e^{-(s-\tau)}\tilde{w}(\tau, x)d\tau.$$

$$(12) \quad = u(x)e^{-s} + \int_0^s e^{-(s-\tau)}\tilde{w}(\tau, x)d\tau.$$

$$(13) \quad w_s + w = \tilde{w} \implies w_x = -w_{sx} + \tilde{w}_x.$$

$$(14) \quad \implies g(s)w_x^2 = -g(s)w_xw_{sx} + g(s)w_x\tilde{w}_x.$$

$$(15) \quad \implies g(s)w_x^2 = -\frac{1}{2}g(s)[(w_x)^2]_s + g(s)w_x\tilde{w}_x.$$

$$(16) \quad \implies \int_0^\infty \int_0^\pi g(s)w_x^2 dx ds = -\frac{1}{2} \int_0^\infty \int_0^\pi g(s)(w_x^2)_s dx ds \\ + \int_0^\infty \int_0^\pi g(s)w_x\tilde{w}_x dx ds.$$

$$(17) \quad \int_0^\infty \int_0^\pi g(s)(w_x^2)_s dx ds = \int_0^\pi \lim_{X \rightarrow \infty} [g(s)w_x^2(s, x)]_0^X dx - \int_0^\infty \int_0^\pi g'(s)w_x^2 dx ds.$$

$$(18) \quad = \int_0^\pi -g(0)w_x^2(0, x)dx - \int_0^\infty \int_0^\pi g'(s)w_x^2 dx ds.$$

$$(19) \quad = -g(0) \int_0^\pi u_x^2(t, x)dx - \int_0^\infty \int_0^\pi g'(s)w_x^2 dx ds.$$

$$(20) \quad \int_0^\infty \int_0^\pi g(s)w_x^2 dx ds = \frac{1}{2}g(0) \int_0^\pi u_x^2(t, x)dx$$

$$+\frac{1}{2}\int_0^\infty\int_0^\pi g(s)w_x^2 dx ds + \int_0^\infty\int_0^\pi g(s)w_x\tilde{w} dx ds.$$

$$\forall a, b(a-b)^2 = a^2 + b^2 - 2ab \geq 0.$$

$$ab \leq \frac{1}{2}a^2 + \frac{1}{2}b^2.$$

Using this relation with $a = w_x, b = \tilde{w}_x$, we obtain

$$w_x\tilde{w}_x \leq \frac{1}{2}w_x^2 + \frac{1}{2}\tilde{w}_x^2.$$

So,

(21)

$$\int_0^\infty\int_0^\pi g(s)w_x\tilde{w}_x dx ds \leq \int_0^\infty\int_0^\pi g(s)\left(\frac{1}{2}w_x^2 + \frac{1}{2}\tilde{w}_x^2\right) dx ds.$$

(22)

$$\leq \frac{1}{2}\int_0^\infty\int_0^\pi g(s)w_x^2 dx ds + \frac{1}{2}\int_0^\infty\int_0^\pi g(s)\tilde{w}^2 dx ds.$$

(23)

$$\int_0^\infty\int_0^\pi g(s)w_x^2 dx ds \leq \frac{1}{2}g(0)\int_0^\pi u_x^2 dx + \frac{1}{2}\int_0^\infty\int_0^\pi g(s)w_x^2 dx ds$$

$$+ \frac{1}{2}\int_0^\infty\int_0^\pi g(s)\tilde{w}^2 dx ds + \frac{1}{2}\int_0^\infty\int_0^\pi g'(s)w_x^2 dx ds.$$

(24)

$$\int_0^\infty\int_0^\pi g(s)w_x^2 dx ds - \frac{1}{2}\int_0^\infty\int_0^\pi g(s)w_x^2 dx ds \leq \frac{1}{2}g(0)\int_0^\pi u_x^2 dx$$

$$+ \frac{1}{2}\int_0^\infty\int_0^\pi g(s)\tilde{w}^2 dx ds + \frac{1}{2}\int_0^\infty\int_0^\pi g'(s)w_x^2 dx ds.$$

(25)

$$\frac{1}{2}\int_0^\infty\int_0^\pi g(s)w_x^2 dx ds \leq \frac{1}{2}g(0)\int_0^\pi u_x^2 dx + \frac{1}{2}\int_0^\infty\int_0^\pi g(s)\tilde{w}^2 dx ds.$$

(26)

$$\int_0^\infty\int_0^\pi g(s)w_x^2 dx ds \leq g(0)\int_0^\pi u_x^2 dx + \int_0^\infty\int_0^\pi g(s)\tilde{w}^2 dx ds.$$

Since $\tilde{w} \in L_g^2(\mathbb{R}_+; H_0^1(0, \pi))$, we have

$$\int_0^\infty\int_0^\pi g(s)\tilde{w}^2 dx ds < \infty,$$

and so (26) shows that $\int_0^\infty\int_0^\pi g(s)w_x^2 dx ds < \infty$ and $w \in L_g^2(\mathbb{R}_+; H_0^1(0, \pi))$.

(27)

$$u - \rho^{-1}v = \tilde{u} \implies v = \rho u - \rho\tilde{u}.$$

(28)

$$v - cu_{xx} + \int_0^\infty g(s)w_{xx} ds = \tilde{v}.$$

then

(29)

$$-\rho\tilde{u} + \rho u - cu_{xx} + \int_0^\infty g(s)w_{xx} ds = \tilde{v}.$$

$$(30) \quad w = u(x)e^{-s} + \int_0^s e^{-(s-\tau)} \tilde{w}(\tau, x) d\tau.$$

Then

$$(31) \quad \begin{aligned} &\rho u - cu_{xx} + u_{xx} \int_0^\infty g(s)e^{-s} ds \\ &+ \int_0^\infty g(s)e^{-(s-\tau)} \tilde{w}_{xx}(\tau, x) d\tau ds = \tilde{v} + \rho \tilde{u}. \end{aligned}$$

$$(32) \quad \begin{aligned} \rho u - \left(c - \int_0^\infty g(s)e^{-s} ds \right) u_{xx} &= \tilde{v} + \rho \tilde{u} \\ &- \int_0^\infty g(s)e^{-(s-\tau)} \tilde{w}_{xx}(\tau, x) d\tau ds. \end{aligned}$$

We thus obtain a linear differential equation of the second order with second member. The homogeneous equation has characteristic equation

$$(33) \quad - \left(c - \int_0^\infty g(s)e^{-s} ds \right) r^2 + \rho = 0.$$

The main goal at this level is to obtain two real solutions for equation (33). So, the constant c must be greater than $\int_0^\infty g(s)e^{-s} ds$. That is to say

$$\left(c - \int_0^\infty g(s)e^{-s} ds \right) > 0.$$

Under this condition, we have

$$- \left(c - \int_0^\infty g(s)e^{-s} ds \right) < 0.$$

The characteristic equation (33) admits two real solutions. Consequently the differential equation (32) has a unique solution u . So, for any triplet $(\tilde{u}, \tilde{v}, \tilde{w}) \in H$, there exists a triplet $(u, v, w) \in D(A)$ such that $(I - A)(u, v, w) = (\tilde{u}, \tilde{v}, \tilde{w})$, that is $I - A$ is surjective and $Im(I - A) = H$.

$$(34) \quad \begin{aligned} \langle (u, v, w), (\tilde{u}, \tilde{v}, \tilde{w}) \rangle &= \int_0^\pi \left(au_x \tilde{u}_x + \rho^{-1} v \tilde{v} \right. \\ &\left. + \int_0^\infty g(s)[u_x - w_x][\tilde{u}_x - \tilde{w}_x] ds \right) dx. \end{aligned}$$

$$(35) \quad A(u, v, w) = (\rho^{-1}v, cu_{xx}(t, x) - \int_0^\infty g(s)w_{xx}(t, s, x) ds, w_s).$$

$$(36) \quad \begin{aligned} &\langle A(u, v, w), (u, v, w) \rangle = \\ &\int_0^\pi \left(a\rho^{-1}v_x u_x + \rho^{-1}v(cu_{xx}(t, x) - \int_0^\infty g(s)w_{xx}(t, s, x) ds) \right) dx \end{aligned}$$

$$\begin{aligned}
& + \int_0^\pi \left(\int_0^\infty g(s)(\rho^{-1}v_x + w_{sx})(u_x - w_x)ds \right) dx. \\
\langle A(u, v, w), (u, v, w) \rangle & = \int_0^\pi \left((c - \int_0^\infty g(s)ds)\rho^{-1}v_x u_x \right) dx \\
& + \int_0^\pi \left(\rho^{-1}v(cu_{xx}(t, x) - \int_0^\infty g(s)w_{xx}(t, s, x)ds) \right. \\
& \left. + \int_0^\infty g(s)(\rho^{-1}v_x u_x - \rho^{-1}v_x w_x + w_{sx}u_x - w_{sx}w_x)ds \right) dx.
\end{aligned}$$

$$\begin{aligned}
(37) \quad \langle A(u, v, w), (u, v, w) \rangle & = c\rho^{-1} \int_0^\pi v_x u_x dx \\
& - \int_0^\pi \int_0^\infty g(s)(\rho^{-1}v_x u_x)ds dx + c\rho^{-1} \int_0^\pi v u_{xx} dx - \rho^{-1} \int_0^\pi \int_0^\infty g(s)v w_{xx} ds dx \\
& + \rho^{-1} \int_0^\pi \int_0^\infty g(s)v_x u_x ds dx - \rho^{-1} \int_0^\pi \int_0^\infty g(s)v_x w_x ds dx \\
& + \int_0^\pi \int_0^\infty g(s)w_{sx} u_x ds dx - \int_0^\pi \int_0^\infty g(s)w_{sx} w_x ds dx.
\end{aligned}$$

$$(38) \quad \int_0^\pi v u_{xx} dx = - \int_0^\pi v_x u_x dx.$$

$$(39) \quad \int_0^\pi \int_0^\infty g(s)v w_{xx} ds dx = - \int_0^\pi \int_0^\infty g(s)v_x w_x ds dx.$$

$$\begin{aligned}
(40) \quad \int_0^\pi \int_0^\infty g(s)w_{sx} w_x ds dx & = -\frac{1}{2} \int_0^\pi g(0)u_x^2 dx \\
& - \frac{1}{2} \int_0^\pi \int_0^\infty g'(s)w_x^2 ds dx.
\end{aligned}$$

$$\begin{aligned}
(41) \quad \int_0^\pi \int_0^\infty g(s)u_x w_{sx} ds dx & = - \int_0^\pi g(0)u_x^2 dx \\
& - \int_0^\pi \int_0^\infty g'(s)u_x w_x ds dx.
\end{aligned}$$

$$\begin{aligned}
(42) \quad \langle A(u, v, w), (u, v, w) \rangle & = \frac{1}{2} \int_0^\pi \int_0^\infty g'(s)w_x^2 ds dx \\
& - \frac{1}{2} \int_0^\pi \int_0^\infty g'(s)w_x u_x ds dx - \frac{1}{2} \int_0^\pi g(0)u_x^2 dx.
\end{aligned}$$

$g(0) = - \int_0^\infty g'(s)ds$ then

$$(43) \quad \langle A(u, v, w), (u, v, w) \rangle = \frac{1}{2} \int_0^\pi \int_0^\infty g'(s)w_x^2 ds dx$$

$$\begin{aligned}
 & - \int_0^\pi \int_0^\infty g'(s)w_x u_x ds dx + \frac{1}{2} \int_0^\pi \int_0^\infty g'(s)u_x^2 dx. \\
 (44) \quad & \langle A(u, v, w), (u, v, w) \rangle = \frac{1}{2} \int_0^\pi \int_0^\infty g'(s)(w_x - u_x)^2 ds dx.
 \end{aligned}$$

The main goal at this level for equation (44), is to get $\langle A(u, v, w), (u, v, w) \rangle \leq 0$. So for that, the function g has to be nonincreasing. That is to say $g'(s) \leq 0$. Under this condition, we have $\langle A(u, v, w), (u, v, w) \rangle \leq 0$ and $\langle -A(u, v, w), (u, v, w) \rangle \geq 0$. So the operator $-A$ is monotone. The operator $-A$ is monotone and $Im(I - A) = H$ so the operator $-A$ is maximal monotone. \square

We deduce the following theorem.

Theorem 3.1. *The system 9 admits a unique solution $Z(t)(x) = T(t)\xi(x)$ for all $\xi \in D(A)$ and $x \in]0, \pi[$ where $(T(t))_{t \geq 0}$ is a C_0 -semi group of contraction on H .*

We then turn to the case of $\mu_1 \neq 0$ and $\mu_2 \neq 0$, we obtain the system

$$(45) \quad \begin{cases} Z'(t) = AZ(t) + h(t), & t \geq 0, \\ Z(0) = \xi, \end{cases}$$

with

$$h(t) = \begin{pmatrix} 0 \\ \mu_1 u_t(t, x) + \mu_2 u_t(t - s, x) \\ 0 \end{pmatrix}.$$

Since $u \in H_0^1(0, \pi) \cap H^2(0, \pi)$, we have

$$(46) \quad (\mu_1 u_t(t, x) + \mu_2 u_t(t - s, x)) \in L^1(0, T; H_0^1(0, \pi) \cap H^2(0, \pi)),$$

so $h \in L^1(0, T; H_0^1(0, \pi) \cap H^2(0, \pi))$. Since A is the infinitesimal generator and $h \in L^1(0, T; H_0^1(0, \pi) \cap H^2(0, \pi))$, we deduce the following theorem

Theorem 3.2. *The system (45) admits a unique solution*

$$(47) \quad Z(t)(x) = T(t)\xi(x) + \int_0^t T(t-s)h(s)ds,$$

for all $\xi \in D(A)$ and $x \in [0, \pi]$ where $(T(t))_{t \geq 0}$ is a C_0 -semi group of contraction on H .

3.2. Stability of the solution of the initial value problem.

The stability of the solution is shown in a particular case $\mu_1 = \mu_2 = 0$.

A C_0 -semigroup $T(t)$ on a Banach space X is called strongly stable if

$$\lim_{t \rightarrow +\infty} \|T(t)p\| = 0,$$

for every $p \in X$ [2].

For the strong stability of the system, we have shown that the mapping $\varphi : t \mapsto \|T(t)\xi(x)\|$ is nonincreasing and bounded below on $[0, +\infty[$. So its limit gives $\inf_{t \geq 0} \varphi(t)$. Under the conditions $v(0, x) = 0$ and $w(0, s, x) = ku(0, x)$ with $k = \frac{c}{\int_0^{+\infty} g(s)ds}$, the function φ is nonincreasing. Thus, for the system to be strongly stable, it is necessary that $\inf_{t \geq 0} \|T(t)\xi(x)\| = 0$.

Proposition 3.2.

$$\lim_{t \rightarrow +\infty} \|T(t)\xi(x)\| = 0,$$

for all $\xi \in H$.

Proof.

$$(48) \quad \rho u - au_{xx} = \tilde{v} + \lambda \rho \tilde{u} - \int_0^\infty g(s)e^{-(s-\tau)} \tilde{w}_{xx}(\tau, x) d\tau,$$

with $a = (c - \int_0^\infty g(s)e^{-s} ds) > 0$.

$$(49) \quad u_{xx} - \frac{\rho}{a}u = -\frac{1}{a}(\tilde{v} + \rho \tilde{u} - \int_0^\infty g(s)e^{-(s-\tau)} \tilde{w}_{xx}(\tau, x) d\tau).$$

$$(50) \quad u_{xx} - \omega^2 u = f(t, x),$$

where $\omega = \sqrt{\frac{\rho}{a}}$ and

$$(51) \quad f(t, x) = -\frac{1}{a}(\tilde{v} + \rho \tilde{u} - \int_0^\infty g(s)e^{-(s-\tau)} \tilde{w}_{xx}(\tau, x) d\tau).$$

The characteristic equation admits two real solutions ω and $-\omega$.

$$(52) \quad u(t, x) = c_1 e^{\omega x} + c_2 e^{-\omega x} + \int_0^x (c_1 e^{\omega(x-y)} + c_2 e^{-\omega(x-y)}) f(t, y) dy.$$

$u(t, 0) = 0 \implies c_1 + c_2 = 0$ then $c_1 = -c_2$.

$$(53) \quad u(t, x) = c_1 (e^{\omega x} - e^{-\omega x}) + c_1 \int_0^x (e^{\omega(x-y)} - e^{-\omega(x-y)}) f(t, y) dy.$$

$$(54) \quad u(t, x) = c_1 [e^{\omega x} - e^{-\omega x} + \int_0^x (e^{\omega(x-y)} - e^{-\omega(x-y)}) f(t, y) dy].$$

$$(55) \quad v(t, x) = \rho u_t = c_1 \frac{d}{dt} \int_0^x (e^{\omega(x-y)} - e^{-\omega(x-y)}) f(t, y) dy.$$

$$(56) \quad \begin{aligned} w(t, s, x) &= u(t - s, x), \\ &= c_1 [e^{\omega x} - e^{-\omega x} + \int_0^x (e^{\omega(x-y)} - e^{-\omega(x-y)}) f(t - s, y) dy]. \end{aligned}$$

$cs h(x) = \frac{e^x + e^{-x}}{2}$ and $sh(x) = \frac{e^x - e^{-x}}{2}$. Then

$$(57) \quad u(t, x) = 2c_1 sh(\omega x) + 2c_1 \int_0^x sh(\omega(x-y))f(t, y)dy.$$

$$(58) \quad v(t, x) = 2c_1 \frac{d}{dt} \int_0^x sh(\omega(x-y))f(t, y)dy.$$

$$(59) \quad w(t, s, x) = 2c_1 sh(\omega x) + 2c_1 \int_0^x sh(\omega(x-y))f(t-s, y)dy.$$

$$(60) \quad T(t)\xi(x) = Z(t)(x) = (u(t, x), v(t, x), w(t, x)),$$

with $\xi(x) = Z(0)(x) = (u_0(x), v_0(x), w_0(x))$.

$$(61) \quad \frac{d^+}{dt} \|T(t)\xi(x)\| = \lim_{h \rightarrow 0^+} \frac{\|T(t+h)\xi(x)\| - \|T(t)\xi(x)\|}{h}.$$

$$\forall x, y, \|x\| - \|y\| \leq \|x - y\|.$$

$$\|T(t+h)\xi(x)\| - \|T(t)\xi(x)\| \leq \|T(t+h)\xi(x) - T(t)\xi(x)\|.$$

$$\forall h > 0, \frac{\|T(t+h)\xi(x)\| - \|T(t)\xi(x)\|}{h} \leq \frac{\|T(t+h)\xi(x) - T(t)\xi(x)\|}{h}.$$

$$\forall h > 0, \frac{\|T(t+h)\xi(x)\| - \|T(t)\xi(x)\|}{h} \leq \frac{\|T(t) \cdot T(h)\xi(x) - T(t)\xi(x)\|}{h}.$$

$$\forall h > 0, \frac{\|T(t+h)\xi(x)\| - \|T(t)\xi(x)\|}{h} \leq \|T(t)\| \left\| \left(\frac{T(h) - I}{h} \right) \xi(x) \right\|.$$

$$(62) \quad \lim_{h \rightarrow 0^+} \left\| \left(\frac{T(h) - I}{h} \right) \xi(x) \right\| = \|A\xi(x)\|.$$

$$(63) \quad \frac{d^+}{dt} \|T(t)\xi(x)\| \leq \|A\xi(x)\|,$$

since $(T(t))_{t \geq 0}$ is a C_0 -semi group of contraction.

$$(64) \quad A\xi(x) = \frac{d}{dt} \Big|_{t=0} Z(t)(x) = \begin{pmatrix} \rho^{-1}v(0, x) \\ cu_{xx}(0, x) - \int_0^\infty g(s)w_{xx}(0, s, x)ds \\ w_s(0, s, x) \end{pmatrix}.$$

We have

$$(65) \quad w_s(0, s, x) = -u_t(0, x) = \rho^{-1}v(0, x).$$

$v(0, x) = 0$, then $w_s(0, s, x) = 0$ and so $w(0, s, x)$ is constant with respect to s .

$$(66) \quad \int_0^\infty g(s)w_{xx}(0, s, x)ds = w_{xx}(0, s, x) \int_0^\infty g(s)ds.$$

$w(0, s, x) = ku(0, x)$, then $w_{xx}(0, s, x) = ku_{xx}(0; x)$. So

(67)

$$cu_{xx}(0, x) - \int_0^\infty g(s)w_{xx}(0, s, x)ds = cu_{xx}(0, x) - w_{xx}(0, s, x) \int_0^\infty g(s)ds.$$

(68)

$$= cu_{xx}(0, x) - ku_{xx}(0, x) \int_0^\infty g(s)ds.$$

(69)

$$= cu_{xx}(0, x) - cu_{xx}(0, x).$$

(70)

$$= 0.$$

With $k = \frac{c}{\int_0^{+\infty} g(s)ds}$. We have

$$(71) \quad cu_{xx}(0, x) - \int_0^\infty g(s)w_{xx}(0, s, x)ds = 0,$$

and

$$v(0, x) = w_s(0, s, x) = 0,$$

then $A\xi(x) = 0$ and $\|A\xi\| = 0$.

So,

$$(72) \quad \frac{d^+}{dt} \|T(t)\xi(x)\| \leq 0.$$

Let $\varphi(t) = \|T(t)\xi(x)\|$. Then φ is nonincreasing and bounded below so,

$$(73) \quad \lim_{t \rightarrow +\infty} \varphi(t) = \inf_{t \geq 0} \varphi(t) = 0,$$

and $(T(t))_{t \geq 0}$ is strongly stable. \square

4. Conclusion

We have proved in our work, the existence, uniqueness and stability of the solution to integro differential system (6) that is motivated evolution linear viscoelasticity body. Our results coincide with conditions $\rho = 1$, $c = 1$ of Kirane and Houari.

5. Author Contributions

The results of this research are obtained with the strong contribution of all authors.

6. Acknowledgement

We would like to thank the anonymous reviewers for their thoughtful comments and efforts towards improving our manuscript.

7. Funding

The authors would like to thank African Center of Excellence in Mathematical Sciences, IT and Applications (CEA-SMIA).

8. Conflict of interest

The authors certify that they have no affiliations with or involvement in any organization or entity with any financial interest (such as honoraria; educational grants; participation in speakers' bureaus; membership, employment, consultancies, stock ownership, or other equity interest, and expert testimony or patent-licensing arrangements), or non-financial interest (such as personal or professional relationship, affiliations, knowledge or beliefs) in the subject matter or materials discussed in this manuscript.

References

- [1] Amendola G., S. Carillo, J.M. Golden and A. Manes, Viscoelastic fluids : free energies, differential problems and asymptotic behaviour, *Discrete and Continuous Dynamical Systems- Series B*, vol. 19, pp.1815-1835, 2014.
- [2] Brezis H. *Analyse fonctionnelle*, Masson, 1987
- [3] Carillo S. , Valente V. and G. Vergara Caffarelli, A result of existence and uniqueness for an integro-differential system in magneto-viscoelasticity, *Appllicable Analisis*, vol. 90, pp. 1791-1802, 2010.
- [4] Carillo S. , V. Valente and G. Caffarelli, A linear viscoelasticity problem with a singular memory kernel : an existence and uniqueness result, *Differential and integral Equations*, vol. 26, pp. 1115-1125, 2013.
- [5] Carillo S. A 3-dimensional singular kernel problem in viscoelasticity : an existence result, *Atti della Accademia Peloritana dei Pericolanti, Classe di scienze Fisiche, Matematiche Naturali*, vol. 97(SI), 13 pp, 2019.
- [6] Carillo S. , M. Chipot, Valente, and G. Caffarelli, A magneto-viscoelasticity problem with a singular memory kernel, *Nonlinear Analysis Series B: Real world Applications*, vol. 35C, pp. 200-210, 2017.
- [7] Chipot M. , I. Shafir, V. Valente, and G. Caffarelli, A nonlocal problem arising in the study of magneto-elastic interactions, *Boll UMI Serie IX, I*, pp. 197-222, 2008.
- [8] Chipot M. , I. Shafir, V. Valente, and G. Caffarelli, One a hyperbolic-parabolic system arising in magnetoelasticity, *J. Math. Anal. App*, vol. 352, pp. 120-131, 2009.
- [9] Kirane M. and B.S. Houari, Existence and asymptotic stability of a viscoelastic wave equation with a delay, *Z. Angew. Math. Phys*, 62 (2011), 1065-1082.
- [10] Mahiuddin Md. , Md. Imran H. Khan, Nghia Duc Pham and M.A. Karim, Developpment of fractional viscoelastic model for characterising viscoelastic properties of food material during drying, *Food Bio-science*, vol. 23, pp. 45-53, 2018.
- [11] Nicaise S. and Pignotti C., Stability and instability results of the wave equation with a delay term in the boundary or internal feedbacks, *SIAM J. Control Optim*, 45(5) (2006), 1561-1585.
- [12] PAZY A., *Semi group of linear operator and applications to partial differential equations*.
- [13] Rassoli A. , N. Fatourae and R. Guidoin, Structural model for viscoelastic properties of péricardial bioprosthetic valves, *Artificial Organs*, vol. 42, pp. 630-639, 2018.
- [14] Shahin-Shamsabadi A. and al, Mechanical, material, and biological study of a PCL/ bioactive glass bone scaffold: Importance of viscoelasticity, *Materials Science and Engineering: C*, vol. 90, pp. 280-288, 2018.
- [15] Valente V. and G. Vergara Caffarelli, On the dynamics of magneto-elastic interactions: existence of solutions and limit behavior, *Asymptotic Analysis*, vol. 51, pp. 319-333, 2007.

GUY DEGLA

ORCID NUMBER: 0000-0003-1162-6140

INSTITUT DE MATHÉMATIQUES ET DE SCIENCES PHYSIQUES

PORTO-NOVO, BÉNIN

Email address: gdegla@gmail.com

PASCAL HINNILO ZINSOU

ORCID NUMBER: 0000-0002-7026-8630

INSTITUT DE MATHÉMATIQUES ET DE SCIENCES PHYSIQUES

PORTO-NOVO, BÉNIN

Email address: hinpascal51@gmail.com