

Full Length Research Paper

Repellent and insecticidal effects of essential oils of *Petroselinum crispum* (Mill.) Fuss and *Pimenta racemosa* var. *racemosa* (Mill.) J.W. Moore leaves on *Dinoderus porcellus* Lesne (Coleoptera: Bostrichidae)

Loko Yêyinou Laura Estelle^{1*}, Bogninou Gbêdossou Sophie Reine², Toffa Joelle¹, Kassa Parfait¹, Douro Kpindou Ouorou Kobi³ and Chougourou C. Daniel⁴

¹ Laboratory of Applied Entomology (LEnA), Ecole Nationale Supérieure des Biosciences et Biotechnologies Appliquées (ENSBB), National University of Sciences, Technologies, Engineering and Mathematics, Abomey, BP 14 Dassa, Benin.

² Laboratoire d'Etude et de Recherche en Chimie Appliquée (LERCA), EPAC/UAC, 01 BP 2009 Cotonou, Bénin.

³ International Institute of Tropical Agriculture, 08 BP 0932 Cotonou, Benin.

⁴ Laboratory of Research in Applied Biology, Department of Environment Engineering, EPAC/UAC, 01 BP 2009 Cotonou, Benin.

Received 3rd September, 2020; Accepted 17th May, 2021

Essential oils from leaves of *Petroselinum crispum* (Mill.) Fuss (Apiaceae) and *Pimenta racemosa* var. *racemosa* (Mill.) J.W. Moore (Myrtaceae) were evaluated for their repellent properties, insecticidal and fumigant activities against *Dinoderus porcellus* Lesne (Coleoptera: Bostrichidae). The essential oils of these plants were obtained by Clevenger-type hydro distillation method with yield of 0.47 and 1.09% w/w for *P. crispum* and *P. racemosa*, respectively. Essential oils of both plants exhibited a low repellent activity (class I and II of repellency) against *D. porcellus* at all tested concentrations. The *P. crispum* and *P. racemosa* essential oils were toxic to *D. porcellus* in contact, ingestion and fumigant assays. The *D. porcellus* adults were more susceptible to contact action of *P. crispum* ($LD_{50} = 1.15 \mu\text{L/adult}$) than *P. racemosa* ($LD_{50} = 414.38 \mu\text{L/adult}$) at 72 h, whereas in ingestion assay, *P. crispum* essential oil was the most effective toxin ($LC_{50} = 3.88 \mu\text{L/g}$) at 21 days interval time. Strong feeding deterrence (68.97%) was achieved in *D. porcellus* adults by using *P. crispum* essential oil at a concentration of 0.4 $\mu\text{L/g}$ of yam chips. In the fumigation assays, *P. crispum* ($LC_{50} = 7.22 \mu\text{L/L air}$) essential oil was more toxic than *P. racemosa* ($LC_{50} = 3.39 \times 10^6 \mu\text{L/L air}$) against *D. porcellus* adults within 7 days. These findings, suggest that *P. crispum* essential oil was more active against *D. porcellus* adults and showed its potential for development as natural anti-feeding agent, and fumigant insecticide for managing *D. porcellus* adults in stored yam chips.

Key words: Contact toxicity, feeding deterrence, fumigant toxicity, repellent activity, storage insects.

INTRODUCTION

Yam (*Dioscorea* species, Dioscoreaceae) is largely cultivated in West Africa for their starchy tubers, which contribute to food security in this region (Kiba et al., 2020). In Benin, fourth worldwide producer, yam contributes to 441 kcal/capita/day (FAO, 2018). However,

yam tubers have high water content, which makes them highly perishable causing severe losses which can reach 85% of stocks (Umogbai, 2013). Traditionally, yam tubers are processed and dehydrated on sun to obtain dried chips whose flour are cooked to obtain a very nourishing

dough (Omohimi et al., 2019). In northern Benin, yam chip flour is also used to prepare a type of couscous namely wassa-wassa, which is recommended to diabetic patients (Behanzin et al., 2018).

Unfortunately, insects attacked yam chips, cause important losses, reduce their market value and the quality of paste obtained with infested yam chips (Omohimi et al., 2019). Among these pests, *Dinoderus porcellus* Lesne (Coleoptera: Bostrichidae) is far the most abundant and damageable pest in stored yam chips (Loko et al., 2013). Currently, control of *D. porcellus* population is done by some farmers with synthetic insecticides (Loko et al., 2013) which lead to death of some persons (Adedoyin et al., 2008; Adeleke, 2009).

It is so urgent to find alternatives to synthetic insecticides for protection of stored yam chips. Potential alternatives to synthetic insecticides are plant-derived insecticides, which are less persistent in the environment, and often less toxic to mammalian (Smith and Thomas, 2020).

Essential oils exhibit various insecticidal and repellent properties. Due to their diverse biologically active compounds, the essential oils of parsley *Petroselinum crispum* (Mill.) Fuss and bay rum tree, *Pimenta racemosa* var. *racemosa* (Mill.) J.W. Moore have been suggested as suitable alternatives for controlling storage insect pests. Indeed, it is known from previous works that essential oils of *P. crispum* leaves have strong adverse effects on *Callosobruchus maculatus* (F.) (Massango et al., 2017), *Ephestia kuehniella* Zeller (Lepidoptera: Pyralidae), and *Plodia interpunctella* Hübner (Lepidoptera: Pyralidae) (Maroufpoor et al., 2016) as a toxic fumigant and reproductive inhibitor. Similarly, *P. racemosa* essential oils show a high toxicity against *Prostephanus truncatus* (Horn) (Noudogbessi et al., 2008) and *Tribolium castaneum* (Herbst) (Lee et al., 2002). However, the biological activities of essential oils of leaves of these plants have not been investigated yet on *D. porcellus*.

The composition of essential oils of parsley and bayrum tree leaves from different origins has been studied by several authors. In general, the essential oils of parsley leaves have β -phellandrene, and 1,3,8-p-menthatriene as the main constituents (Petropoulos et al., 2004; Mulugeta et al., 2015; El-Zaeddi et al., 2016).

While essential oils of *P. racemosa* leaves composed of about twenty components with eugenol, myrcene, and chavicol as major components (Noudogbessi et al., 2008; Contreras-Moreno et al., 2016). The objective of this study was to test the possible repellent and insecticidal properties of essential oils of *P. crispum* and *P. racemosa* leaves against *D. porcellus*.

MATERIALS AND METHODS

Plant and extraction of essential oils

The fresh leaves of *P. crispum* and *P. racemosa* were collected in the town of Abomey-Calavi. The extraction was carried out using a Clevenger type apparatus at the Laboratory of Study and Research in Applied Chemistry (LERCA) of the Polytechnic School of Abomey-Calavi (EPAC) according to Saroukolai et al. (2014). Fresh leaves were subjected to hydrodistillation. Anhydrous sodium sulphate was used to remove water after extraction. Extracted oil was stored in airtight containers in a refrigerator at 4°C until experiment. The yield of essential oils (RD), defined as the ratio between the mass of essential oil obtained after the extraction (M') and the fresh or dry mass of the plant material used (M), was calculated according to the formula:

$$RD(\%) = \frac{M'}{M} \times 100$$

Insect culture

The adult of *D. porcellus* were collected from infested yam chips purchased at the Kpota market in the town of Abomey-Calavi. The rearing of *D. porcellus* was done according to the methodology described by Loko et al. (2020). Five hundred grams of healthy yam chips were infested with 50 adults of *D. porcellus* and maintained in the laboratory (temperature: $25 \pm 2^\circ\text{C}$, RH: $70 \pm 5\%$, and 12L/12D) (Oni and Omoniyi, 2012). F1 progeny were used for all experiments.

Repellence bioassay

The method described by Babarinde et al. (2008) was used for testing repellence of essential oils of *P. crispum* and *P. racemosa* leaves against *D. porcellus*. The experimental design consisted of Petri dishes (9 cm diameter) containing two-half filter paper of Whatman No. 1. Four doses (5, 10, 15, and 20 μL of stock solution dissolved in 0.2 mL of ethanol) of each essential oil was applied to a half paper disc (30 cm^2). Each untreated half paper was joined to a treated using clear adhesive tape and place in Petri dish. The control experiment was to treat one paper half with pure ethanol (0.2 mL). After waiting for 10 min for the solvent to evaporate, 20 unsexed adults of *D. porcellus* (3-7 days old) were deposited in the centre of each Petri dish. The covered Petri dishes were placed in a dark room in the laboratory. The number of *D. porcellus* present on the control (C) and treated (T) filter papers was recorded after 30 min, 2, 4 and 8 h of exposure (Guruprasad and Akmal, 2014). Each treatment was replicated three times. The repulsion percentage (RP) was calculated using the formula:

$$RP = \left[\frac{(C - T)}{(C + T)} \right] \times 100$$

The mean repulsion percentage of each treatment was assigned to the repellent classes: class 0 (RP $\leq 0.1\%$), class I (0.1 - 20%), class II (20.1 - 40%), Class III (40.1 - 60%), Class IV (60.1 - 80%) and Class V (80.1 - 100%). The bioactivity of each dose of essential oil

*Corresponding author. E-mail: lokoestelle@yahoo.fr.

was classified (values < 1 repellency; 1 neutral; > 1 attractant) according to the repellence indices (RI) calculated following the formula:

$$RI = 2PT / PT + PA$$

where PT = percentage of insects attracted by treated paper with essential oil and PA = percentage of insects attracted by paper treated with solvent.

Contact toxicity by topical application

The contact toxicity of each essential oil was evaluated following the methodology described by Mukanga et al. (2010). Using a microsyringe, 1 µL of the essential oils of leaves of each plant was applied at concentrations 0.1, 0.25, 0.5 and 1% (v/v) on the dorsal side of the thorax of 20 newly emerged adults (3-7 days old) *D. porcellus*. The treated insects were transferred individually in a plastic box containing 10 g of untreated yam chips. Solvent extracts were applied to control insects and each treatment was replicated four times. Insects were examined after 24, 48 and 72 h and the number of dead insects was recorded (Caballero-Gallardo et al., 2012).

The percentage of adult mortality was calculated and corrected according to Schneider-Orelli's formula (Püntener, 1981):

$$\text{Corrected mortality} = \frac{\% MT - \% MC}{100 - \% MC} \times 100 \quad (1)$$

where MT = mortality in the treatment, MC = mortality in the control.

Ingestion toxicity and feeding deterrence

To evaluate the ingestion toxicity of each essential oil the methodology described by Loko et al. (2020) was used. For that, 10 g of yam chips were placed in plastic boxes (7 cm × 3.7 cm) and then treated with an acetone solution containing essential oils of *P. crispum* and *P. racemosa* leaves at different doses. The doses were prepared by diluting each time with 1 mL of acetone the respective volumes of 2, 4, 6, 8, and 10 µL of essential oils. They were then dispersed homogeneously in the yam chips using a micropipette. Ten pairs of *D. porcellus* adults (3-7 days old) were introduced in plastic boxes containing treated yam chips. The tests were repeated 4 times for each essential oil, each dose and each control. The counts of dead insects were carried out at 1, 3, 5, 7, 14 and 21 days after exposure (Othira et al., 2009). The percentage of insect mortality was calculated using Schneider-Orelli's formula (Equation 1). The percentage of weight loss was calculated according to the formula:

$$\text{Weight loss (\%)} = \frac{\text{initial weight} - \text{final weight}}{\text{initial weight}} \times 100 \quad (2)$$

The feeding deterrence index (FDI) was calculated with FDI positive value corresponding to feeding deterrent effect and FDI negative value a feeding stimulant effect (Stefanazzi et al., 2011):

$$FDI(\%) = \frac{CC - CT}{CC} \times 100 \quad (3)$$

where CC = Consumption of control yam chips, CT = Consumption

of treated yam chips.

Fumigant toxicity

The fumigant activity of essential oils of *P. crispum* and *P. racemosa* leaves against *D. porcellus* was evaluated using the methodology described by Liu and Ho (1999). 25 ml of each concentration (0, 5, 10, 20 and 40 µL dissolved in acetone) of essential oil were applied on filter paper Whatman No. 1 (2 cm diameter) and placed on the underside of the screw cap of a glass vial (2.5 cm × 5.5 cm). After waiting 1 min for the solvent to evaporate, 10 adults of *D. porcellus* were introduced into the glass vial, which was subsequently closed tightly with the cap. Glass vials were placed in a dark room in the laboratory at ambient conditions. Each treatment was replicated four times and acetone was used as control. Insect mortality was observed daily for a week (Chu et al., 2012). The insect mortality and the weight loss were calculated using respectively Equations 1 and 2.

Data analysis

The normality of mortality and repellence percentage data were tested before subjected to analysis of variance (ANOVA) or general linear model (GLM) using IBM SPSS Statistics version 25 software package. Data not following a normal distribution and not having homogeneity of variances were arcsine-transformed. Student Newman Keuls test was used to identify difference between treatments. Probit regression analysis was done on log-transformed data of various concentration-response experiments using Mathematica software version 11.3.0.0 (Wolfram Research Inc., Champaign, IL, USA). Lethal dose or concentration that kills or repels 50% of exposed insects (LD₅₀, LC₅₀ or RD₅₀) was obtained from derived regression equations.

RESULTS

Repellent activity

The results showed that *P. crispum* and *P. racemosa* essential oils acted as a repellent to *D. porcellus* adults, even at low concentrations (5 µL/cm²) (Table 1). However, the repellence of both essential oils was not significantly different ($P \geq 0.05$) at all corresponding doses. Repellent activity of *P. crispum* and *P. racemosa* essential oils on *D. porcellus* adults were not dose-dependent ($P \geq 0.05$). In addition, there was no significant ($P \geq 0.05$) interaction between oil concentration and exposure time on the repellence of both essential oils. After 30 min of exposure time, the essential oil of *P. crispum* at 5 and 10 µL/cm², showed a decrease of repellent effect with an exposure time increase (Table 1). However, with the high concentrations (15 and 20 µL/cm²) of *P. crispum* essential oil, repellent effect had increased with exposure time. In general, the repellence of *P. racemosa* essential oil against *D. porcellus* decreased with increase of concentration. Results showed that *P. racemosa* essential oil possessed high repellent activity (54.6% after an 8-h exposure) against *D. porcellus* adults only at the lowest concentration of 5 µL/cm². Essential oils of *P. crispum* and *P. racemosa* at

Table 1. Repellence of *P. racemosa* and *P. crispum* essential oils on *D. porcellus*.

Plant	Concentrations ($\mu\text{L}/\text{cm}^2$)	Repellence (%) after				Mean	Class	Repellence index	Classification
		30 min	2 h	4 h	8 h				
<i>P. crispum</i>	5	36.7 ± 40.9^a	25.6 ± 22.5^a	21.6 ± 39.5^a	29.5 ± 11.8^a	28.3 ± 13.7^a	II	0.7 ± 0.1	Repellent
	10	22.4 ± 17.2^a	12.3 ± 11.7^a	-7.8 ± 11.1^a	19.6 ± 16.9^a	11.7 ± 7.1^a	I	0.8 ± 0.0	Repellent
	15	17.1 ± 21.2^a	38.9 ± 14.5^a	37.8 ± 29.6^a	53.5 ± 15.1^a	36.8 ± 9.8^a	II	0.6 ± 0.1	Repellent
	20	14.7 ± 20.4^a	30.3 ± 15.1^a	33.3 ± 33.9^a	43.7 ± 19.7^a	30.5 ± 10.7^a	II	0.7 ± 0.1	Repellent
<i>P. racemosa</i>	5	29.8 ± 10.6^a	42.6 ± 22.8^a	31.1 ± 28.9^a	54.6 ± 4.6^a	39.5 ± 8.8^a	II	0.6 ± 0.1	Repellent
	10	23.2 ± 10.6^a	10.4 ± 12.8^a	9.3 ± 32.9^a	27.9 ± 18.9^a	17.7 ± 9.1^a	I	0.8 ± 0.1	Repellent
	15	4.6 ± 9.5^a	12.5 ± 6.7^a	22.9 ± 23.3^a	9.4 ± 12.1^a	12.3 ± 6.4^a	I	0.8 ± 0.0	Repellent
	20	3.7 ± 3.7^a	20.7 ± 27.1^a	16.7 ± 38.4^a	26.9 ± 17.3^a	17.0 ± 11.3^a	I	0.8 ± 0.1	Repellent

There is no significant difference between the values in a column followed by the same letter ($p < 0.05$).

Table 2. Dose of *P. racemosa* and *P. crispum* essential oils that repel 50% of exposed insects.

Essential oil	Exposure time	RD ₅₀ ($\mu\text{L}/\text{cm}^2$)	Slope ($\pm$ S.E.)	Intercept	χ^2	P-value
<i>P. crispum</i>	30 min	2.39	-0.583 ± 0.35	5.219	7.68	0.768
	2 h	26.37	0.782 ± 0.25	3.871	1.43	0.013
	4 h	9.91	1.159 ± 0.17	3.847	8.54	0.000
	8 h	10.19	1.000 ± 0.20	3.980	2.22	0.047
<i>P. racemosa</i>	30 min	4.77	-2.405 ± 0.09	6.623	9.03	0.287
	2 h	5.62	-1.546 ± 0.13	6.155	6.49	0.138
	4 h	6.41	-1.515 ± 0.14	6.215	6.81	0.178
	8 h	8.44	-2.294 ± 0.09	7.095	6.40	0.007

CI: Confidence interval, S.E.: standard error, χ^2 : Chi-square.

various concentrations ranged in two class of repellency (class I and II). The repellent action (RD₅₀) of *P. crispum* and *P. racemosa* essential oils on *D. porcellus* in function of exposure time is shown in Table 2. The RD₅₀ value indicates that essential oil of *P. crispum* at 30 min interval time was found to be significantly more repellent than that of *P. racemosa* with a RD₅₀ value of 2.39

$\mu\text{L}/\text{cm}^2$. However, after 2, 4 and 8 h interval times, the RD₅₀ values of *P. racemosa* essential oil were lower than those of *P. crispum* (Table 2).

Contact toxicity

Essential oils of *P. crispum* and *P. racemosa* at diverse concentrations had a topical insecticidal

effect on *D. porcellus* (Table 3). The effect concentration of *P. crispum* and *P. racemosa* essential oils on the mortality of *D. porcellus* was not significant ($P \geq 0.05$). However, contrary to *P. crispum*, the time of essential oil *P. racemosa* exposure had significant ($P \leq 0.001$) effect on the mortality of *D. porcellus*. The interaction between oil concentration of both plants and exposure time

Table 3. Mortality of *D. porcellus* after topical application of *P. crispum* and *P. racemosa* essential oils and yam chips weight loss after 3 days of exposure.

Plant	Concentration (% v/v)	Corrected mortality (%)			Weight loss (%)
		24 h	48 h	72 h	
<i>P. crispum</i>	0.1	0.20 ± 3.63 ^a	6.68 ± 4.10 ^a	21.30 ± 1.87 ^a	16.08 ± 3.40 ^a
	0.25	5.00 ± 3.54 ^a	7.40 ± 7.58 ^a	15.08 ± 5.31 ^a	15.35 ± 2.70 ^a
	0.5	4.99 ± 3.73 ^a	14.72 ± 4.46 ^a	19.87 ± 5.21 ^a	22.48 ± 5.11 ^a
	1	7.70 ± 4.29 ^a	17.24 ± 6.06 ^a	19.59 ± 6.60 ^a	26.45 ± 5.58 ^a
<i>P. racemosa</i>	0.1	6.32 ± 1.23 ^a	0.00 ± 3.13 ^a	7.57 ± 5.94 ^a	18.83 ± 2.27 ^a
	0.25	7.70 ± 2.70 ^a	0.00 ± 2.30 ^a	0.13 ± 3.60 ^a	23.98 ± 2.78 ^a
	0.5	10.07 ± 3.51 ^a	5.45 ± 0.29 ^a	3.62 ± 7.23 ^a	14.20 ± 2.41 ^a
	1	5.06 ± 0.06 ^a	5.28 ± 3.06 ^a	6.32 ± 5.19 ^a	26.03 ± 4.70 ^a
Control	0	2.50 ± 1.34	7.50 ± 2.67	7.50 ± 2.50	40.99 ± 2.07 ^b

There is no significant difference between the values in a column followed by the same letter ($p < 0.05$).

Table 4. Contact lethal dose of *P. crispum* and *P. racemosa* essential oils that kills 50% of exposed *D. porcellus*.

Essential oil	Exposure time (h)	LD ₅₀ (μL/adult)	Slope (± S.E.)	Intercept	χ ²	P-value
<i>P. crispum</i>	24	440.23	0.407 ± 0.66	3.923	7.50	0.10
	48	2.32	0.191 ± 0.26	4.703	9.21	0.83
	72	1.15	0.051 ± 1.03	4.988	1.58	0.66
	24	63.33	0.415 ± 0.97	4.252	9.72	0.98
<i>P. racemosa</i>	48	196.57	0.457 ± 0.58	3.951	1.00	0.00
	72	414.38	0.459 ± 0.63	3.788	8.99	0.02

CI: Confidence interval, S.E.: Standard error; χ²: Chi-square.

significant ($P \geq 0.05$). The mortality of *D. porcellus* did not vary significantly ($P \geq 0.05$) according to the diverse concentrations of both essential oils tested at 24, 48 and 72 h of exposure time. After 72 h of exposure, *P. crispum* essential oil at 0.1% of concentration caused the higher mortality (21.30 ± 1.87%) of *D. porcellus* (Table 3). While, only after 24 h of exposure, *P. racemosa* essential oil at 0.5% of concentration caused the higher mortality (10.07 ± 3.51) of *D. porcellus*. At 24 h after exposure, *P. racemosa* essential oil was more toxic (LD₅₀ = 63.33 μL/adult) than *P. crispum* oil (LD₅₀ = 440.23 μL/adult) (Table 4). However, at 48 and 72 h interval time, a contrary trend was observed. The topical application of essential oils of *P. crispum* and *P. racemosa* at different concentrations significantly decreased consumption of yam chips ($F = 7.125$; $df = 8$; $P \leq 0.001$) by *D. porcellus* (Table 3).

Ingestion toxicity and feeding deterrent activity

The ingestion of treated yam chips with both essential

oils at all concentrations caused a great mortality of *D. porcellus* (Figure 1). The effect of oil concentration of both plants ($P \leq 0.05$) and exposure time ($P < 0.001$) were significant on *D. porcellus* mortality. However, their interaction was not significant ($P \geq 0.05$). The essential oil of *P. crispum* at concentration above 0.2 μL/g, caused high mortalities of *D. porcellus* population with concentrations 0.8 and 1 μL/g causing a corrected mortality rate of 80.82% (Figure 1a). On the other hand, the results obtained with *P. racemosa* essential oil showed that the mortality induced by the highest concentration (1 μL/g) and the lowest concentration (0.2 μL/g) fell respectively in the 7 and 14th experimental day, while mortality increased progressively with mean concentrations (0.4, 0.6, and 0.8 μL/g) (Figure 1b). At 1 day interval time *P. racemosa* essential oil (LC₅₀ = 0.001 μL/g) caused the higher feeding toxicity of *D. porcellus* than *P. crispum* oil (LC₅₀ = 0.002 μL/g). However, at 21 days interval time, a contrary trend was observed (Table 5). The two essential oils significantly protected ($F = 2.680$, $df = 45$, $P \leq 0.05$) the yam chips compared to the control (Figure 2). Essential oil of *P. crispum* at 4 μL/mL




Figure 1. Ingestion toxicity of (a) *P. crispum* and (b) *P. racemosa* essential oils on *D. porcellus*.

Table 5. Toxicity by ingestion of *P. crispum* and *P. racemosa* essential oils on *D. porcellus*.

Essential oil	Exposure time (days)	LC ₅₀ (µL/g)	Slope (± S.E.)	Intercept	χ^2	P-value
<i>P. crispum</i>	1	0.002	-0.574 ± 0.73	3.315	4.38	0.077
	3	18.35	3.388 ± 0.08	0.719	2.73	0.434
	5	178.81	1.051 ± 0.33	2.364	2.02	0.000
	7	312.50	0.746 ± 0.33	3.131	9.21	0.957
	14	33.10	0.753 ± 0.25	3.857	8.84	0.901
	21	3.88	2.325 ± 0.42	1.261	1.30	0.729
<i>P. racemosa</i>	1	0.001	-0.286 ± 1.05	3.547	3.20	0.792
	3	218.79	1.027 ± 0.33	2.408	1.06	0.001
	5	2.96	1.982 ± 0.25	3.952	1.56	0.012
	7	31.28	1.869 ± 0.13	2.205	9.98	0.000
	14	1.41×10 ⁵	0.338 ± 0.84	3.182	3.60	0.002
	21	1.89×10 ⁶¹	-0.030 ± 11.85	3.116	0.01	0.032

weight loss (Figure 3).

Fumigant toxicity

The fumigant activity of *P. crispum* essential oil on *D. porcellus* was significantly ($P \leq 0.001$) higher than those

of *P. racemosa* oil. The increase of oil concentration of both plants and exposure time had a significant ($P \leq 0.05$) effect on the mortality of *D. porcellus*. After 7 days of exposure time, the mortality of *D. porcellus* reached 47.32% when exposed to *P. crispum* essential oil at 20 µL/L air (Figure 4). The essential oil of *P. racemosa* at all concentrations and various periods showed weak




Figure 2. Weight loss caused by *D. porcellus* on yam chips treated with *P. crispum* and *P. racemosa* essential oils at 21 days after exposure.




Figure 3. Feeding deterrence indices of treated yam chips with *P. crispum* and *P. racemosa* essential oils on *D. porcellus* after 21 days of exposure.

fumigant toxicity with higher mortality of *D. porcellus* (28.96%) at the higher oil concentration (160 μL/L air) 7 days after exposure. *P. crispum* essential oil had very lower LC₅₀ values than those of *P. racemosa* (Table 6). *D. porcellus* adults fumigated with 160 μL/L air concentration of the *P. crispum* essential oil caused significant less damage ($F = 1.94$; $df = 8$; $P \leq 0.05$) to yam chips than control at 7 days interval time (Figure 5).

DISCUSSION

The yields obtained in *P. racemosa* and *P. crispum* leaves essential oils are similar to those obtained by Alitonou et al. (2012), Craft and Setzer (2017), which are in the order of 0.9 to 2.4% and 0.2 to 0.6%, respectively. Alitonou et al. (2012) revealed that chemical composition of *P. racemosa* leaves essential oils from Benin contains




Figure 4. Mortality of *D. porcellus* caused by *P. crispum* and *P. racemosa* essential oil fumigants.

24 compounds with eugenol and myrcene as the main important. While, composition of *P. crispum* leaves essential oil cultivated in Africa was revealed by some studies with myristicin and apiol as the main compounds (Nawel et al., 2014; Snoussi et al., 2016; Agyare et al., 2017).

The investigation of bioactivities of *P. crispum* and *P. racemosa* essential oils against *D. porcellus* showed their repellent activity. The repellent activity of *P. crispum* and *P. racemosa* oils could be due to the presence of major


active compounds such as myristicin and eugenol, respectively. Indeed, You et al. (2015) have demonstrated that myristicin and eugenol have strong repellent effect respectively to the stored product insects *Liposcelis bostrychophila* Badonnel and *C. maculatus*. Moreover, myristicin and eugenol were strongly repellent against *Lasioderma serricorne* (Fabricius) (Du et al., 2014) and *Dinoderus bifoveatus* Wollaston (Ojimelukwe and Adler, 2000) adults, respectively. However, in this study both tested essential oil showed lower repellency class against *D. porcellus*. Karahroodi et al. (2009) found similar results and showed that *P. crispum* had weak repellency on adults of *P. interpunctella*. While, it has been reported that *P. crispum* had strong repellency activity against *Trialeurodes vaporariorum* (Westw.) in greenhouse tomato production (Tasli et al., 2017). Knowing that the repellent activity of essential oils quickly decrease due to their low molecular masses and high volatility (Jannatan and Rahayu, 2021), this could be explained by the decrease of repellent effect of lower concentration *P. crispum* oil with time.

The decrease of the repellent activity of *P. racemosa* essential oil with increase of concentration could be explained by the interaction and/or inactivation of some oil components at high concentration. In fact, when eugenol the major bioactive compound of *P. racemosa* essential oil was methylated, methyl-eugenol, which can be contained at over 90% in this oil (Tan and Nishida, 2012), may play a physiological role on the toxicity reduction (Chang et al., 2009). The understanding of interaction among compounds contained in *P. racemosa* but also in *P. crispum* essential oils should be an important topic for further research. In general, these results suggest that the essential oils of both plant leaves cannot be used as repellents in the *D. porcellus* control.

The results showed that, *P. racemosa* and *P. crispum* essential oils had insecticidal activity on *D. porcellus* by topical application. Similar findings have been documented by Leyva et al. (2007a, b) which showed that *P. racemosa* essential oil had higher toxicity against *Musca domestica* L. and *Blatella germanica* L., respectively. Moreover, *P. racemosa* oil caused total mortality of *Spodoptera littoralis* (Boisduval) higher than 70% (Pavela, 2013). The toxic effect of *P. racemosa* essential oil to *D. porcellus* could be attributed to the presence of eugenol. In fact, eugenol applied topically on *Sitophilus granarius*, *Sitophilus zeamais*, *Tribolium castaneum* and *P. truncatus* was highly toxic (Obeng-Ofori and Reichmuth, 1997; Huang et al., 2002). Contrary to *P. crispum* essential oil, the contact toxicity of *P. racemosa* oil did not persist with time. The lower persistence of eugenol (Obeng-Ofori and Reichmuth, 1997) could explain these phenomena. Indeed, the loss of toxicity of *P. racemosa* essential oil with exposure time could be due to their high volatility and the quick degradation of active compounds. Concerning *P. crispum* essential oil, its insecticidal activity has been reported

Table 6. Lethal concentration of *P. crispum* and *P. racemosa* essential oil fumigants that kills 50% of exposed *D. porcellus*.

Plant	Exposure (days)	LC ₅₀ (μL/L air)	Slope (± S.E.)	Intercept	χ ²	P-value
<i>P. crispum</i>	1	0.000	-0.552 ± 5.48	3.692	0.67	0.907
	2	0.000	-0.094 ± 2.82	3.912	0.15	0.274
	3	0.004	0.215 ± 1.07	4.460	3.48	0.789
	4	0.078	-0.323 ± 0.72	4.639	8.69	0.958
	5	2.479	-0.535 ± 0.41	5.221	9.69	0.969
	6	2.678	-0.431 ± 0.49	5.185	8.79	0.935
	7	7.224	-0.452 ± 0.45	5.388	6.86	0.804
<i>P. racemosa</i>	1	1.32×10 ⁵	0.519 ± 0.72	2.261	6.00	0.758
	2	2.28×10 ¹²	0.120 ± 2.86	3.102	0.44	0.108
	3	3382.68	0.877 ± 0.38	1.768	5.71	0.074
	4	5775.21	0.773 ± 0.40	2.092	1.00	0.999
	5	2.05×10 ⁹	0.072 ± 4.08	3.445	0.05	0.010
	6	5.07×10 ⁸	0.183 ± 1.47	3.411	1.02	0.997
	7	3.39×10 ⁶	0.134 ± 1.81	3.718	0.17	0.002

**Figure 5.** Weight loss of yam chips caused by *D. porcellus* fumigated with essential oils of *P. crispum* and *P. racemosa* after 7 days of exposure.

against *Aedes aegypti* L. (Intirach et al., 2016). The insecticidal activities of *P. crispum* essential oil can be attributed to the presence of monoterpenoids such as myristicin, which can penetrate into insects and affect their physiological functions (Ebadollahi, 2011). Indeed, the compounds of essential oils exert their activities on insects through neurotoxic effects (Regnault-Roger et al., 2012). Świech and Połec (2013) have reported that myristicin from *P. crispum* essential oil

showed an insecticidal activity against housefly (*M. domestica* L.) and oriental cockroach (*Blatta orientalis* L.). Moreover, high contact toxicity of myristicin against *L. serricorne* adults has been found by Du et al. (2014). Despite the low toxicity of both essential oils, the topical application of these oils on *D. porcellus* had significantly decreased yam chips weight loss showing their potential role in management of this pest.

In this study, *P. crispum* and *P. racemosa* essential oils

demonstrated high insecticidal activity against *D. porcellus* via ingestion of treated yam chips. These results are in accordance with Noudogbessi et al. (2008) which showed high toxicity of treated maize with *P. racemosa* essential oil against *P. truncatus*. The toxicity of both essential oil could be due to the penetration of essential oil compounds in the insect body through the digestive system. These essential oil compounds could act by increasing the gut pH or a reduction in α -amylase activity, which lead to insect mortality (Stefanazzi et al., 2011). Moreover, in treated yam chips, apart from the ingestion toxicity, *D. porcellus* could be rapidly coated with essential oil compounds which lead to contact toxicity by penetration into insect cuticle. The results of this study showed that at 21 days interval time, the mortality of *D. porcellus* due to *P. racemosa* essential oil was absent for the lowest and highest tested concentrations. To understand this phenomenon, on the one hand, the identification of the bioactive compounds in the *P. racemosa* essential oil against *D. porcellus* is necessary and on the other hand the interaction effect of these bioactive compounds must be investigated.

This study revealed that the essential oil of *P. crispum* and *P. racemosa* act as feeding deterrents against *D. porcellus*. Similar investigations were made by Sharaby and El-Nojiban (2015) who showed the feeding deterrence activity of *P. crispum* on *Agrotis ipsilon* (Hufnagel) larvae. Moreover, similarly to this study, Sharaby and El-Nojiban (2015) demonstrated that *P. crispum* essential oil caused high feeding deterrence against *A. ipsilon* but a low contact mortality. The ability of essential oils to reduce feeding of *D. porcellus* is probably due to the presence of main compounds myristicin and eugenol in *P. crispum* and *P. racemosa* essential oils, respectively. Indeed, myristicin exhibited strong antifeedant effect on larvae and imagoes of *Brontispa longissima* (Gestro) (Qin et al., 2010). Similarly, eugenol from *Laurus nobilis* L. essential oil had a feeding deterrent activity against *Mythimna unipuncta* (Haworth) (Muckensturm et al., 1982). Knowing that, oil compounds reduce feeding activity of *D. porcellus* acting on gustatory chemoreceptors (Akhtar et al., 2012), it is important to identify the role of each oil compound to the feeding deterrent effect of both tested oils.

The essential oil of both plants exhibited fumigant activities against *D. porcellus* with different efficacies. According to Jannatan and Rahayu (2021) essential oil fumigants enter the insect body through the tracheae. The fumigant toxicity of *P. crispum* essential oil against *D. porcellus* was dose-dependent and higher than those of *P. racemosa*. This is not surprising because it is known from previous works that *P. crispum* essential oils have strong fumigant toxicity on some storage insect pests such as *C. maculatus* (Massango et al., 2017; Maroufpoor et al., 2016), *Ephestia kuehniella* Zeller and *Plodia interpunctella* (Hübner) (Maroufpoor et al., 2016), but also on field crop insects such as *Trialeurodes vaporariorum* (Westwood) (Mahmoodi et al., 2014). The

fumigant toxic activity of *P. crispum* essential oil could be related to the presence of myristicin. Indeed, the fumigant toxicity of myristicin contained in *P. crispum* essential oil had been shown against *T. vaporariorum* (Mahmoodi et al., 2014). The insecticidal activity of *P. racemosa* essential oils inhaled by *D. porcellus* was weak and more or less stable. Furthermore, contrasting results were reported by Leyva et al. (2007a) and Lee et al. (2002), which showed that, *P. racemosa* had high fumigant toxicity against *M. domestica* and *T. castaneum*, respectively. These discrepancies in biological activity of *P. racemosa* could be related to the differences of chemical composition and tested insect species. The fumigant toxicity of *P. racemosa* could be attributed to the presence of eugenol as major constituent. Indeed, eugenol has been shown to exhibit a fumigant property toward *C. maculatus* (Ajayi et al., 2014). The results showed that the fumigation of *D. porcellus* had significantly reduced the yam chips weight loss. The reduction in food intake by fumigated *D. porcellus* in comparison with control gave a clear picture of the efficacy of both essential oils used as fumigant (Jaya et al., 2012). Fumigant activity of *P. crispum* essential oil is quite promising and they show potential to be developed as possible natural fumigants for control of stored product insects.

Conclusion

In this paper, we report repellent, contact, feeding deterrence, and fumigant activities of essential oils of *P. crispum* and *P. racemosa* leaves against *D. porcellus* for the first time. Both essential oils showed low repellence against *D. porcellus*. While, when they are in contact with *D. porcellus*, both essential oils not only induced mortality but also significantly reduced yam chips weight loss caused by this pest compared to control. However, the ingestion toxicity of essential oil of *P. crispum* was higher than *P. racemosa*. Moreover, *P. crispum* essential oil showed feeding deterrence and fumigant activity against *D. porcellus* adults showing its potential to be developed into antifeedant and fumigants to control against this pest. However, further investigations are necessary to determine which compounds contributing to the toxicity of both essential oils against *D. porcellus*.

CONFLICT OF INTERESTS

The authors have not declared any conflict of interests.

REFERENCES

- Adedoyin OT, Ojuawo A, Adesiyun OO, Mark F, Anigilage EA (2008). Poisoning due to the yam flour consumption in five families in Ilorin, Central Nigeria. *West African Journal of Medicine* 27(1):41-43.

- Adeleke SI (2009). Food poisoning due to yam flour consumption in Kano (Northwest) Nigeria. *Online Journal of Health and Allied Sciences* 8(2):10. <http://www.ojhas.org/issue30/2009-2-10.htm>
- Agyare C, Appiah T, Boakye YD, Apenteng JA (2017). *Petroselinum crispum*: a Review. *Medicinal Spices and Vegetables from Africa*, pp. 527-547.
- Ajayi OE, Appel AG, Fadamiro HY (2014). Fumigation Toxicity of Essential Oil Monoterpenes to *Callosobruchus maculatus* (Coleoptera: Chrysomelidae: Bruchinae). *Journal of Insects*, Article ID 917212, 7 p.
- Akhtar Y, Pages E, Stevens A, Bradbury R, da Camara CAG, Isman MB (2012). Effect of chemical complexity of essential oils on feeding deterrence in larvae of the cabbage looper. *Physiological Entomology* 37:81-91.
- Alitonou GA, Noudogbessi JP, Sessou P, Tonouhewa A, Avlessi F, Menut C, Sohounhloue DCK (2012). Chemical composition and biological activities of essential oils of *Pimenta racemosa* (Mill.) J. W. Moore from Benin. *International Journal of Biosciences* 22(9):1-12.
- Babarinde SA, Sosina A, Oyeyiola EI (2008). Susceptibility of the selected crops in storage to *Sitophilus zeamais* Motschulsky in southwestern Nigeria. *Journal of Plant Protection Research* 48(4):541-549.
- Behanzin JG, Chabi BC, Chabi WN, Ogbon EA, Konfo TRC, Ahissou H, Sezan A, Baba-Moussa L (2018). Evaluation of the Importance of yam cossettes flour couscous (wassa-wassa tchigan) in the diabetics diet in Benin. *American Journal of Food Science and Technology* 6(1):1-6.
- Caballero-Gallardo K, Olivero-Verbel J, Stashenko EE (2012). Repellency and toxicity of essential oils from *Cymbopogon martini*, *Cymbopogon flexuosus* and *Lippia origanoides* cultivated in Colombia against *Tribolium castaneum*. *Journal of Stored Products Research* 50:62-65
- Chang CL, Cho IK, Li QX (2009). Insecticidal Activity of Basil Oil, trans-Anethole, Estragole, and Linalool to Adult Fruit Flies of *Ceratitis capitata*, *Bactrocera dorsalis*, and *Bactrocera cucurbitae*. *Journal of Economic Entomology* 102(1):203-209.
- Chu SS, Du SS, Liu QZ, Liu QR, Liu ZL (2012). Composition and insecticidal activity of the essential oil of *Artemisia igniaria* Maxim. flowering aerial parts against *Sitophilus zeamais* Motschulsky (Coleoptera: Curculionidae). *Journal of Medicinal Plants Research* 6(16):3188-3192.
- Contreras-Moreno BZ, Velasco JJ, Rojas JC, Méndez LC, Celis MT (2016). Antimicrobial activity of essential oil of *Pimenta racemosa* var. *racemosa* (Myrtaceae) leaves. *Journal of Pharmacy and Pharmacognosy Research* 4(6):224-230.
- Craft DJ, Setzer WN (2017). The volatile components of parsley, *Petroselinum crispum* (Mill.) Fuss. *American Journal of Essential Oils and Natural Products* 5(1):27-32.
- Du S, Yang K, Wang C, You C, Geng Z, Guo S, Deng Z, Liu Z (2014). Chemical Constituents and Activities of the Essential Oil from *Myristica fragrans* against Cigarette Beetle *Lasioderma serricorne*. *Chemistry and Biodiversity* 11:1449-1456. doi:10.1002/cbdv.201400137
- Ebadollahi A (2011). Antifeedant activity of essential oils from *Eucalyptus globulus* Labill and *Lavandula stoechas* L. on *Tribolium castaneum* Herbst (Coleoptera: Tenebrionidae).
- El-Zaieddi H, Martínez-Tomé J, Calín-Sánchez A, Francisco B, Carbonell-Barrachina AA (2016). Volatile composition of essential oils from different aromatic herbs grown in Mediterranean regions of Spain. *Foods* 5(41):1-13.
- Food and Agriculture Organization (FAO) (2013/2018). FAOSTAT Database. Food and Agriculture Organization, Roma, Italy. Available online at URL: www.fao.org
- Guruprasad BR, Akmal P (2014). Assessment of repellency and insecticidal activity of *Ajuga parviflora* (Benth) and *Trichilia connaroides* (W&A) leaf extracts against stored product insects. *Journal of Entomology and Zoology Studies* 2(4):221-226.
- Huang Y, Ho SH, Lee HC, Yap YL (2002). Insecticidal properties of eugenol, isoeugenol and methyleugenol and their effects on nutrition of *Sitophilus zeamais* Motsch. (Coleoptera: Curculionidae) and *Tribolium castaneum* (Herbst) (Coleoptera: Tenebrionidae). *Journal of Stored Products Research* 38(5):403-412.
- Intirach J, Junkum A, Lumjuan N, Chaithong U, Jitpakdi A, Riyong D, Wannasan A, Champakaew D, Muangmoon R, Chansang A, Pitasawat B (2016). Antimosquito property of *Petroselinum crispum* (Umbelliferae) against the pyrethroid resistant and susceptible strains of *Aedes aegypti* (Diptera: Culicidae). *Environmental Science and Pollution Research* 23:23994-24008.
- Jannatan R, Rahayu R (2021). Fumigant toxicity and repellency of citronella grass essential oil (*Cymbopogon nardus* (L.) Rendle) to German cockroaches (*Blattella germanica* L.). *European Journal of Biological Research* 11(3):267-273. <http://dx.doi.org/10.5281/zenodo.4670508>
- Jaya SP, Prakash B, Dubey NK (2012). Insecticidal activity of *Ageratum conyzoides* L., *Coleus aromaticus* Benth. and *Hyptis suaveolens* (L.) Poit essential oils as fumigant against storage grain insect *Tribolium castaneum* Herbst. *Journal of Food Science and Technology* 51(9):2210-2215. 10.1007/s13197-012-0698-8
- Karahroodi ZR, Moharrampour S, Rahbarpour A (2009). Investigated repellency effect of some essential oils of 17 native medicinal plants on adults *Plodia interpunctella*. *American-Eurasian Journal of Sustainable Agriculture* 3(2):181-184.
- Kiba DI, Hgaza VK, Aighewi B, Aké S, Barjolle D, Bernet T, N'Diby L, Ilboudo LJ, Nicolay G, Oka E, Ouattara YF, Pouya N, Six J, Frossard, E (2020). A Transdisciplinary approach for the development of sustainable yam (*Dioscorea* sp.) production in West Africa. *Sustainability* 12(10):4016. <http://dx.doi.org/10.3390/su12104016>
- Lee BH, Lee SE, Annis PC, Pratt SJ, Park BS, Tumaalii F (2002). Fumigant toxicity of essential oils and monoterpenes against the red flour beetle, *Tribolium castaneum* Herbst. *Journal of Asia-Pacific Entomology* 5(2):237-240.
- Leyva M, Tacoronte JE, Marquetti MC (2007b). Chemical composition and lethal effect of essential oil from *Pimenta racemosa* (Myrtales: Myrtaceae) on *Blattella germanica* (Diptera: Blattellidae). *Revista Cubana de Medicina Tropical* 59(2):154-158.
- Leyva MI, Tacoronte JE, Navarro A, Montada D, Bello A, Marquetti MC (2007a). Estudio de laboratorio del aceite esencial de *Pimenta racemosa* (Myrtales: Myrtaceae) y su posible utilización para el control de *Musca domestica* (Diptera: Muscidae) *Revista CENIC Ciencias Biológicas* 38(1):18-19.
- Liu ZL, Ho SH (1999). Bioactivity of the essential oil extracted from *Evodia rutaecarpa* Hook f. et Thomas against the grain storage insects, *Sitophilus zeamais* Motsch. and *Tribolium castaneum* (Herbst). *Journal of Stored Products Research* 35:317-328.
- Loko YL, Dansi A, Tamo M, Bokoron-Ganta AH, Assogba P, Dansi M, Vodouhè R, Akoegninou A, Sanni A (2013). Storage insects on yam chips and their traditional management in Northern Benin. *The Scientific World Journal*, Article ID 484536, 11p. <http://dx.doi.org/doi:10.1155/2013/484536>
- Loko YLE, Medegan Fagla S, Kassa P, Ahouansou CA, Toffa J, Glinma B, Dougnon V, Koukou O, Djogbenou SL, Tamò M, Gbaguidi F (2020). Bioactivity of essential oils of *Cymbopogon citratus* (DC) Stapf and *Cymbopogon nardus* (L.) W. Watson from Benin against *Dinoderus porcellus* Lesne (Coleoptera: Bostrichidae) infesting yam chips. *International Journal of Tropical Insect Science* (2020). <https://doi.org/10.1007/s42690-020-00235-3>
- Mahmoodi L, Valizadegan O, Mahdavi V (2014). Fumigant toxicity of *Petroselinum crispum* L. (Apiaceae) essential oil on *Trialeurodes vaporariorum* (Westwood) (Hemiptera: Aleyrodidae) adults under greenhouse conditions *Journal of Plant Protection Research* 54(3):294-298.
- Maroufpoor M, Ebadollahi A, Vafaei Y, Badiie E (2016). Chemical Composition and Toxicity of the Essential Oil of *Coriandrum sativum* L. and *Petroselinum crispum* L. Against against tThree Storedstored-pProduct ilnsect pPests. *Journal of Essential Oil Bearing Plants* 19(8):1993-2002.
- Massango HGLL, Faroni LRA, Haddi K, Heleno FF, Jumbo LOV, Oliveira EE (2017). Toxicity and metabolic mechanisms underlying the insecticidal activity of parsley essential oil on bean weevil, *Callosobruchus maculatus*. *Journal of Pest Science* 90(2):723-733.
- Muckensturm B, Duplay D, Mohammadi F, Moradi A, Robert PC, Simonis MT, Kielen JC (1982). Role of natural phenylpropanoids as antifeeding agents for insects. *Les Colloques de l'INRA (médiateurs chimiques agissant sur le comportement des insectes)* 7:131-135.

- Mukanga M, Deedat Y, Mwangala FS (2010). Toxic effects of five plant extracts against the larger grain borer, *Prostephanus truncatus*. African Journal of Agricultural Research 5(24):3369-3378.
- Mulugeta T, Unnithan CR, Tesfay D (2015). Phytochemical screening, characterization and biological activities of *Petroselinum crispum* (Parsley) leaf oil. World Journal of Pharmacy and Pharmaceutical Sciences 4:142-151.
- Nawel O, Ahmed H, Douniazad EA (2014). Phytochemical analysis and antimicrobial bioactivity of the Algerian parsley essential oil (*Petroselinum crispum*). African Journal of Microbiology Research 8(11):1157-1169.
- Noudogbessi J-P, Kossou D, Sohounloulé DCK (2008). Effet Insecticide, Ovicide et Larvicide des Huiles Essentielles de *Pimenta racemosa* (Miller) et de *Chromolaena odorata* (L. Robinson) sur le Grand Capucin (*Prostephanus truncatus* (Horn)) du Maïs. Journal de la Société Ouest-Africaine de Chimie 26:41-51.
- Obeng-Ofori D, Reichmuth Ch (1997). Bioactivity of eugenol, a major component of essential oil of *Ocimum suave* (Wild.) against four species of stored-product Coleoptera, International Journal of Pest Management 43(1):89-94. <http://dx.doi.org/DOI:10.1080/096708797229040>
- Ojimelukwe PC, Adler C (2000). Toxicity and repellent effects of eugenol, thymol, linalool, menthol and other pure compounds on *Dinoderus bifoveatus* (Coleoptera: Bostrichidae). Journal of Sustainable Agriculture and the Environment 2(1):47-54.
- Omohimi C, Piccirillo C, Ferraro V, Roriz MC, Omemu MA, Dias Santos SM, Da Ressurreição S., Abayomi L., Adebawale A, Vasconcelos MW, Obadina O, Sanni L, Pintado MME (2019). Safety of Yam-Derived (*Dioscorea rotundata*) Foodstuffs—Chips, Flakes and Flour: Effect of Processing and Post-Processing Conditions. Foods 8(1):12. <http://dx.doi.org/10.3390/foods8010012>
- Oni MO, Omoniyi AO (2012). Studies on temperature influence on oviposition and development of immature stages of the yam beetle *Dinoderus porcellus* Lesne (Coleoptera: Bostrichidae) on dried yam species. Journal of Agricultural Science 4:213–218.
- Othira JO, Onek LA, Deng LA, Omolo EO (2009). Insecticidal potency of *Hyptis spicigera* preparations against *Sitophilus zeamais* and *Tribolium castaneum* (herbst) on stored maize grains. African Journal of Agricultural Research 4:187-192.
- Pavela R (2013). Sublethal Effects of Some Essential Oils on the Cotton Leafworm *Spodoptera littoralis* (Boisduval). Journal of Essential Oil Bearing Plants 15 (1):144-156.
- Petropoulos SA, Daferera D, Akoumianakis CA, Passam HC, Polissiou MG (2004). The effect of sowing date and growth stage on the essential oil composition of three types of parsley. Journal of the Science of Food and Agriculture 84:1606-1610.
- Püntener W (1981). Manual for field trials in plant protection. 2nd ed. Agricultural Division; Ciba-Geigy Limited.
- Qin W, Huang S, Li C, Chen S, Peng Z (2010). Biological activity of the essential oil from the leaves of *Piper sarmentosum* Roxb. (Piperaceae) and its chemical constituents on *Brontispa longissima* (Gestro) (Coleoptera: Hispididae). Pesticide Biochemistry and Physiology 96(3):132-139.
- Regnault-Roger C, Vincent C, Arnason TJ (2012). Essential oils in insect control: low-risk products in a high-stakes world. Annual Review of Entomology 57:405-424.
- Saroukolai A, Nouri-Ganbalani G, Rafiee-Dastjerdi H, Hadian J (2014). Antifeedant activity and toxicity of some plant essential oils to Colorado potato beetle, *Leptinotarsa decemlineata* Say (Coleoptera: Chrysomelidae). Plant Protection Science 50:207-216.
- Sharaby A, El-Nojiban A (2015). Evaluation of some plant essential oils against the black cutworm *Agrotis ipsilon*. Global Journal of Advanced Research 2(4):701-711.
- Smith CJ, Perfetti TA (2020). A comparison of the persistence, toxicity, and exposure to high-volume natural plant-derived and synthetic pesticides. Toxicology Research and Application 4:1-15.
- Snoussi M, Dehmani A, Noumi E, Flamini G, Papetti A (2016). Chemical composition and antibiofilm activity of *Petroselinum crispum* and *Ocimum basilicum* essential oils against *Vibrio spp.* strains. Microbial Pathogenesis 90:13-21. doi:10.1016/j.micpath.2015.11.004
- Stefanazzi N, Stadler T, Ferrero A (2011). Composition and toxic, repellent and feeding deterrent activity of essential oils against the stored-grain pests *Tribolium castaneum* (Coleoptera: Tenebrionidae) and *Sitophilus oryzae* (Coleoptera: Curculionidae). Pest Management Science 67:639-646.
- Świech K, Poleć I (2013). Comparison of the insecticidal effectiveness of synthetic and natural myristicin against housefly (*Musca domestica* L.) and oriental cockroach (*Blatta orientalis*). Chemik 67(11):1115-1120.
- Tan KH, Nishida R (2012). Methyl eugenol: Its occurrence, distribution, and role in nature, especially in relation to insect behavior and pollination. Journal of Insect Science 12:56.
- Tasli M, Yoldas Z, Öztekin GB, Tüzel Y (2017). Effects of some repellent plants on greenhouse whitefly *Trialeurodes vaporariorum* (Westw.) in greenhouse tomato production. Acta Horticulturae 1164:407-412
- Umogbai VI (2013). Design, Construction and Performance Evaluation of an Underground Storage Structure for Yam Tubers. International Journal of Scientific and Research Publications 3(5):1-7.
- You CX, Jiang HY, Zhang WJ, Guo SS, Yang K, Lei N, Ma P, Geng ZF, Du SS (2015). Contact Toxicity and Repellency of the Main Components From the Essential Oil of *Clausena anisum-olens* Against Two Stored Product Insects. Journal of Insect Science 15(1):87. <https://doi.org/10.1093/jisesa/iev071>