



E-ISSN: 2320-7078

P-ISSN: 2349-6800

JEZS 2016; 4(6): 408-414

© 2016 JEZS

Received: 23-09-2016

Accepted: 25-10-2016

**Roland Eric Yessinou**

University of Abomey-Calavi (UAC),  
Polytechnic School of Abomey-Calavi  
(EPAC) Production and Animal Health  
Department, Applied Biology Research  
Laboratory (LARBA) 01 Po. Box: 2009  
Cotonou, Benin

**Yao Akpo**

University of Parakou (UP), Faculty of  
Agronomy, Po Box 123 Parakou, Benin

**Camus Adoligbe**

University of Abomey-Calavi (UAC),  
Polytechnic School of Abomey-Calavi  
(EPAC) Production and Animal Health  
Department, Applied Biology Research  
Laboratory (LARBA) 01 Po. Box: 2009  
Cotonou, Benin

**Justin Adinci**

University of Abomey-Calavi (UAC),  
Polytechnic School of Abomey-Calavi  
(EPAC) Production and Animal Health  
Department, Applied Biology Research  
Laboratory (LARBA) 01 Po. Box: 2009  
Cotonou, Benin

**Marc Napoléon Assogba**

University of Abomey-Calavi (UAC),  
Polytechnic School of Abomey-Calavi  
(EPAC) Production and Animal Health  
Department, Applied Biology Research  
Laboratory (LARBA) 01 Po. Box: 2009  
Cotonou, Benin

**Benoît Koutinhoun**

University of Abomey-Calavi (UAC),  
Polytechnic School of Abomey-Calavi  
(EPAC) Production and Animal Health  
Department, Applied Biology Research  
Laboratory (LARBA) 01 Po. Box: 2009  
Cotonou, Benin

**Issaka Youssao Abdou Karim**

University of Abomey-Calavi (UAC),  
Polytechnic School of Abomey-Calavi  
(EPAC) Production and Animal Health  
Department, Applied Biology Research  
Laboratory (LARBA) 01 Po. Box: 2009  
Cotonou, Benin

**Souaïbou Farougou**

University of Abomey-Calavi (UAC),  
Polytechnic School of Abomey-Calavi  
(EPAC) Production and Animal Health  
Department, Applied Biology Research  
Laboratory (LARBA) 01 Po. Box: 2009  
Cotonou, Benin

**Correspondence****Roland Eric Yessinou**

University of Abomey-Calavi (UAC),  
Polytechnic School of Abomey-Calavi  
(EPAC) Production and Animal Health  
Department, Applied Biology Research  
Laboratory (LARBA) 01 Po. Box: 2009  
Cotonou, Benin

## Resistance of tick *Rhipicephalus microplus* to acaricides and control strategies

**Roland Eric Yessinou, Yao Akpo, Camus Adoligbe, Justin Adinci, Marc Napoléon Assogba, Benoît Koutinhoun, Issaka Youssao Abdou Karim and Souaïbou Farougou**

### Abstract

Ticks are bloodsucking arthropods mandatory and represent a real problem for cattle. The aim of this review is to make an inventory of acaricide resistance and ways to fight against the tick *Rhipicephalus microplus*. The main classes of synthetic miticides are arsenical, organochlorine, organophosphates, amidines, macrocyclic lactones, growth regulators and phenylpyrazoles. Pyrethroids are the most widespread and used acaricide in the fight against ticks. Resistance to these classes acaricide was reported in all countries where the tick *Rhipicephalus microplus* is present. To deal with these ectoparasites, a variety of control methods are used. These include ecological, biological, and genetic approaches. Proper control of the ticks will allow farmers to save and limit the losses related to diseases transmitted by ticks. It is urgent to propose and develop new approach to fight against tick *Rhipicephalus microplus*.

**Keywords:** *Rhipicephalus microplus* resistance, control method, cattle

### 1. Introduction

*Rhipicephalus microplus* [1] is a cattle tick, whose development cycle is monoxenous. Although it is originally from India and Indonesia, this tick has shown easy adaptation to tropical and subtropical conditions [2]. It is one of the major obstacles to the development of cattle breeding. Historically, it was introduced in Eastern and Southern Africa from Southern Asia through Madagascar after the outbreak of rinderpest in 1896 [3]. In West Africa, it was discovered for the first time in Ivory Coast by imports of live cattle from Brazil in 2007 [4, 5] and recently in Benin, Burkina Faso and Mali [6]. An economic loss (reduction of weight, decrease of meat and milk quality) due to presence of tick is huge. In Brazil for example, losses due to its direct and indirect impact have been estimated to two billion US dollars in 2000 [7]. In Australia, they are estimated to 100 million Australian Dollars a year [8] and USD 48 million a year in Mexico [9]. Besides, tick can transmit pathogens such as viruses, bacteria and protozoa. The prevalence of tick borne diseases is 1.7% to 36% in the Caribbean [10], 6.25% in Zambia, and 25.0% in Zimbabwe [11]. A prevalence of 9% and 10.8% was recorded in Burkina Faso and Benin respectively [12]. The use of synthetic acaricide currently remains the only way to fight [13]. Introduced in 1895, Arsenical compounds were first used to control ticks [14], and in late 1930s, the first organochlorine compounds were introduced in the market to control ticks that became resistant to arsenic [15]. Synthetic pyrethroids were introduced in the 1970s [15] and they are currently adopted by most of the breeders. However, tick resistance, particularly *Rhipicephalus microplus* to this class of acaricide is common in many countries including Brazil [16, 17], Australia [18, 19], Argentina, New Caledonia, South Africa [20, 21], Benin, Cote d'Ivoire [4, 5], Burkina Faso and Mali [6]. Pyrethroids have their action site localized in the nervous system of arthropods [22]. Their molecular targets include the voltage-dependent sodium channel [23] and acetylcholinesterase (AChE). It is shown that occurrence of mutations in the genes encoding for these target proteins result in structural changes that intend to reduce their affinity with miticides. The detection of genes involved in ticks resistance and their mechanisms is essential to establish a sustainable strategy of control of the tick [24].

### Material and methods

In this study, we made a summary of different research works carried out on tick resistance to

acaricide particularly *Rhipicephalus microplus* resistance and the control methods that are commonly used. In this review, we explained the concept of ticks' resistance to acaricide, the resistance mechanism particularly to pyrethroids which is the most used family of acaricide. Control methods were also investigated with a focus on methods that are environmental friendly and sustainable.

### Concept of tick resistance to acaricide

Resistance is the occurrence, in a population, of individuals with the ability to tolerate doses of toxic substance that are lethal for the majority of individuals in a normal population of the same species<sup>[25]</sup>. The toxicity of acaricide is the result of their interaction with biological targets present in the tick. Several steps are needed before the acaricide exerts its action. It must first be in contact with the arthropod, enter his body, be converted in some cases into the active metabolite and then transported to the action site. Each of these steps is under the control of one or more genes. Any mechanism that changes one of these steps can therefore lead to a resistance<sup>[26, 27]</sup>. Ticks' resistance to acaricide may therefore result from several mechanisms, which can be for instance a change in absorption or excretion of the acaricide, a change in metabolic pathways that allow the acaricide degradation or finally a modification of its action site. The metabolic resistance is due to the increase of the enzymes activity in charge of miticides detoxification. These include sterases, oxidases cytochrome P450 and glutathione S-transferases<sup>[28]</sup>. The increase in activity of the enzyme may be due to a change in the conformation of the enzyme making it more efficient or due to its over production<sup>[29]</sup> that may sometimes be up to 12% of total protein of an individual<sup>[30]</sup>. The overproduction of enzyme may be due to a modification of a regulatory gene controlling the expression level of the enzyme<sup>[31, 32]</sup>, an increase in the number of copies of the genes which code for these enzymes<sup>[33]</sup>; this is a complementary mechanism to the *kdr* mutation. As for the changes in the acaricide action site, they are due to point mutations occurrence in the genes coding for target proteins. This causes structural changes that reduce their affinity with miticides. These mechanisms can also cause cross-resistance to all miticides acting on the same site<sup>[34]</sup>. In most species, the mutation points are generally found on genes encoding for proteins, as that the amino acid change must not only reduce the affinity of the acaricide with the action site, but also preserve the initial functions of the protein to a level that is compatible with the survival of the tick. In most cases, it is likely that genes that confer resistance are already present at very low level in the tick population prior to the introduction of a new acaricide. The lack of standardized techniques for diagnosing the resistance or lack of resistance monitoring system appears to be the main challenge in countries where *Rhipicephalus microplus* is introduced. Actually, resistance suspicions occur when there is a treatment failure in the fight against cattle ticks. Although treatment failure is often the result of inappropriate preparation or application of acaricide, ticks persistence after frequent and correctly applied treatments indicate that resistance to the acaricide is permanent<sup>[35]</sup>. The intensive use of miticides led to the development of resistance to almost all major classes of miticides in many countries including Australia, Mexico, Brazil, New Caledonia, South Africa, USA etc. the failure of tick control is not always due to the resistance, but can be caused by factors such as faulty equipment, improper dose or use of outdated and therefore ineffective chemicals. Therefore, when the failure of the fight

against ticks is observed, the suspected resistance should be confirmed before selecting a new acaricide. Once resistance is confirmed, breeders and livestock sector actors should be informed about the use of alternative chemicals for the control of resistant populations. Therefore, the sensitivity of resistant ticks potential replacement product must be tested<sup>[36]</sup>. In addition, since the cross-resistance is known to occur between compounds of the same class, but not always, more than one representative of each class to be tested. Overall control of the ticks should facilitate the detection of resistance at an early stage is essential to slow its spread early in avoiding resistant individuals selection. It is impossible to understand the mechanisms involved in the development of resistance of ticks without understanding the mode of action of acaricide.

### *Rhipicephalus microplus* resistance to synthetic pyrethroids

Synthetic pyrethroids have been introduced in the 1970s<sup>[15]</sup> and are currently widely used in our farms. Pyrethroids syntheses are highly effective insecticidal and acaricide, they are not very toxic to mammals and are highly biodegradable. Already in the late 1980s, resistance has been reported in Australia<sup>[37]</sup>, Brazil<sup>[38]</sup> and in Mexico in 1994<sup>[39]</sup>. Today pyrethroids resistance is extremely common and is prevalent in all countries where resistance studies were performed<sup>[16, 17]</sup>. The voltage-gated sodium channel is the target site for the activity of pyrethroids and has been studied in many species of arthropods. Three types of mutation in the sodium channel are associated with resistance to pyrethroids ticks<sup>[19, 40]</sup>. A study on *Rhipicephalus microplus* Mexico populations showed a substitution type of mutation in domain III, section 6, and position of nucleotide 2134. The replacement of thymine by adenine has transformed the amino acid, isoleucine to phenylamine<sup>[41, 42]</sup>. This led to the occurrence of a change of structure of acaricide receptor located on the sodium channel gene. As a result, ticks that are subjected to such changes become insensitive to pyrethroid. Thus, two other alternative types of mutations have been identified in the domain II in the segment 4, the sodium channel gene: The leucine is replaced by isoleucine, which results in the substitution of cytosine by adenine and still on the segment 4 we see the replacement of the amino acid glycine by valine due to the substitution at position 215 of guanine to thymine in the Australian population<sup>[43, 44]</sup>. The mutation of domain II is also found in the population of Brazil<sup>[18]</sup>, South Africa and Argentina<sup>[21]</sup>. This mutation gives tick acaricide resistance as permethrin, cypermethrin and flumethrin<sup>[43]</sup>. Domain III mutation of the sodium channel gene associated with resistance of *Rhipicephalus microplus* was noted on the stump of the US and Mexico<sup>[16, 45]</sup>. In Mexico, studies show that *Rhipicephalus microplus* is resistant to deltamethrin, cypermethrin and flumethrin<sup>[46]</sup>. The tick *Rhipicephalus microplus* resistance to Cypermethrin, flumethrin was also reported in Brazil<sup>[38]</sup>. The search for a new management approach is important for the control of ticks.

### Tick resistance to other classes of miticides

Control of cattle ticks with chemical compounds began in the late nineteenth century with the use of arsenic. This class of compounds was followed by organochlorines, organophosphates, amidines, synthetic pyrethroids, phenylpyrazoles, macrocyclic lactones, growth regulators and spinosyns. The introduction of each new acaricide class was followed by the emergence of resistance<sup>[21]</sup>. Arsenical compounds were first used to control ticks, introduced in

1895<sup>[14]</sup>. In 1896, a deep immersion based on arsenic has been used in Queensland, Australia, to control the tick by a local farmer. This deep immersion arsenic was quickly adopted in other areas such as USA, South Africa and Cuba and the disposal has become a widespread practice in Australia<sup>[47]</sup>. The immersion should be frequent due to the very short persistence<sup>[24]</sup> of arsenic<sup>[48]</sup>. After about 40 years of use, the first case of resistance to arsenic *Rhipicephalus microplus* was reported in 1936<sup>[49]</sup>. Thus, the first organochlorine compounds were introduced in the market as acaricide in 1939<sup>[15]</sup> to control ticks resistant to arsenic. More compounds became available in the mid-1940; organochlorines have a high efficiency, a long residual activity and a broad spectrum of action. Moreover, they had the advantage of being less toxic and less expensive than arsenic. The first case of resistance to organochlorine has been observed in Brazil in 1952<sup>[50]</sup>. A decade later, in 1962, the use of all organochlorine was forbidden to fight against ticks, due to residues in meat, milk and the environment resulting from its poor biodegradability and affinity for lipids<sup>[15]</sup>. To contain the infestation of ticks, organophosphates began to be used as ectoparasiticides in the mid-1950. They were used to fight against ticks that have become resistant to organochlorines. Organophosphates were less stable and less persistent than organochlorines, organophosphates but some are very toxic to mammals<sup>[51]</sup>. Organophosphate resistance was first appeared in Australia in the mid-1960s<sup>[52]</sup>. In addition, organophosphate resistance was also observed on a Christmas tree Texas strain<sup>[53]</sup>. Thus amidines have been introduced for the control of ticks in the mid 1970<sup>[54]</sup>. Today, amitraz is the main active ingredient in this class of acaricide. The resistance to amitraz appeared 4 to 10 years after its first use in different parts of the world and has been identified for the first time in the early 1980, Australia<sup>[54]</sup>. Since then, resistance has also been observed in Mexico, South America,

South Africa and New Caledonia. However, in 2007,<sup>[40]</sup> reported that amitraz was still one of the most popular miticides to control ticks on cattle in Australia, Latin America and Southern Africa<sup>[21]</sup>. As for macrocyclic lactones, they were introduced to the market in 1981<sup>[55]</sup> and are divided into two categories<sup>[56]</sup>: avermectin and milbemycin oxime. Macrocyclic lactones are systemically active against ticks and have a longer residual activity than synthetic pyrethroids and are active against a range of arthropods and nematodes<sup>[56]</sup>. However, the long wait for meat and milk can limit the use of this class in cattle. Resistance to avermectin (doramectin and ivermectin) was first reported in Brazil in 2001<sup>[57]</sup> and was later also reported in Mexico ivermectin<sup>[58]</sup>. Nowadays, fipronil is the only compound of phenylpyrazoles use in livestock for the fight against cattle ticks and its use began in the mid-90<sup>[59]</sup>. It has a long residual activity<sup>[56]</sup> and continues up to five weeks in the field<sup>[60]</sup>. Resistance to fipronil was reported for the first time in 2007 with Uruguay<sup>[61]</sup>, and later in Brazil<sup>[62]</sup>. Growth regulators are one of the last classes of miticides, with the first compound representing the class, fluzaron available in the market acaricide since 1994 in Australia. The resistance has not been reported in the literature, but a case was presented in a poster session of Congress in 2010<sup>[63]</sup>. A new strategy must be implemented to fight against ticks.

### Control strategies

The fight against ticks is not only to limit the losses they cause, but also those due to diseases transmitted by ticks or associated with ticks. This struggle must be conducted in a concerted and rational manner based on the various genera and species present in the area, but especially in the light of understanding their respective biology. Thus, several control methods are used worldwide (Table 1).

**Table 1:** Control method against ticks

Methods	Processes	References
Fight With Trap	Trap and ticks removal from the host animal	host animals Products invasion against ticks <sup>[66 ; 81]</sup>
Traditional Method	Manual removal	[70 ; 71 ; 69 ; 72 ; 73 ; 74 ; 75 ; 76]
	Application of traditional products containing harmful substances to ticks	
Chemical Method	Use of synthetic miticides against ticks	[35 ; 2 ; 18 ; 61 ; 82]
Ecological Control	rotational grazing to control the infestation	[ 63 ; 66 ; 64 ; 65]
Biological Control	predators to fight against ticks	[ 67 ; 83, 84]
Genetic Control	Selection of animals resistant to ticks	[ 77 ; 79 ; 80 ; 85 ; 86 ; 87]

Several chemical methods of ticks control were developed, including the use of miticides such as organochlorines, organophosphates, macrocyclic lactones amidines and synthetic pyrethroids. At this level, it is equally important to note that misuse of these miticides leads to the development of resistance of ticks to acaricide. Chemical control is now widely used in our farms but it is very costly to farmers, who do not always have the means to face it. It is extremely difficult to control ticks and the diseases they transmit. Today controlling the infestation of cattle is based mainly on synthetic miticides. The indiscriminate use of these substances also poses pollution problems etc. There is therefore great interest in the development of alternative methods of control. Some are described briefly here.

- **Ecological and biological control**

The ecological approach is to modify the habitat of the tick so as to cripple its reproductive cycle<sup>[64]</sup>, based on a driving rotational grazing. Indeed, it amounts to prevent contact

between ticks and host and thus to an interruption of the cycle. According to Barre<sup>[65]</sup> cited by<sup>[66]</sup>, this process is conceivable that in the case of modern farming (intensification) because it is necessary to have large clean pasture land can be withdrawn for several weeks or months of animal feed<sup>[67]</sup>. Many parasites, parasitoids or predators may have a negative impact on the ticks present in the medium. But note that there is little use of these organisms test (bacteria, fungi, nematodes and insects) to reduce tick populations<sup>[68]</sup>. It was recently found that fungal spores (*Beauveria bassiana* and *Metarhizium anisopliae*) had some activity against *Rhipicephalus appendiculatus* and the tropical bond tick infesting rabbits and cattle on which they were sprayed<sup>[68]</sup> regulating ixodienne and the population to which they are associated. This is the case of the ant *Solenopsis geminata*<sup>[67]</sup> predator of the tick *Rhipicephalus microplus*.

### • Traditional methods

The manual diptank is the most commonly used method by farmers in West Africa. It is a method that is easily applicable to a small population of animals. However, it turns out binding for high parasite load, which is the case of *Rhipicephalus microplus*. It is time consuming and labor intensive, especially when the infestation is observed in the beginning of rainy season (farm work), or end of the rainy season [66]. Breeders from the perspective of destroying ticks on animals, use a variety of plants containing active ingredients harmful to ticks such as [69]: *Actaea spicata*, a plant used externally in the form of lotions or mixed with fat pork; it heals and kills scabies mites. Studies on the mixture of essential oils of *Syzygium aromaticum* and *Cymbopogon citratus* gave a very satisfactory result on females engorged *Rhipicephalus microplus* [70]; the essential oil of *Azadirachta indica* share its bitter properties and its characteristic odor, has a potential acaricide [71]; *Ricinus communis* extract had acaricide effect on ticks [72], it helps to fight against many mites and is harmless to higher animals. Studies on *Tephrosia vogelii* in Benin [73], Zimbabwe [74] and Congo [75] revealed that the leaves, seeds and other parts of the plant contain active ingredients with acaricide properties. The leaves of *Mitracarpus villosus* and *Thevetia nerifolia* have larvicidal activity against *Rhipicephalus microplus* [76], the extract of *Annona squamosa* an acaricide property on the tick [77]. View the number of plants with activity against ticks, the use of plant extracts for control seems to be a promising alternative.

### • Genetic resistance

The external parasitism by ticks is a major constraint to the cattle. The use of genetically resistant animals is a complementary method to fight these ectoparasites. This alternative measure is introduced in Australia [78] to combat ticks. Studies are underway to highlight the breeds resistant to ticks. For example, resistance to *Rhipicephalus microplus* varies according to the breed but is usually more pronounced for example in Zebu cattle (*Bos indicus*) than in cattle (*Bos taurus*) [79]. In Australia, and Sahiwal breeds Brahm ( *Bos indicus* ), respectively selected for meat and milk showed a natural resistance to *Rhipicephalus microplus* higher than *Bos taurus* breeds [80]. The selection of resistant animal breed tick occurs by crossing between animals in order to obtain a product having a natural immunity to these ectoparasites [81].

### Conclusion

Ticks, for their direct impact and the diseases they transmit are a major constraint to the development of cattle breeding in infested cattle herds. The use of synthetic acaricide for control leads to a buildup of resistance. Control of *Rhipicephalus microplus* by the selection of resistant cattle or by the use of plant extracts appears to be a promising alternative to reduce the invasion of ticks.

### Acknowledgements

The authors declare that there is no conflict of interests.

### Reference

- Canestrini G. Prospetto dell'acarofauna italiana Famiglie: Tarsonemini Tiroglifini appendice agli Analgesini. Padova. 1888; 3:309-425.
- Barré N, Uilenberg G. Propagation de parasites transportés avec leurs hôtes: cas exemplaires de deux espèces de tiques du bétail. In Revue scientifique et technique de l'office international. 2010; 29(1):135-147.
- Hoogstraal H. African Ixodoidea ticks of the Sudan with special Reference to Equatorial Province and with preliminary reviews of the genera *Boophilus*, *Margaporus* and *Hyalomma*. Bureau of Medicine and Surgery, Navy Medicine, Washington. 1956; 1:1101.
- Madder M, Adehan S, De Deken R, Adehan R, Lokossou R. New foci of *Rhipicephalus microplus* in West Africa. Experimental and Applied Acarology. 2012; 56(4):385-390.
- Madder M, Thys E, Geysen D, Baudoux C, Horak I. *Boophilus microplus* ticks found in West Africa. Experimental and Applied Acarology. 2007; 43:233-234.
- Adakal H, Biguezoton A, Zoungrana S, Courtin F, De Clercq EM, Madder M. Alarming spread of the Asian cattle tick *Rhipicephalus microplus* in West African another three countries are affected: Burkina Faso, Mali and Togo. Experimental and Applied Acarology. 2013; 61: 383.
- Lovis L, Guerrero FD, Miller RJ, Bodine DM, Betschart B, Sager H. Distribution patterns of three sodium channel mutations associated with pyrethroid resistance in *Rhipicephalus (Boophilus) microplus* populations from North and South America, South Africa and Australia. International Journal for Parasitology: Drugs and Drug Resistance. 2012; 2:216-224.
- Norval RAI, Sutherst RW, Jorgenson O, Gibson JD, Kerr JD. The effect of the brown ear-tick *Rhipicephalus appendiculatus* on the growth of Sanga and European breed protein *tipE* and toxin pharmacology. Journal of General Physiology. 1988; 110:119-133.
- Norval RAI, Meltzer MI, Burridge MJ. Distribution, economic importance and control measures for *Cowdria ruminantium*. Proceedings of a workshop held at ILRAD Nairobi. 1991, 12-27.
- Vega M. Current importance of cattle haemoparasite diseases. In Seminario Internacional de Parasitología Animal. Garrapatas y enfermedades que transmiten. Morelos, México. 1991; 144-150.
- Peter TF, Barbet AF, Alleman AR, Simbi BH, Burridge MJ, Mahan SM. Detection of the Agent of Heartwater, *Cowdria ruminantium*, in *Amblyomma* Ticks by PCR: Validation and Application of the Assay to Field Ticks. Journal of Clinical Microbiology. 2000; 38:1539-1544.
- Farougou S, Adakal H, Biguezoton AS, Boko C. Prévalence de l'infection d'*Amblyomma variegatum* par *Ehrlichia ruminantium* dans les élevages extensifs du Bénin. Revue de Médecine Vétérinaire. 2012; 163 (5):261-266.
- Mendes MC, Duarte FC, Martins JR, Klafke GM, Fiorini LC, Barros ATM. Characterization of the pyrethroid resistance profile of *Rhipicephalus (Boophilus) microplus* populations from the states of Rio Grande do Sul and Mato Grosso do Sul, Brazil. Revista Brasileira de Parasitologia Veterinária. 2013; 22:379-384.
- Waltisbuhl D, Farnsworth B, Kemp D. Pesticides used against ectoparasites ticks, flies, lice of cattle in Australia. In P. A. Holdsworth (ed.), Ectoparasiticide use in contemporary Australian livestock production. Avcare Limited, Canberra. 2005, 45-57.
- Graf IF, Gogolewski R, Leach-Bing N, Sabatini GA, Molento MB, Bordin EL *et al.* Tick control: an industry point of view Parasitology. 2004; 129:427-442.
- Andreotti R, Guerrero FD, Soares MA, Barros JC, Miller RJ, Leon AP. Acaricide resistance of *Rhipicephalus (Boophilus) microplus* in State of Mato Grosso does Sul,

- Brazil. *Revista Brasileira de Parasitologia Veterinária*. 2011; 20:127-133.
17. Mendes MC, Lima CK, Nogueira AH, Yoshihara E, Chiebao DP. Resistance to cypermethrin, deltamethrin and chlorpyrifos in populations of 83 *Rhipicephalus (Boophilus) microplus* Acari: Ixodidae from small farms of the State of São Paulo. *Veterinary Parasitology*. 2011; 178:383-388.
  18. Domingues LN, Bello ACPP, Cunha AP, Brasil BSAF, Barros ATM, Leite RC *et al.* Survey of pyrethroid and 603 organophosphate resistance in Brazilian field populations of *Rhipicephalus (Boophilus) microplus*: detection of C190A mutation in domain II of the para-605 type sodium gene. *Veterinary Parasitology*. 2012; 189(2-4):327-332.
  19. Guerrero FD, Miller RJ, Perez de Leon AA. Cattle tick vaccines: Many candidate antigens, but will a commercially viable product emerge. *International Journal for Parasitology*. 2012; 42:421-427.
  20. Kearney. Acaricide resistance in cattle ticks. Northern Territory Government. 2011; 58:1-4.
  21. Lovis L. Evaluation of acaricide resistance in the cattle tick, *Rhipicephalus (Boophilus) microplus* using a new *in vitro* test and molecular tools. Doctorat University Neuchatel. 2012, 204.
  22. Dong K. Insect sodium channels and insecticide resistance. *Invertebrate Neuroscience*. 2007; 7:17-30.
  23. Lombet A, Mourre C, Ladzunski M. Interaction of insecticides of the pyrethrinoid family with specific binding sites on the voltage dependent sodium channel from mammalian. *Brain Research*. 1988; 459:44-48.
  24. FAO Resistance management and integrated parasite control in ruminants: guidelines 31 In: FAO, Module I Ticks: Acaricide Resistance: Diagnosis, Management and Prevention. Rome, Italy. 2004, 25-77.
  25. WHO-World Health Organization. Expert Committee on Insecticide Resistance of Vectors and Reservoirs of Diseases to Pesticides. Criteria and meaning of tests for determining the susceptibility or resistance, unpublished document, annex of 585 WHO technical, report series. 1976, 81- 6.
  26. Corley SW, Jonsson NN, Piper EK, Cutullé C, Stear MJ, Seddon JM. Mutation in the Rm  $\beta$ AOR gene is associated with amitraz resistance in the cattle tick *Rhipicephalus microplus*. *Proceedings of the National Academy of Sciences*. 2013; 110(42):772-7.
  27. Meyer JM, Ejendal KF, Avramova LV, Garland-Kuntz EE, Giraldo-Calderon GI, Brust TF *et al.* A genome-to-lead approach for insecticide discovery: pharmacological characterization and screening of *Aedes aegypti* D (1)-like dopamine receptors. *PLOS Neglected Tropical Diseases*. 2012; 6:1478.
  28. Bellgard MI, Moolhuijzen PM, Guerrero FD, Schibeci D, Rodriguez-Valle M, Peterson DG. Cattle Tick Base: Une ressource de bioinformatique intégrée basée sur Internet pour *Rhipicephalus (Boophilus) microplus*. *International Journal for Parasitology*. 2012; 42(2):161-169.
  29. Vaughan A, Hemingway J. Mosquito carboxylesterase EST alpha 2(1) (A2). Cloning and sequence of the full-length cDNA for a major insecticide resistance gene worldwide in the mosquito *Culex quinquefasciatus*. *Journal of Biological Chemistry*. 1995; 270:44-49.
  30. Fournier D, Bride JM, Mouches C, Raymond M, Magnin M, Berge JB *et al.* Biochemical characterization of the esterase A1 and B1 associated with organophosphate resistance in the *Culex pipiens* L. complex. *Pesticide Biochemistry and Physiology*. 1987; 27:211-217.
  31. David JP, Strode C, Vontas J, Nikou D, Vaughan A, Pignatelli PM *et al.* The Anopheles gambiae detoxification chip: a highly specific microarray to study metabolic-based insecticide resistance in malaria vectors. *Proceedings of the National Academy of Sciences*. 2005; 102: 4080-4.
  32. Hernandez R, He H, Chen AC, Waghela SD, Ivie GW, George JE *et al.* Identification of a point mutation in an esterase gene in different populations of the southern cattle tick, *Boophilus microplus*. *Insect Biochemistry and Molecular Biology*. 2000; 30:969-977.
  33. Hemingway J, Hawkes NJ, Mccarroll L, Ranson H. The molecular basis of insecticide resistance in mosquitoes. *Insect Biochemistry and Molecular Biology*. 2004; 34:653-65.
  34. Mendes MC, Duarte FC, Martins JR, Klafke GM, Fiorini LC, Barros ATM. Characterization of the pyrethroid resistance profile of *Rhipicephalus (Boophilus) microplus* populations from the states of Rio Grande do Sul and Mato Grosso do Sul, Brazil. *Revista Brasileira de Parasitologia Veterinária*. 2013; 22:379-384.
  35. FAO. Practical considerations of the use of resistant breeds of cattle for improved control of ticks in Latin America. *Animal Production and Health*. 1989; 75:165-176.
  36. Lovis LD, Perret JL, Bouvier J, Fellay JM, Kaminsky R, Betschart B *et al.* A new *in vitro* test to evaluate the resistance level against acaricides of the cattle tick *Rhipicephalus microplus*. *Veterinary Parasitology*. 2011; 182:269-280.
  37. Nolan J, Wilson JT, Green PE, Bird PE. Synthetic pyrethroid resistance in field samples in the cattle tick (*Boophilus microplus*). *Australian Veterinary Journal*. 1989; 66:179-182.
  38. Leite RC. *Boophilus microplus* susceptibilidade, uso atual retrospectivo de carrapaticidas em propriedades das regiões fisiográficas da Baixada do Grande Rio de Janeiro: uma abordagem epidemiológica. PhD thesis. Universidade Federal Rural do Rio de Janeiro. 1988.
  39. Santamaria EM, Frago SH. Resistencia en garrapatas *Boophilus microplus*, a los ixodididas en Mexico. XIV Pan American Congress of Veterinary Science. 1994, 473-474.
  40. Jonsson NN, Hope M. Progress in the epidemiology and diagnosis of amitraz resistance in the cattle tick *Boophilus microplus*. *Veterinary Parasitology*. 2007; 146:193-198.
  41. Guerrero FD, Davey RB, Miller RJ. Use of an allele-specific polymerase chain reaction assay to genotype pyrethroid resistant strains of *Boophilus microplus* (Acari: Ixodidae). *Journal of Medical Entomology*. 2001; 38:44-50.
  42. He H, Chen AC, Davey RB, Wayne IG, George JE. Characterization and molecular cloning of a glutathione S-transferase gene from the from the tick, *Boophilus microplus* (Acari: Ixodidae). *Insect Biochemistry and Molecular Biology*. 1999; 29:737-743.
  43. Jonsson NN, Cutullé C, Corley SW, Seddon LM. Identification of a mutation in the para-sodium channel gene of the cattle tick *Rhipicephalus microplus* associated with resistance to flumethrin but not to cypermethrin. *International Journal for Parasitology*. 2010; 40:1659-1664.

44. Morgan JAT, Corley SW, Jackson LA, Lew-Tabor AL, Moolhuijzen PM, Jonsson NN. Identification of a point mutation in the *para*-sodium channel gene of the cattle tick *Rhipicephalus (Boophilus) microplus* associated with resistance to synthetic pyrethroid acaricides. *International Journal for Parasitology*. 2009; 39(7):775-779.
45. Stone NE, Olafson PU, Davey RB, Buckmeier G, Bodine D. Multiple mutations in the *para*-sodium channel gene are associated with pyrethroid resistance in *Rhipicephalus microplus* from the United States and Mexico. *Parasites & Vectors*. 2014; 7:456.
46. Rodriguez-Vivas RI, Rivas AL, Chowell G, Fragoso SH, Rosario CR, Garcia Z *et al.* Spatial distribution of acaricide profiles (*Boophilus microplus* strains susceptible or resistant to acaricides) in southeastern Mexico. *Veterinary Parasitology*. 2007; 146:158-169.
47. Angus BM. The history of the cattle tick *Boophilus microplus* in Australia and achievements in its control. *International Journal for Parasitology*. 1996; 26:1341-1355.
48. Mitchell M. Acaricide resistance back to basics. *Tropical Animal Health and Production*. 1996; 28:53-58.
49. Legg I. Recent developments in tick and buffalo-fly control. *Australian Veterinary Journal*. 1947; 23:181-185.
50. Freire JJ. Arseno e cloro resistência emprego de tiosfato de dietilparanitrofenila (Parathion) na luta anticarrapato *Boophilus microplus*. *Boletim Produção Animal*. 1953; 9:3-31.
51. Cremllyn R. *Pesticides Preparation and mode of action*. John Wiley and Sons, Hoboken. 1979; 114-115.
52. Roulston WJ, Stone BF, Wilson JT, White LI. Chemical control of an organophosphates and carbamate-resistant strain of *Boophilus microplus* (Can.) from Queensland. *Bulletin of Entomological Research*. 1968; 58:379-392.
53. Miller RJ, Davey RB, George JE. First report of organophosphate-resistant *Boophilus microplus* (Acari: Ixodidae) within the United States. *Journal of Medical Entomology*. 2005; 42:912-917.
54. Nolan J. Current developments in resistance to amidine and pyrethroid tickicides in Australia: Tick biology and control. *Proceedings of an International Conference held from Rhodes University, South Africa*. 1981, 109-114.
55. Geary TG. Ivermectin 20 years on: maturation of a wonder drug. *Trends in Parasitology*. 2005; 21:530-532.
56. Taylor MA. Recent developments in ectoparasiticides. *The Veterinary Journal*. 2001; 161:253-268.
57. Martins JR. Furlong I AVermectin resistance of the cattle tick *Boophilus microplus* in Brazil. *Veterinary Record*. 2001, 149-164.
58. Pérez CLC, Rodríguez Vivas RI, Ramírez CGT, Miller RJ. First report of the cattle tick *Rhipicephalus microplus* resistant to ivermectin in Mexico. *Veterinary Parasitology*. 2010; 168:165-9.
59. Davey RB, Ahrens EH, George JE, Hunter JS, Jeannin P. Therapeutic and persistent efficacy of fipronil against *Boophilus microplus* (Acari: Ixodidae) on cattle. *Veterinary Parasitology*. 1998; 74:261-276.
60. Davey RB, George JE, Hunter JS, Jeannin P. Evaluation of a pour-on formulation of fipronil against *Boophilus annulatus* (Acari: Ixodidae) under natural South Texas field conditions. *Experimental and Applied Acarology*. 1999; 123:351-364.
61. Cuore U, Trelles A, Sanchis L, Gayo V, Solari MA. Primer diagnostico de resistencia al Fipronil en la garrapata comun del ganado *Boophilus microplus*. *Veterinaria*. 2007; 42:35-41.
62. Castro-Janer E, Martins JR, Mendes MC, Namindome A, Klafke GM, Schumaker TT. Diagnoses of fipronil resistance in Brazilian cattle ticks (*Rhipicephalus microplus*) using *in vitro* larval bioassays. *Veterinary Parasitology*. 2010; 173:300.
63. Jackson L, Stutchbury R. Fluazuron resistance in cattle ticks. 2010.
64. Cuisance D, Barré N, De deken R. Ectoparasites des animaux : méthodes de lutte écologique, biologique, génétique et mécanique. *Revue scientifique et technique (International Office of Epizootics)*. 1994; 13:05-13.
65. Barré N, Garrís G, Borel G, Camus E. Hosts and population dynamics of *Amblyomma variegatum* (Acari: Ixodidae) on Guadeloupe, French West Indies. *Journal of Medical Entomology*. 1988; 25(2):111-115.
66. Stachurski F. Modalités de la rencontre entre la stase adulte de la tique *Amblyomma variegatum* (acari, ixodida) et les bovins: applications potentielles à la lutte contre ce parasite. Thèse doctorale, parasitologie. Université Montpellier de Sciences et techniques du Languedoc. 2000, 264.
67. Barré N. Biologie et écologie de la tique *Amblyomma variegatum* (Acarina : Ixodina) en Guadeloupe (Antilles françaises). Thèse de Doctorat d'Etat. Université de Paris-Sud, Paris. 1989, 267.
68. Kaaya GP, Mwangi EN, Ouna EA. Prospects for biological control of livestock ticks, *Rhipicephalus appendiculatus* and *Amblyomma variegatum*, using the entomogenous fungi *Beauveria bassiana* and *Metarhizium anisopliae*. *Journal of Invertebrate Pathology*. 1996; 67:15-20.
69. Adinci KJ, Yessinou RE, Adehan SB, Yovo M, Ahounou S, Sessou *et al.* *In vitro* evaluation of the acaricidal effect of three essential oils extracted from aromatic local plants on *Rhipicephalus (Boophilus) microplus* ticks in Benin. *International Journal of Advanced Research*. 2015; 11:1514-1520.
70. Yessinou RE, Adinci J, Sessou P, Adehan S, Tonouweha A, Akpo Y *et al.* *In vitro* acaricidal effect of *Syzygium aromaticum* and *Cymbopogon citratus* essential oils on engorged female of cattle tick *Rhipicephalus microplus* in Benin. *Scientific Journal of Veterinary Advance*. 2016; 5:80-86.
71. Nawaz M, Sajid MS, Zulfiqar A, Muhamma W, Tanveer A, Abid H *et al.* Anti-Tick Activity of Leaves of *Azadirachta indica*, *Dalbergia sissoo* and *Morus alba* against *Rhipicephalus microplus* (Acari: Ixodidae). *Acta Parasitologica Globalis*. 2015; 6:60-64.
72. Amosti C, Fuchs BM, Amann R, Passow U. Contrasting extracellular enzyme activities of particle associated bacteria from distinct provinces of the North Atlantic Ocean. *Frontiers in Microbiology*. 2012; 3:425.
73. Dougnon TJ, Adéhan S, Houessionon J, Farougou S. *In vitro* affect of the ethanolic extract of *Tephrosia Vogelii* on *Rhipicephalus Sanguineus* in Abomey-Calavi, Avicenna *Journal of Phytomedicine*. 2015; 5(3):247-259.
74. Gadzirayi CT, Gadzirayi E, Mutandwa M, Mwale T, Chindundu. Utilization of *Tephrosia vogelii* in controlling ticks in dairy cows by small-scale commercial farmers in Zimbabwe. *African Journal of Biotechnology*. 2009; 8:4134-4136.
75. Kalume MK, Losson B, Angenot L, Titsch M, Wauters JN, Frédéric M *et al.* Rotenoid content and *in vitro* acaricidal activity of *Tephrosia vogelii* leaf extract on the

- tick *Rhipicephalus appendiculatus*. Veterinary Parasitology - Journal. 2012; 06-015.
76. Kumar A, Singh S, Mahour K, Vihan VS. *In vitro* and *in Vivo* Acaricidal Activity of Some Indigenous Plants Under Organized and Farmer Flock. Pharmacology online. 2011; 3:361-369.
  77. Vihan VS, Agarwal A. Development of a novel herbal ectoparasitological product under DST fund. Project report. 2007.
  78. Sutherst RW, Utech KBW. Controlling livestock parasites with host resistance. CRC Handbook of Pest Management in Agriculture. 1981; 2:385-40.
  79. Piper EK, Jackson LA, Bagnall NH, Kongsuwan KK, Lew AE, Jonsson NN. Gene expression in the skin of *Bos taurus* and *Bos indicus* cattle infested with the cattle tick, *Rhipicephalus (Boophilus) microplus*. Veterinary Immunology and Immunopathology. 2008; 126:110-119.
  80. Brossard M. The use of vaccines and genetically resistant animals in tick control. *Revue scientifique et technique (International Office of Epizootics)*. 1998; 17:188-199.
  81. Uly M. Médecine Vétérinaire naturelle: lutte contre les ectoparasites tropicaux. Weikersheim: Scientific book, V. J. Margraf, Allemagne, 1993, 183.
  82. Lafia S. Les tiques (Amblyommidae) parasites des bovines en République Populaire du Bénin. Thèse Médecine Vétérinaire Dakar. 1982, 83.
  83. Pérez Cogollo LC, Rodríguez Vivas RI, Ramírez Cruz GT, Rosado-Aguilar JA. Survey of *Rhipicephalus microplus* resistance to ivermectin at cattle farms with history of macrocyclic lactones use in Yucatan, Mexico. Veterinary Parasitology. 2010; 172:109-113.
  84. Maranga RO, Kaaya GP, Mueke JM, Hassanali A. Effects of combining the fungi *Beauveria bassiana* and *Metarhizium anisopliae* on the mortality of the tick *Amblyomma variegatum* (Ixodidae) in relation to seasonal changes. Mycopathologia. 2005; 159:527-532.
  85. Barré NH, Mauléon GI, Garris A, Kermarrec Predators of the tick *Amblyomma variegatum* (Acari: Ixodidae) in Guadeloupe, French West Indies. Experimental & Applied Acarology. 1991; 12:163-170.
  86. Mattioli RC, Bah M, Faye J, Kora S, Cassama AM. Comparison of field tick infestation on N'dama, Zebu and N'dama-Zebu crossed cattle. Veterinary Parasitology. 1993; 47:139-148.
  87. Piper EK, Jackson LA, Bielefeldt-Ohmann H, Gondro C, Lew-Tabor AE, Jonsson NN. Tick-susceptible *Bos taurus* cattle display an increased cellular response at the site of larval *Rhipicephalus (Boophilus) microplus* attachment, compared with tick-resistant *Bos indicus* cattle. International Journal for Parasitology. 2010; 40:431-441.
  88. Ibelli AM, Ribeiro AR, Giglioti R, Regitano LC, Alencar MM, Chagas AC *et al.* Resistance of cattle of various genetic groups to the tick *Rhipicephalus microplus* and the relationship with coat traits. Veterinary Parasitology. 2012; 186:425-30.