



Research article

Digestibility of feeds in rabbit (*Oryctolagus cuniculus* linnaeus, 1758) in Africa: A review

G.S.T. Atchade^{1*}, M.F. Houndonougbo², S.E. Attakpa³, C.A.A.M. Chrysostome² and G. A. Mensah¹

¹Laboratoire de Recherches Zootechnique Vétérinaire et Halieutique, Institut National des Recherches Agricoles du Bénin (INRAB), ²Laboratoire de Recherche Avicole et de Zoo-Economie, Département de Production Animale, Faculté des Sciences Agronomiques de l'Université d'Abomey-Calavi, Bénin. ³Laboratoire de Physiopathologie Moléculaire et Toxicologie, Département de physiologie animale, Faculté des Sciences et Techniques de l'Université d'Abomey-calavi, Bénin

<p>Article history Received: Dec 22, 2018 Revised: Mar 02, 2019 Accepted: Mar 09, 2019</p>	<p>Abstract Feed digestibility indicates how much the body uses feed and its nutrients. This study aims to make a literature review on the digestibility of balanced feed given to farmed rabbits in Africa. Data used for this literature review were collected through many scientific papers published from 2008 to 2018 and uploaded to Google Scholar, Pubmed, Agora search engines and in national university libraries. Numerous scientific studies in this field indicate that more than thirty feedstuffs are commonly used to formulate balanced feed delivered to rabbits in Africa. Feed digestibility is generally assessed <i>in vivo</i> by the direct method in farmed rabbits. The data compiled shows that the digestibility of feed by rabbits varies enormously depending on the feedstuffs in balanced feed and also on their incorporation rate. The data recorded by the various authors' show that balanced feeds with an Apparent Digestibility greater than 50%. Keywords: Balanced feed, In vivo digestibility, Farmed rabbit, Africa</p>
--	--

To cite this article: Atchade GST, MF Houndonougbo, SE Attakpa, CAAM Chrysostome and GA Mensah, 2019. Digestibility of feeds in rabbit (*Oryctolagus cuniculus* linnaeus, 1758) in Africa: A review. Res. Opin. Anim. Vet. Sci., 9(1): 1-7.

Introduction

To meet the animal protein requirements of West African populations, emphasis has been placed on the development of production and the health of short-cycle animals (Gbangboché et al., 2005), including rabbit breeding (*Oryctolagus cuniculus* linnaeus, 1758). Indeed, rabbits breeding is a breeding that generates a lot of income in a short time (Lebas, 2007). Its meat has exceptional qualities that distinguish it from others meats. Its flesh is tasty and resembles that of chicken. It is digestible, rich in water, proteins, essential amino acids, iron and

vitamins. It is low in fat and sodium and has a good index of initiation (Lebas, 2007). In traditional and rational breeding, balanced feed including crop and by-products, minerals and vitamins are used to feed farmed rabbit. Thus, balanced feed delivered to rabbits may vary depending on the available feedstuffs. However, the balanced feed delivered can be digested differently by rabbits. "Feed digestibility" indicates how much the body uses feed or nutrients (Deiltel et al., 2012). The present article summarizes results of different feeding studies on the digestibility of balanced feed with both conventional and unconventional feedstuffs in farmed rabbits.

***Corresponding author:** Laboratoire de Recherche Avicole et de Zoo-Economie, Département de Production Animale, Faculté des Sciences Agronomiques de l'Université d'Abomey-Calavi, Bénin, 01BP 3783 Ganhi Cotonou, Bénin. fredericmh@gmail.com

Materials and Methods

Data used for this literature review were collected and processed mainly through:

- A literature search that collected information and data on feed digestibility in farmed rabbits in Africa, mainly from twenty-nine (29) scientific papers published from 2008 to 2018 and uploaded to Google Scholar, Pubmed, Agora search engines and in national university libraries.
- A bibliographic analysis which consisted of the manual counting, the synthesis of the information and data collected and the compilation of feed digestibility's recorded in the farmed rabbit by the different authors.

Results and Discussion

Concentrated feedstuffs in rabbits balanced feed in Africa

The rich tropical biodiversity, the great diversity of agricultural production systems in African countries in general and the presence of small agro-processing units contribute to a wide range of feedstuffs that can be used by animals (Archimède et al., 2011). The diet of farmed chicken and rabbit generally involves two main types of concentrated feedstuffs (CFS): cereals and by-products. These CFS include crop products; agro-industrial food and animal products and by-products. Their use depends mainly on their availability in the region, the season, the price and their ability to be kept for a long time (Toléba et al., 2007). Thirty-three (33) different CFS used in the feeding of farmed rabbits in Africa and listed in this literature review are: Barley, Berseem hay, Brewers dried grain, Burukutu spent grain, Cassava root meal, Cocoa shell, Copra meal cake, Corn bran, Corn, Cotton meal, Cotton waste cake, Cottonseed cake, Full-fat-soyabean, Fish meal, Groudnut cake, Ground haulms, Groundnut haulms, Groundnut husk, Maize bran, Maize grains, Maize offal, Maize, Millet mash residue, Molasses, Palm kernel cake, Rice bran, Rice husk, Rice straw, Roasted full fat soyabean, Soya bean, Soyabean meal, Wheat bran, Wheat flour, Wheat offal, Whole soyabean, Yellow maize.

Moreover, in the extensive production, the diet of rabbit is also based on the use of some vitamins, minerals amino acids and pigments (N'gom 2004, Guindjombi 2007, Kpodékon et al., 2009). In this category, twenty-one (21) feedstuffs are commonly used in Africa. These feedstuffs are: Bi / Di Calcium Phosphate, Calcium Carbonate Dibasic Calcium, Limestone, DL-Methionine, Lysine, Premix, Vitamin premix, Concentrate flesh, Iron sulfate, Palm oil, Sawdust, Sugar, Blood meal, Bone ash, Bone meal, Common salt, Iodized salt, Oyster shell, Oxyboldin, PCP, PKC.

Form of balanced feed used rabbit rearing

Pellet is the form of the balanced feed highlighted in almost all of the research work considered in this

literature review. Indeed, according to Lebas et al., (2007), the advantage of pellet is that it is manufactured according to the specific needs of the animal and it cannot sort, consumes exactly diet planned for him. Also, the rabbit prefers a pellet to a meal feed.

Feed digestibility in rabbits and balanced experiment

The feed ingested by rabbit is not entirely used by his body; part of "ingestats" passes through the digestive tract and is found in the faeces. The digestible fraction of a feed is the part of "ingestats" that is not found in the faeces. It is a quantitative concept that results in the Digestive Utilization Coefficient (D.U.C) or digestibility coefficient (D.C). In other words, it is the proportion of the various constituents of a feed that is retained by body. Digestibility is therefore an index of the disappearance of the feed in the intestine, but not of the efficiency of use of the feed by the metabolism of the animal. Thus, a feed can be very digestible without providing enough nutrients to meet the requirement of the animal.

Most studies on rabbits in Africa focus on feed apparent digestibility, limiting themselves to the balance between the nutrients of ingestats and excretas. Formula used to calculate the Apparent Digestive Utilization Coefficient is as follows:

$$\text{Apparent digestibility} = [(\% \text{Nutrients in feed} \times \text{FI}) - (\% \text{Nutrients in faeces} \times \text{FO})] / \% \text{Nutrients in feed} \times \text{FI} \times 100;$$

Where FI= feed intake and FO= faecal output Apparent digestibility determination in rabbit

Balanced experiment in rabbit feeding

A typical digestibility experiment is conducted over a period of seven (07) to ten (10) days by feeding daily experimental rabbit. To the experimental period are added stages of droppings drying, grinding and analysis (Bourdillon et al., 1990). Experiment on the assessment of feed digestibility in farmed rabbits is carried out in digestibility cages equipped with a faeces collection device, on rabbits that age and sex are known. This device makes it possible not to mix the faeces with the urine of rabbits. Rabbits are individually arranged per digestibility cage and data is collected daily by rabbit. Thus, the quantities of feed delivered and refused as well as those of faeces are weighed and recorded daily.

Determination of Apparent Digestibility in rabbit

According to Rivière, 1991, feed digestibility can be determined from three (03) groups of methods: (i) In vivo methods; (ii) in vitro methods (laboratory method); (iii) Mathematical or prediction methods. In rabbits, direct method *in vivo* is preferably used for

assessing digestibility. It is based on faecal digestibility measurements, the method of which was standardized in 1995 by the EGRAN group (Guermah, 2012). Representative sampling is used because slight differences can be recorded in digestive function in rabbit of the same breed, age and sex. Repetitions allow the detection of errors (Rivière, 1991).

Feed samples as well as faeces samples are dried and used in laboratory to determine their chemical composition according to the methods recommended by AOAC (1991); AOAC (1997). The chemical composition includes nutrients such as Dry Matter (DM), Organic Matter (OM), Crude Protein (CP), Fatty Matter (EE), and Crude Fiber (CF).

After evaluation of the chemical composition of samples of feed served, refused and faeces, Apparent Digestibility of the experimental diet in its nutrients is calculated by its formula.

Apparent digestibility of balanced feed in rabbit

Several authors in Africa studied feed digestibility in rabbits fed with balanced diets based on conventional

and unconventional concentrated feedstuffs. Table 1 presents the digestibility of balanced feeds in farmed rabbits in Africa. The apparent digestibility varied according to incorporation rate of the feedstuff in study. Indeed, most of these digestibilities have been evaluated following the use of a feedstuff at different levels of incorporation into a basic diet formula. Majority of nutrients in balanced feeds delivered to farmed rabbits in Africa are well digested. Digestibility Coefficient of the dry matter in balanced feed varied greatly from a study to another: the highest are above 50% and range from 53 to 99% while the lowest are between 39 and 48%. The digestibility of crude protein, varies from 55 to 97%. However, a very low protein digestibility around 29% was recorded in studies. The digestibility coefficient of organic matter in balanced feed is above 40% and reaches 85%. The fat digestibility varies from 57% to 93%. Crude fiber that are essential for the proper functioning of the rabbit's digestive system are digested at a rate ranging from 50 to 97%. However, some low digestibilities of crude fiber ranging from 24 to 49% are recorded.

Table 1: Apparent digestibility of some balanced feeds in rabbit in Africa

Experimental Feedstuffs (Rate in the feed)	Feed composition	DDM (%)	DCP (%)	DOM (%)	DEE (%)	DCF (%)	Authors (Country)
Lablab seed (0; 19,92; 23,01; 20,59; 23,43 kg/100 kg)	Lablab seed (raw, decorticated, toasted), Maize, Full-fat-soyabean, Rice husk, Brewers dried grain, Bone ash, Salt, Premix)	61.68 – 80.82	67.38 – 78.92	ND	44.64 – 81.06	24.02 – 62.08	Shaahu et al., 2014 (Nigéria)
<i>Acacia saligna</i> , <i>Leuceana</i> , <i>leucocephala</i> , <i>Moringa. Oleifera</i> dried foliages (15 kg/100 kg)	<i>Acacia saligna</i> , <i>Leuceana</i> , <i>leucocephala</i> , <i>Moringa oleifera</i> dried foliages, Barley, Soybean meal, Wheat bran, Berseem hay, Molasses, Calcium carbonate, Dicalcium phosphate, Salt, Premix, DL-méthionine)	53.0 - 61.8	56.2 - 64.7	56.2 - 64.4	76.3 - 82.3	36.5 - 44.9	Abu Hafsa et al., 2016 (Egypte)
Wild sunflower leaf meal (0; 5; 10; 15 kg/100 kg)	Wild sunflower leaf meal, Maize, Corn bran, Groundnut husk, Palm kernel cake, Fish meal, Bone meal, Premix, Salt	44.30- 60.46	29.4- 55.0	ND	9.70- 10.89	8.33- 12.28	Adam, 2013 (Ghana)
Vitamine E, Vitamine C (0; 200; 400 mg/100 kg)	Vitamine E, Vitamine C, Maize, Wheat offal, Soya bean, PKC, Fish meal, Groundnut cake, Bone meal, Iodized salt, Vit-mineral mix	98.17- 99.33	43.73- 55.03	ND	65.29- 95.43	63.10- 75.13	Okachi and Ani, 2016 (Nigeria)
Pineapple peels (0; 10; 20; 30 kg/100 kg)	Pineapple peels, Maize, cottonseed cake, soybean meal, sawdust, wheat bran, Salt, Oyster shell, Panicum maximum	62- 67.7	ND	ND	ND	ND	Aboh et al., 2013 (Bénin)
Pellets of <i>Moringa oleifera</i> (0; 10; 15 kg/100 kg)	Pellets of <i>Moringa oleifera</i> , Maize, Palm kernel cake, Soybean meal, Rice bran, Oyster, shell, Sawdust, Salt	52.3- 57.07	66.12- 74.97	54.26- 61.25	ND	ND	Doungnon et al., 2012 (Bénin)
Sundried <i>Tridax procumbens</i> (0; 5; 10; 15; 20 kg/100 kg)	Sundried <i>Tridax procumbens</i> , Cassava root meal, Roasted full fat soyabean, Burukutu spent grain, Rice husk, Bone meal, Palm oil, Salt, Sugar, Premix, Coccidiostat	79.68- 82.79	70.76- 81.79	ND	71.47- 87.11	39.33- 43.65	Anthony et al., 2016 (Nigéria)

Experimental Feedstuffs (Rate in the feed)	Feed composition	DDM (%)	DCP (%)	DOM (%)	DEE (%)	DCF (%)	Authors (Country)
Sundried Cocoa Pod Husk Meal (0; 12,50; 25; 37,50 kg/100 kg)	Sundried Cocoa Pod Husk Meal, Yellow maize, Soybean, Fish meal, PKC, Wheat offal, Rice husk, Bone meal, Palm oil, Methionine, Lysine, Salt, Premix	95.41-96.43	75.07-78.54	ND	71.51-83.57	74.50-78.44	Ozung et al., 2017 (Nigeria)
Hot Cocoa Pod Husk Meal (0; 12,50; 25; 37,50 kg/100 kg)	Hot Cocoa Pod Husk Meal, Yellow maize, Soybean, Fish meal, PKC, Wheat offal, Rice husk, Bone meal, Palm oil, Methionine, Lysine, Salt, Premix	96.27-96.43	73.31-77.74	ND	74.77-82.03	77.42-85.97	Ozung et al., 2017 (Nigeria)
Fermented Cocoa Pod Husk Meal (0; 12,50; 25; 37,50 kg/100 kg)	Fermented Cocoa Pod Husk Meal, Yellow maize, Soybean, Fish meal, PKC, Wheat offal, Rice husk, Bone meal, Palm oil, Methionine, Lysine, Salt, Premix	96.33-96.58	72.18-78.54	ND	76.26-82.39	61.14-76.79	Ozung et al., 2017 (Nigeria)
Sorghum "Chakalere" Sorghum "Jigare" (0; 17; 34 kg/100 kg)	Sorghum "Chakalere", Sorghum "Jigare", Maize, Wheat offal, Groundnut cake, Fish meal, Groundnut haulms, Limestone, Salt, Premix	66.47-79.48	69.28-84.47	ND	57.97-74.81	30.37-42.88	Igwebuiké et al., 2013 (Nigeria)
Breadfruit meal (0; 9; 12; 15; 18; 21 kg/100 kg)	Breadfruit meal, Maize, Soybean meal, Wheat offal, Corn bran, Palm kernel meal, Fish meal, Rice husk, Bone meal, Salt, Vit/Premix	60.10-63.50	67.10-70.80	ND	70.25-74.23	49.32-55.10	Oladunjoye and Ojebiyi, 2011 (Nigeria)
Sundried <i>Digitaria iburua</i> (0; 20; 40 kg/100 kg)	Sundried <i>Digitaria iburua</i> , Maize, Soybean meal, Fish meal, Wheat offal, Bone meal, Premix Salt	72.99-76.33	73.22-75.05	ND	ND	64.04-67.35	Oke et al., 2016 (Nigeria)
Raw Kapok seed meal (0; 5; 10; 15; 20 kg/100 kg)	Raw Kapok seed meal, Maize, Soybean meal, Fish meal, Maize offal, Wheat offal, Methionine, Lysine, Salt, Premix	60.45-88.89	60.45-84.84	ND	57.27-74.47	58.14-76.09	Wafar et al., 2017 (Nigeria)
Sundried Sorghum offal (0; 11,94; 24,60;35,62;47,56 kg/100 kg)	Sundried Sorghum offal, Maize, Groundnut cake, Wheat offal, Rice bran, Bone meal, Blood meal, Premix, Lysine, Methionine, Salt	60.14-74.34	58.54-76.83	ND	66.73-76.39	28.14-39.31	Ogunsipe et al., 2014 (Nigeria)
Cassava peel residue (0; 10, 25; 20,5; 30,75, 41 kg/100 kg)	Cassava peel residue, Maize, Soybean meal, Palm kernel cake, Wheat offal, Bone meal, Limestone, Premix, Salt	56.2-65.9	54.2-65.7	ND	79.9-93.6	78.6-90.3	Osakwe et al., 2008 (Nigeria)
Cowpea pods (0; 5; 10; 15 kg/100 kg)	Cowpea pods, Maize grains, Wheat bran, Soybean meal, Cotton meal, Palm-Kernel meal, Oyster shell, Lysine, Methionine; PCP, Salt, Premix, Iron Sulphate	78.78-83.57	55.47-60.57	81.47-84.50	ND	ND	Kpomasse et al., 2015 (Bénin)
Cowpea pod shell (0; 10 kg/100 kg)	Cowpea pod shell, Soybean pod shell, Maize, Wheat bran, Soybean meal, Cotton meal, Palm meal, Oyster shell, Lysine, Methionine, Phosphoric acid, Salt, Concentrate flesh, Iron sulfate	39.02-52.70	ND	ND	ND	ND	Koura et al., 2015 (Bénin)
Soybean pod shell (0, 10 kg/100 kg)	Cowpea pod shell, Soybean pod shell, Maize, Wheat bran, Soybean meal, Cotton meal, Palm meal, Oyster shell, Lysine, Methionine, Phosphoric acid, Salt, Concentrate flesh, Iron sulfate	39.02-52.70	ND	ND	ND	ND	Koura et al., 2015 (Bénin)
Sundried breadfruit meal (0; 10; 20; 30 kg/100 kg)	Sundried breadfruit meal, Maize, Soybean, Wheat offal, Fish meal, Palm Kernel cake, Bone meal, Oyster shell, Premix, Salt	71.20-83.98	70.24-79.79	ND	68.10-74.44	57.50-66.41	Oladele-Oso et al., 2010 (Nigeria)
Raw baobab seed meal (0; 10; 20; 30 kg/100 kg)	Raw baobab seed meal, Maize, Soybean, Ground haulms, Wheat offal, Fishmeal, Bone meal, Premix, Salt, Methionine	55.40-67.20	55.39-64.37	ND	69.03-70.54	55.94-67.77	Abdullahi et al., 2017 (Nigeria)

Experimental Feedstuffs (Rate in the feed)	Feed composition	DDM (%)	DCP (%)	DOM (%)	DEE (%)	DCF (%)	Authors (Country)
Dried <i>Acacia albida</i> pods (10; 20; 30; 40 kg/100 kg)	Dried <i>Acacia albida</i> pods, Maize, Maize bran, Groundnut haulms, Groudnut cake, Palm oil, Bone meal, Salt, Premix	68.16-79.49	71.06-80.38	ND	82.09-84.34	28.40-50.29	Igwebuike et al., 2008 (Nigéria)
Millet residue meal (83; 74; 76 kg/100 kg)	Millet residue meal, Fish meal, Soybean meal, DCP, Vitamin/mineral premix, Common salt	70.8-82.7	82.5-90.7	ND	ND	ND	Karikari et al., 2011 (Ghana)
Soybean pod (0; 5; 10; 15 kg/100 kg)	Soybean pod, Maize, Wheat bran, Soybean meal, Cotton meal, palm Kernel meal, Oyster shell, Lysine, Methionine, Bi-calcium Phosphate, Salt, Premix, Iron sulphate	77.62-85.99	63.20-72.23	77.74-86.07	ND	ND	Badet et al., 2015 (Bénin)
Cottonseed cake (0; 5; 10; 15 kg/100 kg)	Cottonseed cake, Maize, Groundnut cake, Rice husk, Fishmeal, Bonemeal, Oyster shell, Vitamin/mineral premix, Salt	69.92-75.92	83.52-86.30	ND	67.96-90.24	87.25-91.82	Amao et al., 2012 (Nigéria)
<i>Prosopis africana</i> pulp (0; 10; 20; 30, 40%)	<i>Prosopis africana</i> pulp, Maize and conventional ingredient	59.08-66.95	72.26-79.32	ND	61.11-70.27	34.49-38.41	Adamu et al., 2011 (Nigéria)
<i>Moringa</i> leaf meal (0; 5; 10; 15; 20 kg/100 kg)	<i>Moringa</i> leaf meal, Millet mash residue, Soybean meal, DCP, Vitamin-mineral premix, Salt	65.02-78.40	65.10-87.80	ND	70.00-77.50	50.01-55	Nuhu, 2010 (Ghana)
<i>Moringa</i> powder (0; 1,50; 3,0 kg/100 kg)	<i>Moringa</i> powder, Corn, Wheat bran, Rice bran, Rice straw, Cocoa shell, Soya bean meal flour, Cotton waste cake, Copra meal cake, Shells of oysters, Methionine, Lysine, Dibasic calcium, Vitamin premix, Oxyboldine	ND	80.85-85.00	ND	67.00-68.70	ND	Djakalia et al., 2011 (Côte d'Ivoire)
<i>Moringa</i> leaf meal (0; 5; 10; 15; 20 kg/100 kg)	<i>Moringa</i> leaf meal, Maize, Wheat offal, Soya meal Palm kernel cake, Groundnut cake, DCP, Lysine, Methionine, Premix, Salt	56.10-80.7	65.13-89.2	ND	71.20-77.2	50.18-56.2	Olatunji et al., 2016 (Nigéria)
<i>Moringa</i> leaf meal (0; 1,50; 3,00; 4,50 kg/100 kg)	<i>Moringa</i> leaf meal, Maize, Soybean meal, Groundnut cake, Palm kernel cake, Wheat offal, Bone meal, Oyster shell, Salt, Vitamin-mineral premix, Lysine, Methionine	53.17-55.31	68.25-71.36	ND	80.80-83.31	74.61-78.87	Ayo-Ajasa et al., 2017 (Nigéria)
Leafy vegetables (0; 50 %)	Leafy vegetables, Commercial pelleted	65.49-87.86	86.73-96.22	ND	ND	ND	Yao Konan et al., 2016 (Côte d'Ivoire)
<i>Pachyrhizus. erosus</i> chips (0; 10; 20 kg/100 kg)	<i>Pachyrhizus erosus</i> chips, Maize, Rice bran, Wheat flour, Cottonseed cake, Soybean meal, Palm kernel meal, Oyster shell, Salt, Sawdust	48.13-61.58	64.17-69.01	42.12-58.34	ND	ND	Atchadé, 2012 (Bénin)
Maize bran (5,5; 11; 16,5 kg/100 kg)	Maize bran, Maize, Soybean meal, Cottonseed cake, Palm kernel meal, Shell, Lysine, Methionine, DCP, Nacl, Premix, Iron sulphate	60.18-67.23	93.34-97.14	ND	ND	91.94-97.06	Tchibozo et al., 2017 (Bénin)

DDM= Dry Matter Digestibility; DCP= Digestibility Crude Protein; DOM= Organic Matter Digestibility; DEE=: Ether Extract Digestibility; DCF= Crude Fiber Digestibility; ND= Not Determined.

Conclusion

A total of 33 scientific articles were used to provide information on balanced feed digestibility in farmed rabbits in Africa. The data demonstrate that the compound feeds delivered to farmed rabbits are often

digestible. This inventory will be useful to assess the digestibility of each concentrated feedstuff in farmed rabbits to develop a table of their nutritional composition in Benin.

References

- Abdullahi AI, Wafar, RJ, Yusuf SZ (2017) Evaluation of Raw Baobab Seed (*Adanosonia digitata*) Meal on the Performance of Weaned Rabbits. *Journal of Adv Biol Biotechnol* 12(2): 1-8.
- Aboh AB, Zoffoun GA, Djenontin AJP, Babatounde S, Mensah GA (2013) Effect of graded levels of dry pineapple peel on digestibility and growth performance of rabbit. *J Appl Biosci* 67: 5271-5276.
- Oladele-Oso A, Faboro R, Isah O, Oni A, Bamgbose A and Dele P (2010) Potential of bread fruit (*Artocarpus altilis*) an ecologically forest based feed resource in rabbit nutrition. *Trop Subtrop Agroecosystems* 12 (2010): 99-108.
- Abu Hafsa SH, Salem AZM, Hassan AA, Barbabosa A, Kholif AE, Lopez S, Elghandour MMY (2016) Digestion, growth performance and caecal fermentation in growing rabbits fed diets containing foliage of browse trees. *World Rabbit Sci* 2016 24: 283-293.
- Adam N (2013) Digestibility and growth performance of female rabbit fed *Tithonia diversifolia* leaf meal. Thesis of Kwame Nkrumah University of science and Technology, College of agriculture and natural resources, 61 p.
- Adamu L, Igwebuike JU, Kwari ID, Aliyu J (2011) The effects of *Prosopis africana* pulp on nutrient digestibility, carcass components and blood composition of growing rabbit. *J Environ Issues Agr Dev Count* 3: Number 1, April 2011.
- Amao OA, Adejumo DO, Togun VA, Oladunjoye IO (2012) Growth response and nutrient digestibility of prepubertal rabbit bucks fed cottonseed cake-based diets supplemented with vitamin E. *Afr J Biotechnol* 11(76): 14102-14109.
- Anthony TI, Ogwiji O, Oche DA, Wayo N, Sarwuan GD, Torhemen M and Idoga EE (2016) Nutrient intake and digestibility of Rabbits Fed different levels of *Tridax Procumbens* in Cassava – Based Rations. *Schol J Agr Sci* 6: 257-262.
- AOAC (1991) Official Methods of Analysis. Association of Official Agricultural Chemists, 15th Edition, Washington, DC, USA.
- AOAC (1997) Official Methods of Analysis. Association of Official Analytical Chemists, 16th Edition, Washington, DC, USA.
- Archimède H, Bastianelli D, Boval M, Tran G, Sauvart D (2011) Ressources tropicales : disponibilité et valeur alimentaire. *INRA Prod Anim* 24(1): 23-40.
- Atchadé GST (2012) Utilisation des racines tubéreuses de *Pachyrhizus erosus* dans l'aliment granulé des lapins en croissance au Bénin. Mémoire de Master. Faculté des Sciences et Techniques, Université d'Abomey-Clavi, Bénin, 81 p.
- Ayo-Ajasa OY, Egbeyale LT, Sanusi GO, Ibrahim K, Hamzat OA and Falako MF (2017) Performance and cost-benefits of weaner rabbits fed graded levels of *Moringa oleifera* leaf meal. *Mal J Anim Sci* 20(1): 59-68.
- Badet FGJ, Houndonougbo MF and Chrysostome CAAM (2015) The optimal rate of soybean (*Glycine max*) pods in rabbit (*Oryctolagus cuniculus*) feed: digestibility study and growth performances. *Res Opin Anim Vet Sci* 5: 406-410.
- Bourdillon A, Carre B, Conan L, Francesch M, Fuentes M, Huyghebaert G, Janssen WMMA, Leclercq B, Lessire M, McNab J, Rigoni M, Wiseman J (1990) European reference method of *in vivo* determination of metabolisable energy in poultry: reproducibility, effect of age, comparison with predicted values. *Brit Poult Sci* 31: 567-576.
- Deiltel L, Créchet C, Fornier E, Leborgne MC, Montméas L (2012) Nutrition et alimentation des animaux d'élevage. Tome I, 3^{ème} édition. Educagri Editions, Dijon.
- Djakalia B, Guichard BL and Soumaila D (2011) Effect of *Moringa oleifera* on Growth Performance and Health Status of Young Post-Weaning Rabbits. *Res J Poultry Sci* 4(1): 7-13.
- Dougnon TJ, Aboh BA, Kpodékon TM, Honvou S, Youssao I (2012) Effects of substitution of pellet of *Moringa oleifera* to commercial feed on rabbit's digestion, growth performance and carcass trait. *J App Pharm Sci* 2(9): 015-019.
- Gbangboché AB, Hornick J, Adamou-N'diaye, Edorh AP, Farnir F, Abiola FA, Leroy PL (2005) Caractérisation et maîtrise des paramètres de la reproduction et de la croissance des ovins Djallonké (*Ovis amon aries*). *Annales de la Médecine Vétérinaire* 149 : 148-160.
- Guermah H (2012) Nutrition du lapin : Etude de sources alimentaires alternatives. Thèse de l'Université Mouloud Mammeri de Tizi-Ouzou, Faculté des Sciences Biologiques et Sciences Agronomiques, 122 p.
- Guindjombi S (2007) Cuniculture périurbaine dans les Niayes : Situation actuelle et perspectives de développement. Thèse de doctorat en Médecine vétérinaire, Faculté de Médecine, de Pharmacie et d'Odonto-Stomatologie de Dakar, 117 p.
- Igwebuike JU, Anugwa FOI, Raji AO, Ehiobu NG and Ikurior SA (2008) Nutrient digestibility, haematological and serum biochemical indices of rabbits fed graded levels of *Acacia albida* pods. *ARPN J Agr Biol Sci* 3:33-40.
- Igwebuike JU, Medugu CI, Kwari ID & Dauda A (2013) Growth performance and nutrient digestibility of growing rabbits fed two varieties of sorghum as replacement for maize as energy source in tropical environment. *Global J Bio-Sci Biotechnol* 2: 167-172.

- Karikari PK, Darkoh G and Deku G (2011) Evaluation of millet residue meal based diets as feed for the domestic rabbit (*Oryctolagus cuniculus*). Res Opin Anim Vet Sci 1(3): 144-149.
- Koura BI, Houndonougbo F and Houinato M (2015) Effect of incorporation of cowpea and soybean pods in diets on feed intake, digestibility and weight gain performances of rabbit. Rev Cames 3: 32-36.
- Kpodékon TM, Youssao AKI, Koutinhoun BG, Missohou A, Fayomi J, Fagbohoun A et Djago Y (2009) Comparaison des performances de croissance de lapereaux en engraissement nourris par un aliment à base de tourteau de tournesol, soit sous forme farineuse soit sous forme granulée. Livest Res Rural Dev 12 (21)
- Kpomasse CC, Houndonougbo MF and Chrysostome CAAM (2015) Optimal level of cowpea (*Vigna unguiculata*) pods in feed of growing rabbit (*Oryctolagus cuniculus*): Digestibility study and growth performances. Res Opin Anim Vet Sci 5(10): 411-415.
- Lebas F, Djago AY, Kpodékon M (2007) Méthodes et Techniques d'élevage du Lapin. Elevage du lapin en milieu tropical. 2^{ème} édition révisée. Editeur : Association "Cuniculture". Corronsac France, 71 p.
- N'gom S (2004) Ebauche d'un referentiel sur la composition chimique et valeur nutritive des matieres premieres utilisables en alimentation des volailles au senegal. Thèse de doctorat de troisième cycle de chimie et biochimie des produits naturels. Université Cheick Anta Diop de Dakar Sénégal, p: 158.
- Nuhu F (2010) Effect of Moringa leaf meal on nutrient digestibility, growth, carcass and blood indices of weaner rabbits. Thesis of Kwame Nkrumah University of science and technology, Kumasi faculty of agriculture and natural resources, 107 p.
- Ogunsipe MH, Agbede JO and OA Adedeji (2014) Performance response, carcass evaluation and economic benefit of rabbits fed sorghum offal-based diets. Afr J Food Agr Nutr Dev 14: 8586-8601.
- Okachi VCW, Ani AO (2016) Effects of dietary inclusion of vitamins c and e on nutrient digestibility, haematological and carcass characteristics of rabbits in a hot humid tropical environment. Intern J Agr Res Rev 4(3): 494-503.
- Oke OS, Bamgbose AM, Alabi JO, Olagoke KO, Omotoso BO, Ige OS, Muhammad SB (2016) Performance and nutrient digestibility of acha grains (*Digitaria iburua*) as replacement for maize in the diet of female grower rabbits. J Nat Sci Res 9:58-60.
- Oladunjoye IO, Ojebiyi OO (2011) Feeding value of processed breadfruit (*Artocarpus altilis* sosberg) meal for grower rabbit. Afr J Basic Appl Sci 3(3): 98-104.
- Osakwe, Ikechukwu I and Nwuguru NR (2008) Feed intake and nutrient digestibility of weaner rabbits fed cassava peel as replacement for maize. Anim Res Intern 5(1): 770-773.
- Olatunji AK, Alagbe OJ, Hammed MA (2016) Effect of varying levels of *Moringa olifera* leaf meal on performance and blood profile of weaner rabbits. Intern J Sci Res 5:
- Ozung PO, Kennedy OO, Agiang EA, Eburu PO, Evans EI, Ewa CE (2017) Growth Performance and Apparent Nutrient Digestibility Coefficients of Weaned Rabbits Fed Diets Containing Different Forms of Cocoa Pod Husk Meal. Agr Food Sci Res 4(1): 8-19.
- Rivière R (1991) Manuel d'alimentation des ruminants domestiques en milieu tropical. Paris, France : la documentation française, 529 p.
- Shaahu DT, Dzungwe EN and Akpe ME (2014) Digestibility and Growth Performance of Rabbits Fed Raw or Processed *Lablab Purpureus* Seed in Diets. Intern J Adv Agr Sci Technol 2(6): 1-7.
- Tchibozo SVC, Dossa LH, Alowanou GG, Koura BI, Lesse P, Houndonougbo F, Houinato MB (2017) Valorisation des Ressources Alimentaires Locales en Alimentation des Lapins en Engraissement en Afrique: Synthèse bibliographique. J Appl Biosci 120: 12036-12041.
- Toleba SS, Mensah GA, Zougou CGT, Codjo B, Kpera GN et Pomalegni SCB (2007) Inventaire des ingrédients alimentaires simples et composés utilisés pour nourrir l'aulacode d'élevage au sud et au centre du Bénin. Bulletin de la Recherche Agronomique du Bénin, Numéro 57 – Septembre 2007.
- Wafar RJ, Yakubu B, and Lalabe BC (2017) Effect of Feeding Raw Kapok (*Ceiba pentandra*) Seed Meal on the Growth Performance, Nutrient Digestibility, Carcass and Organ Weights of Weaner Rabbits. Asian Res J Agr 5(3): 1-8.
- Yao Konan F, Otchoumou KA, Wognin Legbe RMF, Konan KA, Niamke S (2016) Growth parameters, protein digestibility and health status of rabbit *Oryctolagus cuniculus* fed with palatable leafy vegetables. European Sci J 12: 2016.