

Journal of Chemical, Biological and Physical Sciences



An International Peer Review E-3 Journal of Sciences

Available online at www.jsbsc.org

Section C: Physical Sciences

CODEN (USA): JCBPAT

Research Article

Activity of radionuclides in soils contaminated by scrap iron residues in the first (1st) district of Cotonou (Bénin)

Francis Théotime Mahudjro Hounsou^{1,4}, Alphonse Sako Avocefohou^{2,3}, Alassane Youssao Abdou Karim^{2*}, Christian Akomonla¹, Papin Montcho², Bertin Ahotondji Gbaguidi³, Lamine Baba-Moussa⁴, Kuassi Marcellin Amoussou-Guenou¹,

¹.Unité d'Enseignement et de Recherche de biophysique, médecine nucléaire, Faculté des Sciences de la Santé de Cotonou (FSS), Université d'Abomey-Calavi (UAC), Bénin

².Unité de Recherche en Ecotoxicologie et Etude de Qualité, Bénin/ Laboratoire d'Etude et de Recherche en Chimie Appliquée/ Université d'Abomey-Calavi (UAC), Bénin

³.Laboratoire de Recherche en Biologie Appliquée (LARBA), Ecole Polytechnique d'Abomey-Calavi (EPAC), Université d'Abomey-Calavi (UAC) Bénin

⁴.Laboratoire de Biologie et de Typage Moléculaire en Microbiologie, Faculté des Sciences et Techniques (FAST)/Université d'Abomey-Calavi (UAC), Bénin

Received: 15 October 2022; **Revised:** 24 November 2022; **Accepted:** 05 December 2022

Abstract: The scrap metal waste export activity has proliferated in Bénin and in the West African sub-region in recent decades. It is a sector that employs a growing number of workers, particularly on the former SONICOG site in the first district of Cotonou in Bénin. Unfortunately, this scrap metal waste can constitute chronic sources of radioactive contamination of the soil and of exposure of populations to ionizing radiation. The aim of this research work is to estimate the specific activities of the soil and the radionuclides involved on the site of the former SONICOG scrap metals, which has been in operation for more than two decades. The methodological approach consisted in taking soil samples using the diagonal method and analyzing them by gamma spectrometry using a Baltitic Scientific Instrument, Ltd (BSI) type

device fitted with GammaPro analysis software. The results show a stronger contamination of the superficial horizons by radionuclides such as lead (^{212}Pb), thorium (^{232}Th), cesium (^{137}Cs) in general and compared to the control soil. On the other hand, potassium (^{40}K) is detected only in the surface horizon of the soil₁ while actinium (^{228}Ac) was found in the first two surface horizons of the soil₂; as for americium (^{241}Am), it has been detected in the soil₃ but it is not quantifiable (low concentrations). Moreover, all these radioelements are in low quantity in the control sample (soil₀) chosen as the reference soil. This observation, coupled with an unequal distribution of radioactive sources in the soil horizons, confirms the hypothesis that the radioactive sources come from ferrous and non-ferrous metals from scrap metal waste on the various sites. Workers in the scrap metal sorting-recycling chain risk the effects of direct chronic exposure to low doses of ionizing radiation emitted by these sources and populations from indirect exposure.

Key words: scrap metal, contaminated soil, ionizing radiation, radionuclides, Cotonou

INTRODUCTION

Some radionuclides are produced by humans, such as cesium-137 ^[1-3], and others are present in nature, such as potassium -40, lead -210, uranium -238 and thorium -232^[4-7]. Radionuclides are present everywhere in small quantities, the greatest exposure to radiation comes from radioelements that are naturally present in the environment ^[8-10]. These emanate from the earth's crust and cosmic radiation (from outside the earth's atmosphere). Since the atomic era, the migration of radioelements from anthropogenic sources into the food chain, particularly from contaminated sites and waste, has been a subject of concern for scientists because of the harmful effects of ionizing radiation on the human body ^[11,12]. Indeed, radioelements are characterized by their nuclear instability which is manifested by the particulate radiation and/or photons emissions ^[12] of high energy which gives them this ionizing character^[13-15]. For instance, cesium-137 is found in the environment following accidental releases, aerial tests of nuclear weapons and accidents (1957) in Kyshtym (Chelyabinsk, Russia) and Windscale (United Kingdoms), Chernobyl in 1986 ^[16] and Fukushima Daiichi ^[1,17] on March 11th, 2011. Since potassium is an essential element for life, potassium-40 is the main source of exposure to radioactivity of healthy organisms by ingestion^[18]. Potassium -40 is naturally β^- and α radioactive. It is naturally present in the earth's crust. Currently, about ten radionuclides are regularly measured in the French environment, including cesium -137 as well as americium -241 ^[6]. Actinium -228 is naturally β^- and α radioactive while americium ^{241}Am is naturally α radioactive. Cesium -137 (^{137}Cs) is naturally radioactive β^- and α . As for lead-212 (^{212}Pb), it is naturally radioactive β^- and radiogenic belonging to the radioactive decay chain of thorium -232 (^{232}Th). It is generally found in the impurities of metallic ores and especially in the nuclear fuel cycle ^[19-21]. This radioactive radiation has the property of ionizing matter, unlike other electromagnetic radiation such as visible light, infrared rays or microwaves ^[22]. Human beings are exposed daily to sources of ionizing radiation ^[23] from radioactive materials of natural origin (NORM); but anthropogenic sources due to nuclear accidents are at the origin of the appearance of most pathologies, cancers, alteration of the immune system, genetic heritage and DNA over several generations ^[13,14]. Different sectors of activity can be sources of exposure to radioelements: medicine, production and use of nuclear fuel, armies, industry, mining and extraction of raw materials, food processing, etc. The doses in the human organism depend on the quantities ingested by the individual ^[24-26]. More than 60 radioelements are naturally in soil, air and water ^[23]. Among all the heavy elements that descend

from uranium or thorium, the lead-210 mostly contributes to internal exposure; it is incorporated by inhalation and especially by ingestion [27]. Uranium-238, with its two descendants, thorium-232, protactinium-240 and then uranium-232 are essentially incorporated by ingestion and are mainly concentrated in the kidneys and bones. Thorium-230 and thorium-232 enter mainly through the lungs (inhalation). Their descendant's radium-226 and radium-228 are present in the diet [28,29]. Metallurgical mining is also considered a potential source of contamination due to impurities; thus, the scrap metal waste resulting from the use of industrial products within the framework of a consumption economy on a world scale presents a double scientific and ecological interest. In Benin, and particularly in Cotonou, scrap iron residues (metals) are thrown away everywhere at random. With the risk linked to exposure to anthropogenic radiation from scrap metal waste which pollutes the soil added to sources of terrestrial and cosmic origin, it is necessary to fear the contamination of agricultural products. In fact, the soil of the Béninese coast is recommended for various crops, planting forest species [30] and gardening [31] using compost household and urban waste. In addition, the former SONICOG site, receptacle for residue from refrigerators, bacteria and any type of scrap metal is a reservoir of radioelements [32]. Investigations on the presence of radioelements in this former SONICOG scrap metal dump (Akpakpa, Cotonou) will make it possible to establish the contamination degree of the soil by metallic waste by the radioactivity.

MATERIALS AND METHODS

Choice of sampling sites: Four sites are chosen on the basis of different criteria: a control site₀ (soil₀), a site in the area of the former SONICOG where scrap metal waste is deposited for sorting, recycling and recovery operations before export site₁ (soil₁) and a secondary scrap metal disposal site located 1 km south of site₁. Site₀ is the control site whose characteristics are those of a soil without scrap iron, household nor industrial waste.

Sampling: The material consists of soil samples taken from the site of the former SONICOG scrap metal dump at Akpakpa not far from the Cotonou lagoon (**Figure 1**) and within a radius of 1 km from this site₁. Three of the four soil sampling sites are landfills located on the SONICOG site₁ (soil₁) and respectively on a radius of 1 km to the North (soil₂) and to the south (soil₃) of the SONICOG site₁. Site₂ and Site₃ are secondary burial sites located outside SONICOG. The fourth sampling site denoted by (site₀) is free of scrap scrap; we had taken the control soil (Soil₀) which will be used for the comparison of the specific activities. Soil samples are taken at three depth levels (at the surface or 0 cm level, at 10 cm and at 20 cm depth) in horizontal areas of the identified sites, not subject to runoff, undisturbed and excluding any possibility of stagnation of rainwater.




Figure 1: Location map of the study area

The three sampling sites were randomly chosen. Once collected, the samples are weighed and then crushed. The soil samples were taken using a steel shovel on a delimited surface of 1 meter square (m^2) side. The samples were taken at each corner as well as at the intersection point of the two diagonals of this surface to constitute a composite sample. And then these composite samples are homogenized and treated by appropriate methods in order to ensure their stability.

Activities' determination by gamma spectrometry method: A conventional spectrum acquisition chain comprises apart from the detector and its preamplifier, an amplifier ensuring the amplification and shaping of the signal, a digital converter and a multi-channel analyzer. The principle of gamma spectrometry is based on the interaction of photons emitted by radioactive sources with matter. The interaction of photons with matter is done either by photoelectric effect, by Compton effect or by creation of pairs.

With a reference punctual source consisting of the deposit of a radionuclide, located at a distance d from the detection crystal, a spectrum is recorded for an acquisition period t . For each energy E emitted by the source, a peak is observed in the spectrum whose surface $N(E)$ is proportional to the number of photons emitted by the source and to the efficiency of the detector for the energy considered.

The total absorption efficiency, $RP(E)$ is determined according to:

$$RP(E) = N(E) \times A \times I(E) \times t$$

$N(E)$: number of counts recorded in the total absorption peak; **A :** activity of source at time of measurement (Bq); **$I(E)$:** emission intensity of the energy line E ; **t :** measurement duration (active time); **$RP(E)$:** efficiency for energy E and the source-detector geometric conditions of the measurement.

Depending on the case, the number of counts in the total absorption peak must be corrected by different factors.

Samples' analysis: The various analyses of soils radioactive activities were carried out by gamma spectrometry at the Environmental Study and Monitoring Laboratory (ESML/LESE) in Bénin. Gamma spectrometry is a fast, non-destructive and multi-elements technique that does not require performing radiochemistry on the sample [33]. To succeed in these analyses, we have mixed the samples on the squares of one metre (1 m) sides and the composite sample obtained is dried in an oven (70°C) then crushed and sieved. The fine fraction of the treated sample is passed to the spectrometer [10,34] to determine the activity (in mBq/g) of the six radionuclides detected (^{212}Pb , ^{232}Th , ^{137}Cs , ^{40}K , ^{228}Ac , and ^{241}Am). But the radionuclides ^{234}Th , ^{64}Cu and ^{22}Na have been searched but are not detected.

The analysis of the samples was carried out using a gamma spectrometer of the Baltic Scientific Instrument (BSI) type fitted with GammaPro analysis software; this device has an analyser which converts the energy spectra of radioactive sources into specific activity and applies corrections using the calibration curves in energy and efficiency. This spectrometer has three detection chains for the analysis of radioelements and the quantification of the samples' activity. It is made up of a high-purity germanium-type semiconductor (hyper-pure Germanium HPGe) characterized by a high energy resolution of 1.8 Kev, an excellent peak symmetry and a relative efficiency of 30%, lodged in a vacuum chamber (cryostat) held by an electrical insulator support. A pre amplifier performs the charge-voltage conversion, and delivers a pulse proportional to the charge collected. An Amplifier allows the pulse received from the preamplifier to be converted into a Gaussian signal while the signal is being amplified. A multi-channel analyser made up of a multi-channel analyser card, inserted into the computer and comprising an ADC (Analog to Digital Converter) which collects the

signals and builds the calibration curves in energy (figure 2) and in efficiency (figure 3) from the table of values.




Figure 2: Energy calibration curve




Figure 3: Efficiency calibration curve of the HPGe detector

Analysis and calculation of activity, uncertainty and detection limit of the radionuclides detected in the sample: Using the Spectrum of the soil sample, the specific activity and the limit of detection (LDD or AMD) of the radionuclides are calculated with the formulas below and a counting time of 161982s.

$$A = \frac{S_{real}}{\epsilon \cdot I \cdot t} \quad \text{and} \quad LDD \text{ ou } AMD = \frac{4,66 \cdot \sqrt{Nbf}}{\epsilon \cdot I \cdot t \cdot m}$$

N.bf: Surface of the background noise; **ε:** efficiency calculated using the equation $y = 3.149x^{-0.793}$.

The results of the calculations are presented in tables.

Separation of ^{137}Cs using a soil sample with ^{134}Cs as radioactive tracer

Radiochemical treatment: A mass of 200 g of this sample was analysed by gamma spectrometry and the result shows whether there is the presence of ^{137}Cs or other artificial radioelements in the sample. The radiochemical preparation consists of extracting the ^{137}Cs in a small mass of around 20g from the soil sample with hydrochloric acid for the destruction of the sample and its dissolution. In fact, its purpose is to prevent the fixation of the atoms of ^{137}Cs and of the ^{134}Cs added as a tracer. The correction of the tracer activity was carried out using the initial activity on the date of the analysis, the density of the extract and its half-life which is 2 years. The corrected activity on the injection date is denoted A_1 .

The protocol consists of periodically adding about 50-100 ml of hydrochloric acid to a beaker, stirring the mixture on a hot plate under a temperature between 130-150°C. And then let the sample dry on the hot plate and continue attacking the sample with nitric acid HNO_3 still hot and finally allowing the sample to dry again on the hot plate.

RESULTS AND DISCUSSION

Recognition of radioelements by gamma spectrometry: Different radioelements have been sought on the four types of soil (Table 1) from the characteristic energies; the results compared with the quantification and detection thresholds made it possible to define the levels of soil contamination. Soil sample spectra were used to determine activities. The values of the energies compared to the limits of detection (LDD) and of quantification made it possible to define different levels of evaluation: - not detected; Trace: < quantification threshold; Low: > detection threshold (or

detected). The recognition of radioelements by gamma spectrometry was carried out from their energy in KeV at the surface, at 10 and 20 cm deep (table 1). The device has detected characteristic energies which signify the presence of the radioelements whose energies are presented in table 1 above. This table shows that radioactive lead (^{212}Pb), thorium (^{232}Th) and caesium (^{137}Cs) whose energies are presented were detected in the surface horizons in the four types of soil samples including samples control soils without any contamination by burying scrap metal. On the other hand, potassium ^{40}K was detected only of soil₁ sampling whereas actinium (^{228}Ac) and americium (^{241}Am) were respectively detected in soil₂ and soil₃. However, the distribution of radionuclides in soil horizons is not uniform. In the general case, the superficial horizon (0 cm) presents specific activity values higher than the 0-10 cm horizon and the latter higher than the 10-20 cm one. The three radioelements are lower in the control sample compared to the Soil₁, Soil₂ and Soil₃ where scrap metal waste is exposed and buried.

Table (1): Recognition of radioelements by gamma spectrometry from their energy in KeV at the surface (0 cm), at 10 and 20 cm

| <i>Radionuclides</i> | <i>Soil 0</i> | <i>Soil 1</i> | <i>Soil 2</i> | <i>Soil 3</i> |
|-----------------------------------|---------------|---------------|---------------|---------------|
| ^{212}Pb (Lead-212) | 238.6 | 238.6 | 238.6 | 238.6 |
| ^{232}Th (Thorium -232) | 63.3 | | 63.3 | |
| ^{137}Cs (Cesium -137) | 661.6 | 661.6 | 661.6 | 661.6 |
| ^{40}K (Potassium -40) | - | 1525.3 | - | - |
| ^{228}Ac (Actinium-228) | - | - | 911.1 | - |
| ^{241}Am (Americium-241) | - | - | - | 59.5 |
| ^{234}Th (Thorium -234) | - | - | - | - |
| ^{64}Cu (Copper-64) | - | - | - | - |
| ^{22}Na (Sodium-22) | - | - | - | - |

-: radioelements not found

The main source of these radionuclides is exogenous, but the control samples show values characteristic of the geochemical background. On this control site, the soil horizons at depth are less or not contaminated. This confirms that the source of radioelements is anthropogenic. Except for caesium, the three elements were found in all soil samples.

Distribution of average specific soil activities: The graph in figure 4 below shows the variations in the specific activities of the soils sampled for the three most represented radionuclides, namely Pb-212, Th-232 and Cs-137. The four soil sample collection points are designated by soil₀ to soil₃ in the graph. They show average specific activities of different intensity levels in the three depth levels. Soil₀ or control soil shows the lowest specific activity intensities with average values ranging from 0.15 to 1.30 mBq/g regardless of the sampling depth. It is followed by soil₃ (6.03 to 12.59 mBq /g), soil₁ (6.57 to 14.91 mBq /g), and soil₂ (1.01 to 17.54 mBq /g).

The graph in figure 5 above shows the percentages of the specific activities of the soils sampled for the three most represented radionuclides, namely Pb-212, Th-232 and Cs-137. Considering the three main radionuclides, soil₁ confirms the highest radioactive contamination with 35% of the total radiation of the three sources. It is followed by soil₂ (32%) and soil₃ (31%) of all sources' activity. In addition to the three main radionuclides detected in the control soil, new radionuclides are detected at the other sampling points. These are potassium 40 (^{40}K), actinium 228 (^{228}Ac) and americium 241 (^{241}Am).




Figure (4): Average specific soil activities in mBq/g




Figure (5): Relative distribution (in %) of total specific soil activities of the three main radionuclides (Pb-212, Th-232 and Cs-137) in soil horizons




Figure (6): Relative distribution (in %) of the total specific activities of the radionuclides studied (Pb-212, Th-232, Cs-137, K-40, Ac-228 and Am-241) in soil horizons

Considering all the radionuclides detected, soil₁ confirms the highest radioactive contamination with 51% of the total radiation from the sources, as does soil₂ with 29%. Soil₃ is the latest with 19% of total sources' activity. The control soil (Soil₀) has decreased from 2 to 1%. In fact, the soil₁ revealed on the surface a source of potassium 40 with a high specific activity of 137.501 mBq/g. This potassium would certainly come from recent additions of sand or materials rich in this natural isotope, the most abundant of all the sources of natural human exposure. This is the most important specific activity recorded in this study. As for soil₂, it is characterized by the appearance of actinium on the first two depth levels at concentrations of 34.64 and 12.01 mBq/g. This decrease indicates an anthropogenic source unrelated to the bedrock (the mother rock).

Evolution of control soil activities (soil₀) in function of depth: The device has detected characteristic energies which indicate the presence of Pb-212, Th-232 and Cs-137 on the first two levels, namely level 0-0 on the surface, and level 0-1 at 0 to 10 cm deep. In general, the intensities are decreasing from surface to depth. It should be noted that there is a total absence of radioactive

activity in the horizons of 20 cm and more. We consider by hypothesis that the specific activity values obtained are characteristic of the geochemical soil background of the Akpakpa quarter where all the sampling sites are located.

Table 7a: Activities of radionuclides in the control soil sample (in mBq/g) taken from the SONICOG compound

| <i>Soil</i> | <i>Level</i> | ^{212}Pb | ^{232}Th | ^{137}Cs | ^{40}K | ^{228}Ac | ^{241}Am |
|-------------------------|--------------------|--------------------|--------------------|-------------------|-----------------|-------------------|-------------------|
| <i>Soil₀</i> | 0-0 | 3.15 | 0.357 | 1,802 | - | - | - |
| | 0-1 | 0.75 | 0.09 | 0.802 | - | - | - |
| | 0-2 | Trace | Trace | Trace | - | - | - |
| <i>Average</i> | <i>1.30</i> | <i>0.15</i> | <i>0.87</i> | - | - | - | |

Legend: - not detected; Trace: < quantization threshold.

Sampling levels: on the surface (Level 0-0), 0 to 10 cm (Level 0-1) and 10 to 20 cm (Level 0-2)

The device has detected characteristic energies which indicate the presence of Pb-212, Th-232 and Cs-137 on the first two levels, namely level 0-0 on the surface, and level 0-1 at 0 to 10 cm deep. In general, the intensities are decreasing from surface to depth. It should be noted that there is a total absence of radioactive activity in the horizons of 20 cm and more. We consider by hypothesis that the specific activity values obtained are characteristic of the geochemical soil background of the Akpakpa quarter where all the sampling sites are located.

Evolution of SONICOG burial soil activities in function of depth: The device has detected characteristic energies that confirm the presence of Pb-212, Th-232, Cs-137 and K-40. The samples from soil₁ were taken in the main landfill area of the former SONICOG site in the part where the scrap metal waste was buried for about twenty years. This could explain the fact that samples from this site showed a high intensity of specific activity of radionuclides with a relatively higher specific activity with 34.488 mBq/g for the ^{212}Pb , 47.728 mBq/g for ^{232}Th and 14.802 mBq/g for the ^{137}Cs . These values are significantly higher than those of the control site located in the same enclosure.

Table 7b: Activities of radionuclides in soil₁ (mBq/g) taken from the SONICOG compound

| <i>Soil</i> | <i>Level</i> | ^{212}Pb | ^{232}Th | ^{137}Cs | ^{40}K | ^{228}Ac | ^{241}Am |
|-------------------------|---------------------|---------------------|--------------------|---------------------|-----------------|-------------------|-------------------|
| | 1 -0 | 34,488 | 47,728 | 14,802 | 137,501 | - | - |
| <i>Soil₁</i> | 1 -1 | 12,075 | 11.75 | 10,602 | Trace | - | - |
| | 1 -2 | 0.126 | Trace | -222 | Trace | - | - |
| <i>Average</i> | <i>12.00</i> | <i>14.91</i> | <i>6.57</i> | <i>45.83</i> | - | - | |

Legend: - not detected; Trace: < quantization threshold.

Sampling levels: on the surface (Level 1 -0), 0 to 10 cm (Level 1 -1) and 10 to 20 cm (Level 1 -2)

Evolution of secondary burial soil activities (soil₂) in function of depth: In addition to soil radionuclides, the device has detected characteristic energies that indicate the presence of actinium 228, a transuranic element.

Pb -212, Th-232 and Ac-228 have respectively a specific activity of 37.37 mBq/g, 24.35 mBq/g and 34.64 mBq/g in the surface samples on this site located 1km to the north of SONICOG (far from the sea side). While Cs-137 exhibited a relatively lower specific activity of 2.31 mBq/g at the surface. Pb -212, Th-232, Cs-137 and Ac-228 have respectively a specific activity of 15.24 mBq/g, 11.29

mBq/g, 0.72 mBq/g and 12.01 mBq/g in samples taken at 10 cm deep. We also detected the presence of these nuclides at a depth of 20 cm in traces except for Cs-137 which was not detected at this depth

Table 7c: Activities of radionuclides in soil samples 2 (mBq/g) taken 1 km from SONICOG to the north

| Soil | Level | ²¹² Pb | ²³² Th | ¹³⁷ Cs | ⁴⁰ K | ²²⁸ Ac | ²⁴¹ Am |
|-------------------|--------------|-------------------|-------------------|-------------------|-----------------|-------------------|-------------------|
| Soil ₂ | 2 -0 | 37.37 | 24.35 | 2.31 | - | 34.64 | - |
| | 2 -1 | 15.24 | 11.29 | 0.72 | - | 12.01 | - |
| | 2 -2 | Trace | Trace | - | - | Trace | - |
| Average | 17.54 | 11.88 | 1.01 | - | 15.55 | - | - |

Legend: - not detected; Trace: < quantization threshold.

Sampling levels: on the surface (Level 2 -0), 0 to 10 cm (Level 2 -1) and 10 to 20 cm (Level 2 -2)

Evolution of secondary burial soil activities (soil₂) in function of depth: The device detected characteristic energies which make it possible to confirm the presence not only of Pb-212, of Th-232 and of Cs-137, radionuclides detected in the sol0 reference soil.

In addition to soil radionuclides, the device has detected characteristic energies which indicate the presence of americium 241, a transuranic element. Pb -212, Th-232 and Cs-137 have respectively a specific activity of 24.514 mBq/g, 21.185 mBq/g and 12,830 mBq/g in surface samples at this site located 1km south of SONICOG (sea side). While Am-241 exhibited relatively lower and unquantifiable specific activity at all levels. We also detected the presence of Pb-212 (13.25 mBq/g) , Th-232 (10.12 mBq/g) and Cs-137 (5.25 mBq/g) of weak activity in the 10 cm horizons and traces deeper at 20 cm.

Table 7d : Activities of radionuclides in soil₃ (mBq/g) taken 1 km from SONICOG to the south (near the Atlantic Ocean)

| Soil | Level | ²¹² Pb | ²³² Th | ¹³⁷ Cs | ⁴⁰ K | ²²⁸ Ac | ²⁴¹ Am |
|-------------------|--------------|-------------------|-------------------|-------------------|-----------------|-------------------|-------------------|
| Soil ₃ | 3 -0 | 24,514 | 21,185 | 12.83 | - | - | Low |
| | 3 -1 | 13.25 | 10.12 | 5.25 | - | - | Low |
| | 3 -2 | Trace | Trace | Trace | - | - | Trace |
| Average | 12.59 | 10.44 | 6.03 | - | - | - | |

Sampling levels: at the surface (Level 3 -0), 0 to 10 cm (Level 3 -1) and 10 to 20 cm (Level 3-2)

In this study, on the four types of soil studied, certain radioelements are constantly found, namely lead (²¹²Pb), thorium (²³²Th) and cesium (¹³⁷Cs). However, other types of radioactive elements such as potassium (⁴⁰K), actinium (²²⁸Ac), and americium (²⁴¹Am) appeared only in the scrap metal waste burial soils. From the analysis of these results, we can propose two interpretations, namely, natural and artificial sources. Indeed, lead 212 is radiogenic and belongs to the family of radioactive decay of Th-232. The presence of these two radionuclides in all the samples can be explained by their common origin, which is found in ferrous and non-ferrous metal ores.

They could also be originated from the natural geochemical backgrounds of the soils studied insofar as the control soil has concentration levels in the superficial horizons different from zero. On the other hand, these metals disappear at the 10-20 cm horizon on practically all the soil samples except

soil₁ where the soil has a low specific activity (0.126 mBq /g). This contradicts the first hypothesis based on the geochemical background. This is confirmed by the appearance of potassium 40 only in the main burial area on the former SONICOG site (soil₁) in a very high quantity (137.501 mBq /g). However, this value is lower than that obtained by Chakraborty *et al.*^[9], who has reported a specific activity of 451.90 ± 24.89 mBq/g for this radionuclide in Chittagong city in Bangladesh. This high concentration could be explained by the fact that on this site, scrap metal waste was buried there for about twenty years.

This is in agreement with a higher specific activity of the radionuclides present with 34.488 mBq/g for ²¹²Pb, 47.728 mBq/g for ²³²Th and 14.802 mBq/g for ¹³⁷Cs. It should be noted that the figures of 38.47 ± 2.72 and 2.41 ± 0.18 Bq kg⁻¹ obtained by Chakraborty *et al.*^[9], respectively for Th-232 and Cs-137 are lower than the values recorded for the soil₁ which is the most contaminated by radioactive elements. On the other hand, Kall *et al.*^[35] have reported a much higher level of specific activities in soils for Th-212 (81 ± 20 to 161 ± 23 Bq.kg⁻¹) and K-40 (112 ± 34 to 565 ± 50 Bq.kg⁻¹) on the shore of French Bay in the localities of Ankorikihely , betahitry and Lazaret-Nord. Phosphogypsum (PG) waste associated with phosphoric acid produced in industrial fertilizer and chemical production facilities in Egypt can contaminate soils^[36].

Indeed, El Afifi *et al.*^[36] have found concentrations of 8.3 ± 0.76 and 64.3 ± 4.1 Bq/Kg of Th-232 and K-40 respectively in chemical fertilizer manufacturing residues in Egypt. The concentrations obtained are much higher than our data for Th-232 and lower for K-40. The IAEA exemption limits are 10 Bq/kg for ⁴⁰K and ¹³⁷Cs^[37]. These limits are exceeded in most samples at the surface and at the second depth level^[8,32,38].

There is actinium (²²⁸Ac) and potassium (⁴⁰K) in the soil₁ with a high specific activity of 137.501 mBq/g for potassium and 34.64 mBq/g for ²²⁸Ac while the latter belongs to the ²³²Th family; these results are in accordance with those of Kelly *et al.*^[39]. As for the secondary soil samples located 1 km to the south (towards the sea) and 1 km to the north (on the mainland), we have mainly noted the presence of ²¹²Pb, ²³²Th and ¹³⁷Cs and that of actinium and americium in some soils. Thorium has a specific activity of 21.185 Bq/g on the site 1 km towards the sea (soil₃) on the surface on the secondary site located to the south. On the site 1 km on the mainland to the north (soil₂), the presence of actinium was noted with a specific activity of 34.64 mBq/g.

On the control site on which there was no burial of scrap metal waste, only a few rare radioelements such as Th, Pb and Cs are obtained, but americium (²⁴¹Am) and actinium (²²⁸Ac) have presented a negligible specific activity. Whereas, on the burial site, the presence of other types of radioelements was noted, the concentration of which decreases with depth. It appears that these soils are heavily contaminated by potentially toxic metals such as zinc, lead, chromium, americium and actinium. This contamination would come from sediments or sludge, phosphate residues, battery cells, vehicle batteries, iron and other non-degradable metals such as residues of copper rods, thorium from welders buried in the ground by scrap metal sorters. We also note an increase in the specific activity of the radionuclides which starts from double for lead and triple for thorium and is explained by an accumulation effect.

If plants and/or vegetables were grown in this region, very high concentrations of radionuclides could be found in the leaves of these plants. However, high concentrations of lead have been found in leafy vegetables collected from areas of high concentration of household waste, in fish samples and in aquatic ecosystems in Bénin^[30,40,41] with high risks that these lead residues contain radionuclides. We have noticed that the different mass activities progressively decrease with depth to tend to disappear around 20 cm for certain radionuclides such as ⁴⁰K and ²²⁸Ac and according to the sampling site. It is deduced that the evolution of the concentration in the soils of these

radionuclides is dependent on a large number of parameters, namely the physicochemical and biological nature (texture) of the soils, meteorological conditions (in particular precipitation) and runoff water, and accumulation [42]. It is therefore customary to consider as a first approximation and in particular for the burials of recent radioactive scrap metal from the last 5 years an exponential decrease in the concentration of artificial radioelements according to the depth in the ground [11,43]. There is surely a phenomenon which induces a decrease in the concentration of the radionuclide [44].

This phenomenon evolves over time depending on the radioelement transfer conditions in the soil of sites containing high radioelement proportions such as the site of the former SONICOG company, the site located 1 km to the south and the site located 1 km to the north are the sites polluted by radioactivity. This confirms that highly radioactive substances have been handled and stored in an uncontrolled manner, leading to the dispersion of these substances and presenting a potential risk depending on the use made of the sites: site for sorting and recycling waste from scrap.

In general, almost all soil and water rocks naturally contain small quantities of radioactive materials such as uranium, thorium, potassium and their descendants [45-47]. The work carried out by Cresswell *et al.*, [48] also shows that urban areas have very complex radiative environments due to the diversity of radioactive sources. These authors further demonstrated that the presence of Cs-137 identifies relatively undisturbed soil and a feature called "Naturally Occurring Radioactive Materials" (NORM). They can thus be found in building materials.

These materials containing radionuclides naturally present in the earth's crust called NORM [49] had been mentioned by Desideri *et al.* [6]; these determine the geochemical distribution of natural (U, Th) and anthropogenic (Pu, Am, ¹³⁷Cs) radionuclides in three sediment samples of different origin. TE - NORM stands for Technically enhanced natural radioactive materials for large volumes of waste with low specific activity emitted by several industrial activities such as mineral extraction, beneficiation and chemical treatment of ores, the production of phosphate fertilizers, extraction and purification of trace elements, etc.

The Pb-212 could have its origin in the rocks containing residues of Th-232 from which would come the buried metals. Indeed, Worden *et al.* [45] have demonstrated that sandstones exceptionally enriched in uranium with concentrations of 0 to 70 ppm would be at the origin of the contamination of waters with Pb-210 from which it would come inside the reservoir (rock). These concentrations, generally low, are observed on the control site, that is to say the site without burial of waste from scrap metal and vary between 3.21 mBq/g for thorium 212 and 0 mBq/g for potassium 40. The (TE-NORM) stands [17] for "Enhanced Technology Naturally Occurring Radioactive Materials" is due to human activities operating in various industrial sectors [15].

Human activities can during the handling, storage, burial and storage of products and waste as well as production equipment concentrate materials on a site. Thus, the radioactivity is concentrated in the products and/or residues, including food products [7,9]. And consequently, the level of radioactivity becomes higher than that encountered in the natural state. This is verified on the main scrap metal waste landfill site and on all the secondary sites with respectively thorium which goes to 47.728 Bq/g, lead to 34.488 mBq/g and potassium to 137.501 mBq/g. It is important to note that the two secondary sites alone concentrate 63% of the total activity of the three main radionuclides found in all the sites, namely Pb-212, Th-232 and Cs-137. The natural radioactive elements which are the basis of the TE-NORM are present everywhere in the earth's crust, but the only difference in their distribution will be their affinity with respect to such or such other element and this will define their abundance from one environment to another, from one ore to another, hence their chemical speciation [6].

However, the physical observation of the waste on the various scrap processing sites reveals, before their burial, the presence, among other things, of aluminium residues, burnt copper, batteries and batteries of vehicles, electronic waste, etc. These areas typically contain fine sand particles, corrosion particles, paint chips, battery packs, and other fine particles breaking off from scrap metal deposits that can be the source of Pb-212 found in the TE-NORMs. These results agree with those of Egidi^[38] and El Afifi *et al.*^[50] on radionuclides emitted from NORM and TE-NORM. The agricultural amendment products can significantly influence the chemical properties of sandy soils⁵¹, because they are poor in minerals.

CONCLUSION

This study has been carried out to assess the radiological quality of scrap metal waste on the former SONICOG site in Cotonou. It reveals the presence of radioelements whose activity is generally low in samples taken at a depth of 20 cm. This weak natural radioactivity would be reinforced by the waste which was on the sites and which produces a specific activity far superior to the exemption limits recommended by the IAEA. This observation confirms that the soils are contaminated by scrap metal waste buried underground. Under these conditions, environmental monitoring becomes vital and must above all be a major concern of environmental specialists for the prevention of the populations' health and especially of workers carrying out activities on the SONICOG site. The basic standards relating to the health protection of the population and workers against the dangers resulting from ionizing radiation should be reinforced in Benin and in the sub-region.

REFERENCES

1. IRSN, Exposition de la population française aux rayonnements ionisants-bilan 2014-2019. Rapport 2019
2. H. Nifenecker, Eléments de réflexion sur les pathologies éventuellement induites par le Césium 137. CNRS LPSC/IN2P3. 53 avenue des Martyrs Grenoble, 2006, 38026.
3. H.M. Saleh, T.A. Bayoumi, H.H. Mahmoud, R.F. Aglan, Uptake of cesium and cobalt radionuclides from simulated radioactive wastewater by *Ludwigia stolonifera* aquatic plant. *Nuclear Engineering and Design*, 2017, 315, 194–199
4. J. P. Knochel, L. N. Dotin, R. J. Hamburger. Pathophysiology of intense physical conditioning in a hot climate. I. Mechanisms of potassium depletion. *The Journal of clinical investigation*, 1972, 51(2), 242-255.
5. M.Y. Shoeib and K.M. Thabayneh, Assessment of natural radiation exposure and radon exhalation rate in various samples of Egyptian building materials. *Journal of Radiation Research and Applied Sciences* 2014, journal homepage: <http://www.elsevier.com/locate/jrras>
6. D. Desideri, M.A. Meli, L. Feduzi, C. Roseli, The importance of radiochemistry for the characterization of NORM and of environments contaminated by NORM, *International Journal of Environmental Analytical Chemistry*, 2006, 86, 601-613.
7. K.M. Thabayneh, N.M. Jassar, Natural radioactivity levels and estimation of radiation exposure in environmental soil samples from Tulkarem Province-Palestine. *J Soil Sci*, 2012, 2:7–16.

8. Planinsek Petra, Smodis Borut, Benedik Ljudmila, Uptake of natural radionuclides from contaminated soil into vegetables and consequent dose assessment. *Journal of Radioanalytical and Nuclear Chemistry*, 2018, 318:2373–2379
9. S.R.Chakraborty, R. Azim , A.K.M. Rezaur Rahman, R.Sarker, Radioactivity concentrations in soil and transfer factors of radionuclides from soil to grass and plants in the Chittagong city of Bangladesh. *J Phys Sci*,2013, 24(1):95-111 http://tel.archives-ouvertes.fr/docs/00/82/10/54/PDF/jeambrun_marion_2012_ED413.pdf. [https://doi.org/10.1007/s10967-018-6221-6\(0123456789\(\).,-volV\)\(0123456789\(\).,-volV\)](https://doi.org/10.1007/s10967-018-6221-6(0123456789().,-volV)(0123456789().,-volV))
10. D. Saba, Les radionucléides et le plomb émis par une usine de fertilisants au Liban : Etude de leurs mobilités dans les sols et les plantes pour une évaluation du risque sanitaire. *Environnement et Société*. Université Paris-Saclay, 2019. Thèse de Doctorat Français. ffnNT : 2019a,SACLA025f
11. D. Saba, N. Manouchehri, S. Besançon, O. El Samad, R. Baydoun, R.B. Khozam, P. Cambier, Bioaccessibility and radioisotopes of lead in soils around a fertilizer industry in Lebanon. *Environmental geochemistry and health*,2019b, 41(6), 2749-2762.
12. B.R.S. Minty, Fundamentals of airborne gamma-ray spectrometry, *Journal of Australian Geology and Geophysics*,1997, 17, 2, 39-50.
13. WHO; Effet de la guerre nucléaire sur la santé et les services de santé. 2e Édition. Rapport du groupe de gestion de l'WHO sur la réalisation de la résolution WHA,1987,36.28.
14. T. Ohkita et J. Rohbita, Effets biologiques d'une guerre nucléaire, effets aigus des rayonnements : valeur de la DL50. Annexes. Rapport WHO,1987.
15. N.Mitchell, D. Perez-Sanchez, M.C.Thorne, A review of the behaviour of U-238 series radionuclides in soils and plants, *Journal of Radiological Protection*, 2013, 33, 2, R17-R48.
16. P. Lestaevel, R. Racine, H. Bensoussan, C. Rouas, Y. Gueguen, I. Dublineau, J.M.Bertho, P. Gourmelon , J.R. Jourdain, M. Souidi, Caesium 137: Properties and biological effects resulting of an internal contamination. *Médecine Nucléaire*, 2010, 34,108–118. doi: 10.1016/j.mednuc.2009.12.003
17. IAEA, Naturally Occurring Radioactive Material (NORM VIII) Proceedings of an International Symposium Rio de Janeiro, Brazil,2016,
18. R. Trevisi, S. Risica, M. D'Alessandro, D. Paradiso, C. Nuccetelli, Natural radioactivity in building materials in the European Union: a database and an estimate of radiological significance. *J Environ Radioact*,2012, 105:11–20.
19. Ahamdach Noureddine, Stammose Denise, Isolation and microscopic characterization of nuclear fuel particles from contaminated soil of Chernobyl. *C. R. Acad. Sci. Paris, Sciences de la Terre et des planètes / Earth and Planetary Sciences*,2000, 330, 415–422 © 2000 Académie des sciences / Éditions scientifiques et médicales Elsevier SAS. Tous droits réservés S1251805000001476/FLA
20. B. Chareyron, Situation radiologique dans l'environnement des anciens sites miniers du Lodevois (Hérault),CRIIRAD,septembre,2004 (<http://www.criirad.org/actualites/uraniumfrance/lodeve/notecriiradlodevesept04.doc>)

21. B. Chareyron, Monitoring of the radiological contamination of streams and rivers using aquatic bioindicators. I.S. RIVERS, 2018.
22. P. Guénel, L. De Guire, D. Gauvin, M. Rhains, Rayonnements non ionisants. In : Environnement et santé publique - Fondements et pratiques, 2003, pp. 441-462
23. M. Jeambrun, L'uranium et ses descendants dans la chaîne alimentaire, Thèse de Ph. D., Strasbourg, France, Laboratoire d'Hydrologie et de Géochimie de Strasbourg. En ligne, 2012.
24. Y. Barthe, Les déchets radioactifs à vie longue sont-ils gouvernables ? In Annales des Moines, 1998,
25. M. Puncher & J.D. Harrison, Assessing the reliability of dose coefficients for inhaled and ingested radionuclides. Journal of Radiological Protection, 2012, 32(3), 223.
26. N.N. Jibiri & T.H. Abiodun, Effects of Food Diet Preparation Techniques on Radionuclide Intake and Its Implications for Individual Ingestion Effective Dose in Abeokuta, Southwestern Nigeria. World Journal of Nuclear Science and Technology, 2012, 2(3), 106-113.
27. M. Souidi, E. Tissandie, R. Racine, H.B. Soussan, C., Rouas, E. Grignard & Y. Gueguen, Uranium: propriétés et effets biologiques après contamination interne. In Annales de Biologie Clinique, 2009, 67, 1, p. 23).
28. H.W. Kirby, M.L. Salutsky, The Radiochemistry of Radium, National Academy of Sciences National Research Council, 1964, 205 p. En ligne. <http://library.lanl.gov/cgi-bin/getfile?rc000041.pdf>.
29. J.L. Basdevant, Maîtriser le nucléaire: Sortir du nucléaire après Fukushima. Editions Eyrolles, 2012.
30. P. Drechsel & S. Schmall, Mineral deficiencies and fertilization of coastal reforestation in Bénin, West Africa. Fertilizer research, 1990, 23(3), 125-133.
31. K.P. Nagavallema, S.P. Wani, S. Lacroix, V.V. Padmaja, C. Vineela, M.B. Rao & K.L. Sahrawat, Vermicomposting: Recycling wastes into valuable organic fertilizer. Global Theme on Agroecosystems Report no. 8. 2004.
32. F.T.M. Hounsou, H. Sina, A.S. Avocefohou, P. Aina, A.B. Gbaguidi, K.M., Amoussou-Guenou, L. Baba-Moussa, Radiological Quality and Dangerousness of Ferrous and Non-ferrous Metals Waste in Cotonou (Bénin). Asian Journal of Environment & Ecology, 2020, 13-21.
33. L. Swiderski, M. Moszynski, A. Syntfeld-Kazuch, M. Szawlowski, and T. Szczesniak, Measuring the scintillation decay time for different energy depositions in NaI(Tl), CsI(Tl) and CsBr3 scintillators. Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment, 2014, 749, 68 – 73.
34. M. Annunziata, Handbook of Radioactivity Analysis, 2nd edition, Elsevier, 2003, 1273 p.
35. B. Kall, T. Tombo, M. Rasolonirina, N. Rabesiranana, G. Rambolamanana, Contribution à l'étude de dose due à la radioactivité gamma du sol sur la rive de la baie des Français,

- Antsiranana, Madagascar. *Afrique Science*, 2015, 11(1), 122-135, 122, <http://www.Afriquescience.info>
36. E.M.El Afifi, M.A.Hilal, M.F.Attallah, S.A.EL-Reefy, Characterization of phosphogypsum wastes associated with phosphoric acid and fertilizers production. *Journal of Environmental Radioactivity*, 2009, 100, 407–412. doi: 10.1016/j.jenvrad.2009.01.005
37. IAEA, Notification et autorisation pour l'utilisation des sources de rayonnements. (Supplément à la publication no GS-G-1.5 de la collection Normes de sûreté de l'AIEA). IAEA-TECDOC-1525, 2011.
38. P. Egidi, Introduction to Naturally Occurring Radioactive Material. In Proc. of 42nd Annual Meeting of the Health Physics Society, 1997.
39. E.A.Kelly, M.N. Lam, C. Eiroa-Lledo, S.A. Kozimor, L.M. Lilley, V. Mocko, B.W. Stein, Preparation of an Actinium-228 Generator. <https://dx.doi.org/10.1021/acs.inorgchem.9b03563>
40. E.A.Pazou Yehouenou, A. Soton, D. Azocli, H. Acakpo, M. Boco, L. Fourn, D. Houinsa, J.C.Keke, B. Fayomi, Contamination du sol, de l'eau et des produits maraîchers par des substances toxiques et des métaux lourds sur le site de Houéyiho (Cotonou) en République du Bénin. *Int. J. Biol. Chem. Sci.*, 2010, 4(6): 2160-2168.
41. A.K.A. Youssao, Etude de la distribution des résidus de plomb dans les écosystèmes aquatiques du Chenal de Cotonou et du lac Nokoué au Bénin. Thèse. Soutenue le 1er, 2011.
42. C. Ferry, La migration du Radon 222 dans un sol application aux stockages de résidus issus du traitement des minerais d'uranium. Thèse de Doctorat de l'Université de Paris-Sud U.F.R. Scientifique D'Orsay, 2000, 6127, 166 p.
43. W.L.Lennemann, The management of high-level radioactive wastes. *IAEA Bull*, 1978, 21 (4), 1-16.
44. A.E. Lebatard, Bourlès Didier, Quantification des processus superficiels et datation par les radionucléides cosmogéniques ^{10}Be , ^{26}Al et ^{36}Cl . *Quaternaire*, 2015, 26(3): 193-213.
45. R.H. Worden, D.A.C. Manning, P.R. Lythgoe, The origin and production geochemistry of radioactive lead (^{210}Pb) in NORM-contaminated formation waters », *Journal of Geochemical Exploration*, 2000, 69-70, 695-699.
46. D. Larivière, N. Guérin, Natural Radioactivity », in Atwood, D.A. (éd.), *Radionuclides in the Environment*, Chichester, John Wiley & Sons, 2010, p. 261-272.
47. H.Vandenhove, C. Hurtgen, T.E.Payne, Uranium, in Atwood, D.A. (éd.), *Radionuclides in the Environment*, Chichester, John Wiley & Sons, 2010, 261-272.
48. A.J. Cresswell, D.C.W. Sanderson, M.Harrold, B. Kirley, C.Mitchell, A. Weirb, Demonstration of lightweight gamma spectrometry systems in urban environments. *Journal of Environmental Radioactivity*, 2013, 124, 22e28. <http://dx.doi.org/10.1016/j.jenvrad.2013.03.006>
49. IAEA, Naturally Occurring Radioactive Material (NORM VII). Proceedings of an International Symposium Beijing, China, 2013, 22–26.

50. E.M. El Afifi, M.A. Hilal, S.M. Khalifa, H.F. Aly, Evaluation of U, Th, K and emanated radon in some NORM and TENORM samples. *Radiation Measurements*, 2006, 41, 627 – 633. doi: 10.1016/j.radmeas.2005.09.01
51. J. Mouton, Développement d'un procédé de lavage de sol pour le traitement simultané des HAP et du plomb (Doctoral dissertation, Université du Québec, Institut national de la recherche scientifique) 2008.

*** Corresponding author: Alassane Youssao Abdou Karim;**

Unité de Recherche en Ecotoxicologie et Etude de Qualité, Bénin/ Laboratoire d'Etude et de Recherche en Chimie Appliquée/ Université d'Abomey-Calavi (UAC), Bénin

Online publication Date: 06.12.2022