

## Article 12

Bodjrènou FSU, **Amoussa Hounkpatin W**, Termote C, Dato G, Savy M. (2021) : Determining factors associated with breastfeeding and complementary feeding practices in rural Southern Benin. *Food Science and Nutrition* 9:135–144. DOI: 10.1002/fsn3.1971

Scopus Preview Author search Sources ? [Create account](#) [Sign in](#)

## Sources

ISSN  Enter ISSN or ISSNs [Find sources](#)

ISSN: 2048-7177 x

**i Improved Citescore**

We have updated the CiteScore methodology to ensure a more robust, stable and comprehensive metric which provides an indication of research impact, earlier. The updated methodology will be applied to the calculation of CiteScore, as well as retroactively for all previous CiteScore years (i.e. 2018, 2017, 2016...). The previous CiteScore values have been removed and are no longer available.

[View CiteScore methodology >](#)

**Filter refine list**

[Apply](#) [Clear filters](#)

**Display options**

Display only Open Access Journals

Counts for 4-year timeframe

No minimum selected

Minimum citations

Minimum documents

**1 result** [Download Scopus Source List](#) [Learn more about Scopus Source List](#)

All

View metrics for year: 2020

	Source title ↓	CiteScore ↓	Highest percentile ↓	Citations 2017-20 ↓	Documents 2017-20 ↓	% Cited ↓
1	Food Science and Nutrition <i>Open Access</i>	2.7	55% 140/310 Food Science	4082	1485	61

[Top of page](#)

https://mjlclarivate.com/journal-profile

Web of Science Group **Master Journal List** Search Journals Match Manuscript Downloads Help Center Welcome, BODJRENOU Sam [Settings](#) [Log Out](#)

The power of the Web of Science™ on your mobile device, wherever inspiration strikes. [Dismiss](#) [Learn More](#)

About

General Information

Web of Science Coverage

Journal Citation Report

Open Access Information

Peer Review Information

PubMed® Information

[Return to Search Results](#)

### FOOD SCIENCE & NUTRITION [Share This Journal](#)

ISSN / eISSN 2048-7177  
Publisher WILEY, 111 RIVER ST, HOBOKEN, USA, NJ, 07030-5774

**About**

Food Science & Nutrition is the peer-reviewed journal for rapid dissemination of research in all areas of food science and nutrition. The Journal will consider submissions of quality papers describing the results of fundamental and applied research related to all aspects of human food and nutrition, as well as interdisciplinary research that spans these two fields.

**General Information**

<b>Journal Website</b>	<a href="#">Visit Site</a>	<b>Publisher Website</b>	<a href="#">Visit Site</a>
<b>1st Year Published</b>	2012	<b>Frequency</b>	Bi-monthly
<b>Issues Per Year</b>	6	<b>Country / Region</b>	UNITED STATES OF AMERICA
<b>Primary Language</b>	English	<b>Aims and Scope</b>	<a href="#">Visit Site</a>
<b>Editorial Board</b>	<a href="#">Visit Site</a>	<b>Instructions for Authors</b>	<a href="#">Visit Site</a>
<b>Avg. Number of Weeks from Submission to Publication</b>	24	<b>Plagiarism Screening</b>	<a href="#">Visit Site</a>
<b>Article DOIs</b>	Yes	<b>Host Platform</b>	Wiley Online Library
<b>Full-Text Formats</b>	PDF, HTML		



## ORIGINAL RESEARCH

# Determining factors associated with breastfeeding and complementary feeding practices in rural Southern Benin

Fifali Sam Ulrich Bodjrènou<sup>1,2</sup> | Waliou Amoussa Hounkpatin<sup>2</sup> | Céline Termote<sup>3</sup> | Geoffroy Dato<sup>1</sup> | Mathilde Savy<sup>4</sup>

<sup>1</sup>Alliance of Bioversity International and CIAT, Cotonou, Benin

<sup>2</sup>University of Abomey-Calavi, Faculty of Agricultural Sciences, Abomey-Calavi, Benin

<sup>3</sup>Alliance of Bioversity International and CIAT, Nairobi, Kenya

<sup>4</sup>NUTRIPASS-IRD, Cotonou, Benin

## Correspondence

Fifali Sam Ulrich Bodjrènou, Alliance of Bioversity International and CIAT, Cotonou, Benin; University of Abomey-Calavi, Faculty of Agricultural Sciences, Abomey-Calavi, Bénin.

Email: bodjrenousam@gmail.com

## Funding information

the Ministry of Foreign Affairs of Finland; CGIAR.

## Abstract

This study aimed at characterizing breastfeeding and complementary feeding practices in a food-insecure area of Benin and identifying factors associated with these practices. A cross-sectional study was conducted in the districts of *Bopa* and *Houéyogbé* among  $n = 360$  mother–child pairs. Children aged 0–17 months were considered. Socioeconomic characteristics among children and mothers, *Breastfeeding on demand*, *Breastfeeding frequency during children illness*, and *Positioning and Attachment of children while breastfeeding* were assessed using semi-structured interviews and observations. Qualitative 24-hr recalls were administered to mothers to compute WHO recommended complementary feeding practices indicators namely *minimum dietary diversity* (MDD), *minimum meal frequency* (MMF), and *minimum acceptable diet* (MAD) among 6–17 months old children ( $n = 232$ ). Associations between each feeding practice and mothers' socioeconomic characteristics were tested using multivariate generalized linear models. *Breastfeeding on demand* and *good positioning and attachment for breastfeeding* rates were 59% and 66%, respectively. Only 26% of mothers used to increase breastfeeding frequency when their children were ill. The proportions of children who met MDD, MMF, and MAD were 51%, 75%, and 44%, respectively. Children living in *Houéyogbé* were less likely to be breastfed on demand compared with those living in *Bopa*; however, they had better *breastfeeding frequency during illness* and meal frequency. Socioeconomic factors with significant association with breastfeeding practices were children age and sex and mothers' education, ethnicity, and employment status. Complementary feeding practices were positively associated with children's age but not with other socioeconomic characteristics. Breastfeeding and complementary feeding practices were almost suboptimal or medium and still need to be improved through well designed nutrition intervention program including nutrition education.

## KEYWORDS

Benin, Breastfeeding practices, Complementary feeding practices, Determinants

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2020 The Authors. *Food Science & Nutrition* published by Wiley Periodicals LLC

## 1 | INTRODUCTION

Benin is a West African country where child malnutrition is a public health problem. According to the latest Demographic and Health Survey (DHS), about one-third of children under 5 years old were stunted in 2017-2018 (INSAE & ICF, 2019). A major contributor to this situation is the inadequacy of breastfeeding and complementary feeding practices. About one-half of newborns (54%) benefit from early initiation of breastfeeding (INSAE & ICF, 2019). The median duration of exclusive breastfeeding (EBF) was 2.4 months in rural areas and 1.3 months in urban areas, and the rate of EBF till six months of age barely reached 42% (INSAE & ICF, 2019). The diversity of complementary foods was low with only 28% of 6–23 months children who met the minimum dietary diversity (MDD) (INSAE & ICF, 2019).

A diagnostic study conducted in two districts of southern Benin (namely *Bopa* and *Houéyogbé*) located in a high food-insecure area reported that infants and young children feeding practices were suboptimal (Mitchodigni, et al., 2017a; Mitchodigni, et al., 2017b). Authors suggested that multisectoral interventions should be implemented in this area to improve feeding practices for young children. Following this recommendation, a nutrition education program was planned in this region aiming at optimizing mothers' knowledge on nutrition and subsequently, breastfeeding and complementary feeding practices. Nutrition education is meant to improve knowledge, skills, motivation, and behavior of individuals or communities, leading potentially to subsequent positive effects on nutritional status and health (FAO, 2005). In sub-Saharan Africa, nutrition education interventions showed good contribution to improving breastfeeding practices (Aidam et al., 2005; Tylleskär et al., 2011), complementary feeding practices (Waswa et al., 2015) and also nutritional status of <2 years children (Lassi et al., 2013).

In order to refine the messages' content and take into account the contextual factors as part of the intervention, an in-depth understanding of the situation toward breastfeeding, complementary feeding practices, and associated factors in the area was necessary. Evidence from Benin and other African countries shows that multiple factors have the potential to affect positively or negatively breastfeeding and feeding practices among young children. These factors include, but are not limited to, cultural beliefs and habits (Amoussa Hounkpatin et al., 2014; Aborigo et al., 2012; Aryeetey & Goh, 2013; Issaka et al., 2014; Otoo et al., 2009), households' socioeconomic status (Agho et al., 2011; Mitchodigni, et al., 2017b; Sokan-Adeaga et al., 2019), agriculture practices within the households (Mitchodigni, et al., 2017b), mothers' overall instruction (Olatona et al., 2017; Qureshi et al., 2011; Sokan-Adeaga et al., 2019), mothers' occupation (Amoussa Hounkpatin et al., 2014; Mitchodigni, et al., 2017b; Nkrumah, 2017), as well as knowledge and perceptions of appropriate practices (Aborigo et al., 2012; Agunbiade & Ogunleye, 2012; Aidam et al., 2005; Issaka et al., 2014; Otoo et al., 2009; Qureshi et al., 2011).

The present study, which was conducted before the implementation of the nutrition education program, aimed at (a) characterizing

breastfeeding and complementary feeding practices in the intervention area using indicators which addressed directly the limitations among children feeding practices identified during the diagnostic survey and (b) identifying factors that were associated with these practices specifically in this intervention area. This could help to identify factors other than those relative to the intervention which could also influence the effect of the intervention by affecting the ability of the beneficiaries to adopt or reject recommended practices.

## 2 | MATERIALS AND METHODS

### 2.1 | Setting

The study was conducted in the districts of *Bopa* and *Houéyogbé* which present the highest rates of household food insecurity in the Mono department, respectively 40% and 34% (INSAE & PAM, 2014).

### 2.2 | Sampling

The data presented in this paper were related to the baseline survey of the impact evaluation of the nutrition education intervention implemented in the districts of *Bopa* and *Houéyogbé* in Southern Benin (Bodjrenou et al., 2020). A total of eight villages were randomly selected after stratification by district. In each village, 45 mother-child pairs were randomly selected from an exhaustive list of 0 to 17-months-old children living in the villages using a random number technique. We targeted this age-group firstly to focus on breastfeeding and complementary feeding practices. Secondly, the intervention and its impact evaluation were meant to last for 6 months; hence, children would be 6–23 months old at endline. In total, 360 mother-child pairs were surveyed.

### 2.3 | Data collection

Mothers or primary caregivers of children were interviewed during home visits by trained enumerators. Mothers' socioeconomic characteristics including age, ethnic group, employment status, marital status, level of education, and participation in nutrition education programs were collected. *Breastfeeding on demand* and attitudes toward *breastfeeding frequency during child illness* were recorded, while correct *positioning and attachment of the child during breastfeeding* was observed. We considered six parameters during the observations: breasts held by mother's hand, mother's hand doing a "C-shape," child's chin attached to the breast, child's lip in eversion, child in straight position, and child's whole body facing the mother's chest (Unicef, 2012; Vinther & Helsing, 1997; WHO & UNICEF, 1993). Enumerators observed mothers breastfeeding their children; each correct *breastfeeding positioning and attachment* parameter adopted by mothers was marked "1," and "0" otherwise.

The enumerators stayed in the study villages for about 2 weeks. This gave them several opportunities to observe mothers' practices. If an enumerator did not have the opportunity to observe a breastfeeding episode when interviewing a mother, she/he did not ask the mother to breastfeed her child, as this might disturb the habits. Enumerators had taken advantage of their stay in the village and would come back to make the observation as soon as he would have the opportunity to see the child at the breast. There is no specific duration of a breastfeeding episode. We have observed that youngest children breastfed longer and more frequently than older ones. Enumerators recorded different parameters of breastfeeding positioning and attachment as soon as mother positioned child and started breastfeeding.

The children aged 6–17 months feeding practices were also assessed using a qualitative 24 hr recall (WHO et al., 2008). All foods and drinks that had been used to feed the child the previous day and their constitutive ingredients were listed by the mother or the primary caregiver.

## 2.4 | Data management and statistical analysis

The following binary variables were computed and analysed: (a) *Breastfeeding on demand* (yes/no) (b) *Breastfeeding frequency when the children were ill* (increased versus unchanged or decreased). For each mother, a *breastfeeding positioning and attachment* score was computed by summing up the mark attributed to each of the six positioning parameters that were considered during the observations. The score, which theoretically ranged from 0 to 6 points, was recorded as a binary variable using the median score as a threshold: (a) mothers with a score less than the median and (b) mothers with a score equal or higher than the median (hence considered as “good positioning”).

From the 24 hr recall data, foods were categorized into seven food groups as recommended namely: (a) grains, roots, and tubers; (b) legumes and nuts; (c) dairy products; (d) flesh foods; (e) eggs; (f) vitamin-A rich products; (g) fruits and vegetables different from those rich in vitamin-A (Kennedy et al., 2010; WHO et al., 2008). The minimum dietary diversity (MDD), minimum meal frequency (MMF), and minimum acceptable diet (MAD) were computed following WHO and UNICEF guidelines (Kennedy et al., 2010; WHO et al., 2008). Children aged 6–23 months were considered having met MDD if they had consumed foods from at least four different food groups out of the seven recommended over the day prior to the survey. Children who received solid, semi-solid, or soft foods (including milk feeds for nonbreastfed children) the minimum number of times or more the day prior to the survey were considered having met MMF. A child who met both MDD and MMF was considered having met MAD (WHO et al., 2008). We, then, determined the proportions of children having met MDD, MMF, and MAD.

Characteristics of mothers and feeding practices indicators were described and presented using descriptive statistics: percentages

for categorical variables and mean  $\pm$  SD for continuous variables. Comparisons among the two districts were also performed using chi-square tests for categorical variables and Mann–Whitney test or Student's *t* tests for continuous variables. Associations between socioeconomic characteristics of mothers and breastfeeding and feeding practices outcomes were analysed through a two-step approach. Bivariate analyses (chi-square, Mann–Whitney, and Student's *t* tests) were first performed to identify factors with significant association with each practice. Variables that were significant at 20% (Mickey & Greenland, 1989; Trekpa et al., 2005) were kept for the next step. Secondly, we used generalized linear model (GLM) with binomial distribution probability and logit link function to build models presenting the contribution of the variables (those that were significant in bivariate analysis) to each outcome. All independent variables were entered together into each model in one step. Data presented included odd-ratios, confidence interval associated, *z*-value, and *p*-value.

The relevance of regression models was assessed by displaying the goodness of fit parameters (X-squared and *p*-value) associated with Hosmer–Lemeshow test (*p*-value should be higher than 5%) and the error rate (quality of model's predictions) derived from the confusion matrix (error rate should be <0.5). Where Hosmer–Lemeshow test was not significant, we performed Pearson's residuals test (*p*-value should be <5%).

Descriptive analyses and bivariate analyses were carried out using SPSS 23.0 (IBM-SPSS, 2014) while GLM and relative tests were conducted using R (Zuur et al., 2013). All analyses took into account the study design.

## 3 | RESULTS

### 3.1 | Socioeconomic characteristics of the sample

The average ages were  $8.4 \pm 4.9$  months for children and  $27.5 \pm 5.9$  years for mothers. Boys represented 53.1% of children in the sample. Most of mothers were from the *Sahouè* ethnic group, had no schooling, and lived with their husbands (Table 1). Children's characteristics (gender and age) were similar across districts. Mothers' age, marital status, and ethnic group were also similar across districts. However, the proportion of mothers with no schooling was significantly lower in the district of *Houéyogbé* than *Bopa*, with 45.0% and 77.2% respectively (*p*-value <.001).

Only 13.9% of mothers had no income generating activity. Mothers living in *Bopa* were mainly involved in agriculture (48.3%), trading (39.4%), and food processing (26.7%) while those living in *Houéyogbé* were involved in agriculture (27.2%), trading (23.9%), foods selling (24.4%), food processing (23.3%), and handicraft (20.6%) (Table 1).

Less than 6% of mothers had benefitted from a nutrition education program in the past with a difference between districts (*Bopa*: 2.2%; *Houéyogbé*: 9.4%; *p*-value =.003) (Table 1).

**TABLE 1** Socioeconomic characteristics of children and mothers ( $n = 360$ )

Parameters	n	All	Bopa	Houéyogbé	p-value
<b>Children characteristics</b>					
Age (months)	360	8.4 (4.9)	8.5 (4.8)	8.2 (5.0)	.578 <sup>S</sup>
Gender					
Male	191	53.1	50.6	55.6	.342 <sup>C</sup>
Female	169	46.9	49.4	44.4	
<b>Mothers characteristics</b>					
Age (years)	358 <sup>*</sup>	27.5 (5.9)	28.1 (5.9)	26.9 (5.8)	.054 <sup>S</sup>
Marital status					
Living alone	73	20.3	16.7	23.9	.088 <sup>C</sup>
Living with husband	287	79.7	83.3	76.1	
Formal education level					
No schooling	220	61.1	77.2	45.0	<.001 <sup>C</sup>
Literate or primary school	95	26.4	21.1	31.7	
Secondary school and more	45	12.5	1.7	23.3	
Ethnic group					
Sahoué	295	81.9	82.2	81.7	.89 <sup>1C</sup>
Other	65	18.1	17.8	18.3	
Employment status					
No activity	50	13.9	11.1	16.7	.128 <sup>C</sup>
Food processing	90	25.0	26.7	23.3	.465 <sup>C</sup>
Agriculture	136	37.8	48.3	27.2	<.001 <sup>C</sup>
Animal breeding	29	8.1	7.8	8.3	.846 <sup>C</sup>
Trading	114	31.7	39.4	23.9	.002 <sup>C</sup>
Foods selling	64	17.8	11.1	24.4	.001 <sup>C</sup>
Handicraft	45	12.5	4.4	20.6	<.001 <sup>C</sup>
Number of activities		1.4 (1.0)	1.5 (0.9)	1.3 (1.0)	.178 <sup>M</sup>
Attended nutrition education programs in the past					
At least once	21	5.8	2.2	9.4	.003 <sup>C</sup>
Never	339	94.2	97.8	90.6	

Note: Values presented are percentage for categorical variables and Mean (Standard Deviation) for continuous variables.  $p$ -values presented are probabilities relative to chi-square test (<sup>C</sup>) for categorical variables, Student's  $t$  test for Ages (<sup>S</sup>) and Mann-Whitney test (<sup>M</sup>) for Number of activities.

\*Two mothers did not know and could not estimate their exact age; therefore, we missed this information.

### 3.2 | Feeding practices

All children in our sample were breastfed. Percentage of mothers who used to breastfeed on demand their children were 58.7% (Figure 1); this proportion was slightly higher in *Bopa* than *Houéyogbé* (63.9% and 53.4%, respectively;  $p$ -value = 0.043). The proportion of mothers who had met *good positioning and attachment for breastfeeding* was 66.1% with no significant difference across districts (Figure 1). Only one-quarter of mothers declared increasing *breastfeeding frequency when their children were ill*.


The percentages of children meeting the requirements of MDD, MMF, and MAD were respectively 50.9%, 75.4%, and 43.5% (Figure 2). In *Houéyogbé*, 82.5% of children met the MMF while they were only 68.6% in *Bopa* ( $p$ -value = .015). The other indicators did not differ across districts.

### 3.3 | Factors associated with breastfeeding practices


Mothers who lived in *Houéyogbé* were less likely to breastfeed on demand (OR = 0.59, CI 95% = [0.36; 0.98],  $p = .04$ ) than mothers who lived in the district of *Bopa* (Table 2). Mothers who had higher school level (at least secondary school) were 3.08 more likely to breastfeed on demand than mothers with no schooling (OR = 3.08, CI 95% = [1.40; 6.79],  $p = .005$ ). Likewise, mothers who practiced trading and those who were not from the *Sahoué* ethnic group had higher odds to breastfeed on demand than others (Table 2).

Regarding good positioning and attachment for breastfeeding, only the children's age was associated with this indicator. Mothers of older children tended to have lower odds of having good positioning

**FIGURE 1** Breastfeeding practices in the sample and by district: Percentage of children achieving *Breastfeeding on Demand*, *Good Positioning and attachment for breastfeeding* and *Increased Breastfeeding Frequency during illness*. (% = Percentage; *p*-values presented are probabilities relative to chi-square test comparing the two districts)



**FIGURE 2** Complementary feeding practices in the whole sample and by district: Percentage of children achieving recommended complementary feeding practices. (% = Percentage; *p*-values presented are probabilities relative to chi-square test comparing the two districts)



and attachment for breastfeeding compared with others (OR = 0.94, CI 95% = [0.90; 0.99], *p* = .01) (Table 2).

Several factors were associated with the frequency of breastfeeding during illness episodes. Increased breastfeeding was more likely to happen among mothers who lived in Houéyogbé compared with those living in Bopa (OR = 2.16, CI 95% = [1.12; 4.18], *p* = .02), when children were older (OR = 1.10, CI 95% = [1.02; 1.19], *p* = .02), and when they were boys (OR = 0.51, CI 95% = [0.27; 0.96], *p* = .04). Unexpectedly higher the education level of mothers was, lesser was the likelihood to increase the frequency of breastfeeding illness episodes (OR = 0.19, CI 95% = [0.08; 0.48], *p* = .0004) (Table 2).

### 3.4 | Factors associated with complementary feeding practices

Very few factors were associated with the three indicators on complementary feeding practices (Table 3). Only the age of children was positively associated with the MDD (OR = 1.13,

CI 95% = [1.04; 1.23], *p* = .006) and the MAD (OR = 1.09, CI 95% = [1.01; 1.19], *p* = .04). Children who lived in Houéyogbé were more likely to meet the MMF compared to children who lived in Bopa (OR = 2.47, CI 95% = [1.24; 4.93], *p* = .01). However, the district was not associated with the MMD or the MAD. The variables related to the mothers' education and employment were not associated with any of the complementary feeding practices indicators.

## 4 | DISCUSSION

Results from our study showed that children feeding practices were suboptimal or medium and this situation requires actions for improvement. The definition of adequate interventions requires first to have a good knowledge of the situation and associated factors. Thus, we observed that age and sex of children, district of residence, ethnic group, education level, and employment status of mothers were associated with breastfeeding practices, with some variations depending on the indicators used. Mothers' education was positively

TABLE 2 Results from the GLM on breastfeeding practices

Parameters	OR	Confidence interval associated to OR		Z-value	p-value
		2.50%	97.50%		
Breastfeeding on demand ( $n = 356$ ; Hosmer and Lemeshow goodness of fit: X-squared = 11.03, $df = 8$ , $p$ -value = .1999; Error rate = 38.48)					
(Intercept)	0.62	0.33	1.19	-1.42	.15471
District_Houéyogbé (Ref = Bopa)	0.59	0.36	0.98	-2.04	.04104
Gender_Female (Ref = Male)	0.96	0.61	1.50	-0.19	.84630
Education_Literate or Primary school (Ref = No schooling)	1.44	0.84	2.45	1.34	.18083
Education_Secondary school (Ref = No schooling)	3.08	1.40	6.79	2.79	.00526
Ethnic group_Others (Ref = Sahouè)	1.93	1.04	3.57	2.10	.03583
Agriculture_Yes (Ref = No)	1.72	0.90	3.28	1.63	.10271
Trading_Yes (Ref = No)	1.89	1.07	3.35	2.20	.02817
Children's age (months)	1.05	1.00	1.10	1.91	.05591
Number of activities	1.00	0.70	1.42	-0.01	.99260
Good positioning and attachment for breastfeeding ( $n = 358$ ; Hosmer and Lemeshow goodness of fit: X-squared = 4.81, $df = 8$ , $p$ -value = .7775; Error rate = 32.68)					
(Intercept)	3.48	1.77	6.84	3.62	.00030
District_Houéyogbé (Ref = Bopa)	0.87	0.55	1.39	-0.57	.56751
Gender_Female (Ref = Male)	1.20	0.76	1.90	0.77	.44025
Ethnic group_Others (Ref = Sahouè)	1.53	0.82	2.85	1.33	.18406
Animal breeding_Yes (Ref = No)	0.47	0.18	1.23	-1.54	.12467
Trading_Yes (Ref = No)	0.61	0.36	1.05	-1.79	.07306
Children age	0.94	0.90	0.99	-2.47	.01369
Number of activities	1.14	0.84	1.55	0.86	.38758
Food selling_Yes (Ref = No)	0.54	0.29	1.00	-1.97	.04859
Increasing breastfeeding frequency during illness ( $n = 256$ ; Hosmer and Lemeshow goodness of fit: X-squared = 9.68, $df = 8$ , $p$ -value = .2881; Error rate = 26.17)					
(Intercept)	0.07	0.01	0.48	-2.71	.00674
District_Houéyogbé (Ref = Bopa)	2.16	1.12	4.18	2.29	.02179
Gender_Female (Ref = Male)	0.51	0.27	0.96	-2.10	.03585
Marital status_Living With husband (Ref = Living Alone)	1.88	0.78	4.54	1.40	.16268
Education_Literate or Primary school (Ref = No schooling)	0.19	0.08	0.48	-3.53	.00042
Education_Secondary school (Ref = No schooling)	0.54	0.19	1.57	-1.13	.25849
Food processing_Yes (Ref = No)	1.38	0.70	2.74	0.92	.35523
Animal breeding_Yes (Ref = No)	1.32	0.49	3.51	0.55	.58170
Children age	1.10	1.02	1.19	2.33	.01985
Mother age	1.01	0.95	1.07	0.29	.76961

Abbreviations: OR, odd-ratio; Ref, modality of reference.

associated with *breastfeeding on demand* but was negatively associated with *breastfeeding frequency during illness*. Educated mothers are likely to have a better knowledge of the importance of good breastfeeding practices (Al Ketbi et al., 2018), and we would expect better practices among them. On the other hand, educated mothers are also more likely to be employed in public or private companies

and they may have less time to apply good practices compared to mothers with informal jobs or no job at all. Employed mothers may also be stressed and tired, resulting in little time or energy at the end of the day to nurse their babies properly (Netshandama, 2002). Moreover, home-based activities like agriculture or small trading offer to mothers the opportunity to stay at home or nearby. Other

**TABLE 3** Results from the GLM on complementary feeding practices

Parameters	OR	Confidence interval associated to OR		Z-value	p-value
		2.50%	97.50%		
Minimum dietary diversity MDD (n = 230; Hosmer and Lemeshow goodness of fit: X-squared = 5.91, df = 8, p-value = .6573; Error rate 41.30)					
(Intercept)	0.37	0.12	1.14	-1.73	.08421
District_Houéyogbé (Ref = Bopa)	0.95	0.55	1.64	-0.17	.86438
Gender_Female (Ref = Male)	1.04	0.60	1.79	0.14	.89179
Nutrition education_Yes (Ref = No)	2.37	0.67	8.42	1.34	.18186
Food processing_Yes (Ref = No)	0.69	0.35	1.36	-1.06	.29011
Children age (months)	1.13	1.04	1.23	2.75	.00589
Number activities	0.82	0.59	1.14	-1.17	.24133
Minimum meal frequency MMF (n = 230; Hosmer and Lemeshow goodness of fit: X-squared = 10.76, df = 8, p-value = .2157; Error rate = 24.35)					
(Intercept)	1.64	0.51	5.32	0.83	.40960
District_Houéyogbé (Ref = Bopa)	2.47	1.24	4.93	2.57	.01020
Gender_Female (Ref = Male)	1.33	0.71	2.50	0.90	.37000
Education_Literate or Primary school (Ref = No schooling)	1.34	0.60	2.99	0.72	.47360
Education_Secondary school (Ref = No schooling)	0.63	0.22	1.78	-0.88	.38110
Children age	1.01	0.92	1.11	0.20	.84050
Minimum acceptable diet MAD (n = 230; Hosmer and Lemeshow goodness of fit: X-squared = 31.33, df = 8, p-value = .0001229; Pearson's Residuals: 0.3878219; Error rate = 38.26)					
(Intercept)	0.31	0.10	0.97	-2.01	.04430
District_Houéyogbé (Ref = Bopa)	1.20	0.69	2.09	0.64	.52060
Gender_Female (Ref = Male)	1.22	0.71	2.10	0.71	.47850
Food processing_Yes (Ref = No)	0.79	0.39	1.60	-0.65	.51370
Children age	1.09	1.01	1.19	2.07	.03810
Number of activities	0.79	0.56	1.11	-1.38	.16860
Food selling_Yes (Ref = No)	1.88	0.92	3.85	1.73	.08300

Abbreviations: OR, odd-ratio; Ref, modality of reference.

activities such as food selling require to leave home, and this may affect time spent with children and breastfeeding practices (Amoussa Hounkpatin et al., 2014; Nkrumah, 2017). Mother's marital status was not associated with breastfeeding practices in our study. Yet, a study in Ghana revealed better breastfeeding practices among women who lived with their husband (Rose, 2007). Indeed, when husbands have good financial situation, women could benefit from these resources and devote more time to nursing roles and less to income generating activities.

We also found that mothers living in Bopa, who were mostly involved in home-based activities, were more likely to breastfeed on demand than those living in Houéyogbé. The fact that Houéyogbé is more urbanized than Bopa may explain these results (Mitchodigni, et al., 2017b). These results are in line with national DHS (INSAE & ICF, 2019; INSAE, 2015) and other studies from low- and middle-income countries (Hitachi et al., 2019; Iffa & Serbesa, 2018; Kumar et al., 2017) which reported that most of breastfeeding

practices, especially *breastfeeding on demand* or EBF, were better in rural than urban areas. On the other hand, level of urbanization could be positively associated with other practices. For example, when a child is sick, Beninese mothers living in rural areas would preferentially treat their children with medicinal herbs in the form of tisanes (Allabi et al., 2011; Towns et al., 2014); the intakes of these tisanes could reduce breastmilk consumption. This echoed with our finding: mothers living in Houéyogbé were more likely to increase *breastfeeding frequency when children were ill* than mothers living in Bopa.

*Breastfeeding on demand* varied according to mothers' ethnic groups. Children from the *Sahouè* ethnic group were less likely to benefit from *breastfeeding on demand* than others. As shown in other sub-Saharan contexts (Asare et al., 2018; Jacdonmi et al., 2016; Tawiah-Agyemang et al., 2008; Wanjohi et al., 2017), social and cultural beliefs were determinant for breastfeeding practices. Finally, children's age was negatively associated with the likelihood to adhere

to the recommended positions for breastfeeding. Mothers explained during informal discussions that as children were growing up, they became stronger and more agitated; consequently, it was more difficult to control them during breastfeeding.

Regarding complementary feeding, as expected, we observed that children's age was positively associated with the MDD and MAD. Generally, mothers used to replace breast milk with porridge and then family foods progressively as children are growing up. Traditional porridges that are first given to young children consist of a cereal mixed with water and sometimes sugar, which provide low dietary diversity. The family meals that are eaten by older children offer higher dietary diversity. Thus, as children are growing up and start eating family meals, their dietary diversity increases (Amoussa Hounkpatin et al., 2014).

Living in Houéyogbé also seemed to be more favorable to higher meal frequency among children than living in Bopa; probably because of the difference of urbanization (Mitchodigni, et al., 2017b). No other factors were found to be associated with the complementary feeding indicators we investigated.

We observed that some intercepts were significant (*Increasing Breastfeeding Frequency during Illness, Good Positioning, and Attachment for Breastfeeding*, minimum acceptable diet). Thus, there were possibly other factors influencing feeding practices, but these factors were not collected in the present study.

Mitchodigni, et al. (2017b) showed also that agriculture, especially the diversity of food groups grown by households, increased the likelihood of meeting the MDD among 6–23 months old children. Moreover, we expected that participating in a nutrition education program in the past would have been associated with present feeding practices. Indeed, many studies had shown that nutrition education interventions targeting breastfeeding or dietary habits had good contribution to improving breastfeeding and complementary feeding practices in low- and middle-income countries especially in Africa (Aidam et al., 2005; Lassi et al., 2013; Tylleskär et al., 2011; Waswa et al., 2015). In other cases, nutrition education helped to increase knowledge and attitudes but not practices (Mojisola et al., 2019; Ruzita et al., 2007). However, in the present study few mothers (<6%) participated this program and this could not allow good analysis or conclusions. Moreover, participating a nutrition program does not necessarily lead to behavior change and achieving the adequate knowledge does not guarantee the adoption of good practices. Some other factors such as limited access to nutritious foods (Dang et al., 2005), financial constraints (Aborigo et al., 2012; Otoo et al., 2009), and pressures or support from families including food taboos (Kakute et al., 2005; Amoussa Hounkpatin et al., 2014; Issaka et al., 2015) play an important role. In the other hand, nutrition education programs have to be well planned and delivered (Chapman-Novakofski, 2014) and intensive (Bukusuba, 2005; Roy et al., 2005) to ensure their efficiency.

#### 4.1 | Limitation

We could have used qualitative method in the present study to have in-depth understanding of the situation. Qualitative research

approaches allow a detailed description of participants' feelings, opinions, experiences, and interpretations of their actions (Rahman, 2017). However, they have some limitations. Results cannot be generalized to the entire research population with the same degree of certainty as quantitative measures. Sample size used in qualitative research is often smaller than that used in quantitative research methods (Dworkin, 2012). Qualitative research results are not tested to determine whether they are statistically significant or due to chance (Atieno, 2009). Therefore, we decided to use quantitative method to ensure measurability and comparability (statistical) of indicators since we aim to collect data related to the same indicators at the end of the intervention.

## 5 | CONCLUSION

Breastfeeding and complementary feeding practices were not optimal in the districts of Bopa and Houéyogbé, department of Mono, Southern Benin. These results supported the importance of a nutrition education program at the community level in order to improve mothers' and community members' knowledge and attitudes toward children feeding practices. Moreover, socioeconomic, cultural, and demographic factors such as age and sex of children, district of residence, ethnic group, education level, and employment status identified in the present paper as influencing infants and children feeding practices will be taken into account when designing the community-based nutrition education program. According to these factors, different categories of people within our target population will be defined. Thus, key messages will be adapted to address each of these categories considering their specific backgrounds in terms of knowledge, occupation, food habits, culture, and beliefs.

#### ACKNOWLEDGMENTS

Research activities were funded by the Ministry of Foreign Affairs of Finland and CGIAR; The corresponding authors received also a PhD allowance from IRD.

#### CONFLICT OF INTEREST

"No conflicts of interest."

#### ETHICAL APPROVAL

The study was conducted in accordance with the Declaration of Helsinki. Ethical clearance was obtained from the Benin National Ethics Committee for Scientific Research (N°45/MS/DC/SGM/DFR/CNERS/SA). Administrative authorities of the two districts were informed and approved the study. Written informed consent of participants was obtained after receiving complete information about the study in the local language.

#### ORCID

Fifali Sam Ulrich Bodjrènou  <https://orcid.org/0000-0002-5619-3409>  
Waliou Amoussa Hounkpatin  <https://orcid.org/0000-0002-2902-8248>

Céline Termote  <https://orcid.org/0000-0003-3217-0226>  
 Mathilde Savy  <https://orcid.org/0000-0003-0562-9601>

## REFERENCES

- Aborigo, R. A., Moyer, C. A., Rominski, S., Adongo, P., Williams, J., Logonia, G., Affah, G., Hodgson, A., & Engmann, C. (2012). Infant nutrition in the first seven days of life in rural northern Ghana. *BMC Pregnancy Childbirth*, 12, 76. <https://doi.org/10.1186/1471-2393-12-76>
- Agho, K. E., Dibley, M. J., Odiase, J. I., & Ogbonmwan, S. M. (2011). Determinants of exclusive breastfeeding in Nigeria. *BMC Pregnancy and Childbirth*, 11, 2. <https://doi.org/10.1186/1471-2393-11-2>
- Agunbiade, M. O., & Ogunleye, V. O. (2012). Constraints to exclusive breastfeeding practice among breastfeeding mothers in south-west Nigeria: Implications for scaling up. *International Breastfeeding Journal*, 7, 5. <https://doi.org/10.1186/1746-4358-7-5>
- Aidam, A. B., Pérez-Escamilla, R., & Lartey, A. (2005). Lactation counseling increases exclusive breastfeeding rates in Ghana. *Community and International Nutrition*, 135(7), 1691–1695. <https://doi.org/10.1093/jn/135.7.1691>
- Al Ketbi, M. I., Al Noman, S., Al Ali, A., Darwish, E., Al Fahim, M., & Rajah, J. (2018). Knowledge, attitudes, and practices of breastfeeding among women visiting primary healthcare clinics on the island of Abu Dhabi, United Arab Emirates. *International Breastfeeding Journal*, 13, 26. <https://doi.org/10.1186/s13006-018-0165-x>
- Allabi, A. C., Busiac, K., Ekanmiana, V., & Bakiono, F. (2011). The use of medicinal plants in self-care in the Agonlin region of Benin. *Journal of Ethnopharmacology*, 133(11), 234–243. <https://doi.org/10.1016/j.jep.2010.09.028>
- Amoussa Hounkpatin, W., Blum, L., Ledain, A., & Badou-Savi, A. (2014). *Rapport final de l'enquête qualitative : Etude des déterminants des pratiques d'alimentation complémentaire des enfants de moins de 2 ans, Cas de 3 localités au Bénin: Boukoubé, Djakotomey et Kétou*. Cotonou, Benin: Ministère de la Santé du Bénin, Faculté des Sciences Agronomiques de l'Université d'Abomey-Calavi, UNICEF-Bénin.
- Aryeetey, R. N. O., & Goh, Y. E. (2013). Duration of exclusive breastfeeding and subsequent child feeding adequacy. *Ghana Medical Journal*, 47(1), 24–29. <https://pubmed.ncbi.nlm.nih.gov/23661852/>
- Asare, B.-Y.-A., Preko, J. V., Baafi, D., & Dwumfour-Asare, B. (2018). Breastfeeding practices and determinants of exclusive breastfeeding in a cross sectional study at a child welfare clinic in Tema Manhean, Ghana. *International Breastfeeding Journal*, 13, 12. <https://doi.org/10.1186/s13006-018-0156-y>
- Atieno, P. O. (2009). An analysis of the strengths and limitation of qualitative and quantitative research paradigms. *Problems of Education in the 21st Century*, 13, 13–18. <http://oaji.net/articles/2014/457-1393665925.pdf>
- Bodjrenou, F. S. U., Amoussa Hounkpatin, W., Mitchodigni, I., Salako, V., Glele Kakai, R., Dadele, Y., Ntandou Bouzitou, G., Schneider, L., Mutanen, M., Savy, M., Kennedy, G., Hounhouigan, J. D., & Termote, C. (2020). Comparing video and poster based education for improving 6–17 months children feeding practices: A cluster randomized trial in rural Benin. *Progress in Nutrition*, 22(1), 330–342. <https://doi.org/10.23751/pn.v22i1.9177>
- Bukusuba, J. (2005). Nutrition education: The vital step that will predict success of the fortification programme in Uganda. *African Journal of Food Agriculture and Nutritional Development*, 5, 1. <http://www.bioline.org.br/pdf?nd05016>
- Chapman-Novakofski, K. (2014). Does nutrition education "work"? *Journal of Nutrition Education and Behavior*, 46(2), 89. <https://doi.org/10.1016/j.jneb.2014.01.004>
- Dang, S., Yan, H., Yamamoto, S., Wang, X., & Zeng, L. (2005). Feeding practice among younger Tibetan children living at high altitudes. *European Journal of Clinical Nutrition*, 59(9), 1022–1029. <https://doi.org/10.1038/sj.ejcn.1602207>
- DeRose, L. F. (2007). Women's work and breastfeeding simultaneously Rise in Ghana. *Economic Development and Cultural Change*, 55(3), 583–612. <https://www.jstor.org/stable/10.1086/511193>
- Dworkin, S. L. (2012). Sample size policy for qualitative studies using in-depth interviews. *Archives of Sexual Behavior*, 41, 1319–1320. <https://doi.org/10.1007/s10508-012-0016-6>
- FAO (2005). *Nutrition education in primary schools* (Vol. 1). The Reader. Rome, Italy: FAO. ISBN 92-5-105454-1. P64. <http://www.fao.org/3/a-a0333e.pdf>
- Hitachi, M., Honda, S., Kaneko, S., & Kamiya, Y. (2019). Correlates of exclusive breastfeeding practices in rural and urban Niger: A community-based cross-sectional study. *International Breastfeeding Journal*, 14, 32. <https://doi.org/10.1186/s13006-019-0226-9>
- IBM-SPSS (2014). *IBM SPSS advanced statistics 23*. P126. IBM-SPSS.
- Iffa, M. T., & Serbesa, M. L. (2018). Assessment of the influence of mother's occupation and education on breastfeeding and weaning practice of children in public hospital, harari regional state Ethiopia. *Family Medicine and Medical Science Research*, 7, 3. <https://doi.org/10.4172/2327-4972.1000234>
- Institut National de la Statistique et de l'Analyse Économique (INSAE), & ICF (2019). *Enquête Démographique et de Santé au Bénin, 2017–2018* (pp. 1–675). Cotonou, Benin and Rockville, Maryland, USA: INSAE and ICF.
- Institut national de la Statistique et de l'Analyse Economique du Bénin (INSAE) (2015). *Enquête par grappes à indicateurs multiples 2014, Rapport final* (pp. 1–492). Cotonou, Bénin: INSAE.
- Issaka, A. I., Agho, K. E., Page, A. N., Burns, P. L., Stevens, G. J., & Dibley, M. J. (2015). The problem of suboptimal complementary feeding practices in West Africa: What is the way forward? *Maternal and Child Nutrition*, 11(Suppl 1), 53–60. <https://doi.org/10.1111/mcn.12195>
- Issaka, I. A., Agho, K. E., Page, A. N., Burns, P., Stevens, G. J., & Dibley, M. J. (2014). Determinants of early introduction of solid, semi-solid or soft foods among infants aged 3–5 months in four Anglophone West African Countries. *Nutrients*, 6(7), 2602–2618. <https://doi.org/10.3390/nu6072602>
- Jacdonmi, I., Suhainizam, M. S., Suriani, I. B., Zoakah, A. I., & Jacdonmi, G. R. (2016). Determinants of exclusive breastfeeding continuity among mothers of infants under six months in Plateau State, Nigeria. *International Journal of Health Sciences Research*, 6(4), 18–25. [https://pdfs.semanticscholar.org/f807/9bd86787dc567fac1781cb182941ab4d6c5b.pdf?\\_ga=2.95754460.1732896802.1591175391-1477554726.1551255001](https://pdfs.semanticscholar.org/f807/9bd86787dc567fac1781cb182941ab4d6c5b.pdf?_ga=2.95754460.1732896802.1591175391-1477554726.1551255001)
- Kakute, P. N., Ngum, J., Mitchell, J., Kroll, K. A., Forgwei, G. W., Ngwang, L. K., & Meyer, D. J. (2005). Cultural barriers to exclusive breastfeeding by mothers in a rural area of Cameroon. *Africa. Journal of Midwifery Womens Health*, 50(4), 324–328. <https://doi.org/10.1016/j.jmwh.2005.01.005>
- Kennedy, G., Ballard, T., & Dop, M.-C. (2010). *Guidelines for measuring household and individual dietary diversity* (pp. 1–56). Roma, Italy: Nutrition and Consumer Protection Division, FAO. ISBN: 9789251067499. <http://www.fao.org/publications/card/en/c/5aacbe39-068f-513b-b17d-1d92959654ea>
- Kumar, M. V., Acharya, I., Acharya, J. P., Shrivani, P. R., & Ramya, S. (2017). A comparison of breast feeding practices among urban and rural areas of Rangareddy district of Telangana. *International Journal Community Medicine and Public Health*, 4(6), 2136–2140. <https://doi.org/10.18203/2394-6040.ijcmph20172190>
- Lassi, Z. S., Das, J. K., Zahid, G., Imdad, A., & Bhutta, Z. A. (2013). Impact of education and provision of complementary feeding on growth and morbidity in children less than 2 years of age in developing countries: A systematic review. *BMC Public Health*, 13(Suppl 3), S13. <https://doi.org/10.1186/1471-2458-13-S3-S13>
- Mickey, R. M., & Greenland, S. (1989). The impact of confounder selection criteria on effect estimation. *American Journal of Epidemiology*,

- 129(1), 125–137. <https://doi.org/10.1093/oxfordjournals.aje.a115101>
- Mitchodigni, I. M., Amoussa Hounkpatin, W., Ntandou-Bouzitou, G., Termote, C., Kennedy, G., & Hounhouigan, D. J. (2017a). Complementary feeding practices of children aged 6–23 months in rural Area, Southern-Benin: Challenges and opportunities. *International Journal of TROPICAL DISEASE & Health*, 24(2), 1–12. <https://doi.org/10.9734/IJTDH/2017/33877>
- Mitchodigni, M., Amoussa Hounkpatin, W., Ntandou-Bouzitou, G., Avohou, H., Termote, C., Kennedy, G., & Hounhouigan, D. J. (2017b). Complementary feeding practices: Determinants of dietary diversity and meal frequency among children aged 6–23 months in Southern-Benin. *Food Security*, 9, 1117–1130. <https://doi.org/10.1007/s12571-017-0722-y>
- Mojisola, K., Macintyre, U. E., Gericke, G. J., & Becker, P. (2019). A contextual nutrition education program improves nutrition knowledge and attitudes of South African teachers and learners. *Frontiers in Public Health*, 7, 258. <https://doi.org/10.3389/fpubh.2019.00258>
- Netshandama, V. (2002). Breastfeeding practices of working women. *Curationis*, 25(1), 21–27. <https://doi.org/10.4102/curationis.v25i1.698>
- Nkrumah, J. (2017). Maternal work and exclusive breastfeeding practice: A community based crosssectional study in Efutu Municipal, Ghana. *International Breastfeeding Journal*, 12, 10. <https://doi.org/10.1186/s13006-017-0100-6>
- Olatona, F. A., Adenihun, J. O., Aderibigbe, S. A., & Adeniyi, O. F. (2017). Complementary feeding knowledge, practices, and dietary diversity among mothers of under-five children in an urban community in Lagos State, Nigeria. *International Journal of MCH and AIDS*, 6(1), 46–59. <https://doi.org/10.21106/ijma.203>
- Otoo, G. E., Lartey, A. A., & Pérez-Escamilla, R. (2009). Perceived incentives and barriers to exclusive breastfeeding among periurban Ghanaian women. *Journal of Human Lactation*, 25(1), 34–41. <https://doi.org/10.1177/0890334408325072>
- Programme Alimentaire Mondial, Service de l'Analyse de la Sécurité Alimentaire (VAM) (2014). *République du Bénin: Analyse Globale de la Vulnérabilité et de la Sécurité Alimentaire (AGVSA)* (p. 1–146). Rome, Italie: PAM.
- Qureshi, A. M., Oche, O. M., Sadiq, U. A., & Kabiru, S. (2011). Using community volunteers to promote exclusive breastfeeding in Sokoto State, Nigeria. *Pan African Medical Journal*, 10, 8. <https://doi.org/10.4314/pamj.v10i0.72215>
- Rahman, S. (2017). The advantages and disadvantages of using qualitative and quantitative approaches and methods in language “testing and assessment” research: a literature review. *Journal of Education and Learning*, 6(1), 102–112. <https://files.eric.ed.gov/fulltext/EJ1120221.pdf>
- Roy, S. K., Fuchs, G. J., Mahmud, Z., Ara, G., Islam, S., Shafique, S., Akter, S. S., & Chakraborty, B. (2005). Intensive nutrition education with or without supplementary feeding improves the nutritional status of moderately-malnourished children in Bangladesh. *Journal of Health Population Nutrition*, 23(4), 320–330.
- Ruzita, A. T., Wan Azdie, M. A. B., & Ismail, M. N. (2007). The effectiveness of nutrition education programme for primary school children. *Malaysian Journal of Nutrition*, 13(1), 45–54.
- Sokan-Adeaga, M. A., Sokan-Adeaga, A. A., & Sokan-Adeaga, E. D. (2019). A systematic review on exclusive breastfeeding practice in Sub-Saharan Africa: Facilitators and barriers. *Acta Scientific Medical Sciences*, 3(7), 53–65.
- Tawiah-Agyemang, C., Kirkwood, B. R., Edmond, K., Bazzano, A., & Hill, Z. (2008). Early initiation of breast-feeding in Ghana: Barriers and facilitators. *Journal of Perinatology*, 28, S46–S52. <https://doi.org/10.1038/jp.2008.173>
- Towns, A. M., Mengue Eyi, S., & van Anel, T. (2014). Traditional medicine and childcare in Western Africa: Mothers' knowledge, folk illnesses, and patterns of healthcare-seeking behavior. *PLoS One*, 9(8), e105972. <https://doi.org/10.1371/journal.pone.0105972>
- Trekpa, M. J., Pekovic, V., Santana, J. C., & Zhang, G. (2005). Risk factors for lead poisoning among cuban refugee children. *Public Health Reports*, 120(2), 179–185. <https://doi.org/10.1177/003335490512000212>
- Tylleskär, T., Jackson, D., Meda, N., Engebretsen, I. M. S., Chopra, M., Diallo, A. H., Doherty, T., Ekström, E.-C., Fadnes, L. T., Goga, A., Kankasa, C., Klungsoyr, J. I., Lombard, C., Nankabirwa, V., Nankunda, J. K., Van de Perre, P., Sanders, D., Shanmugam, R., Sommerfelt, H., ...PROMISE-EBF Study Group (2011). Exclusive breastfeeding promotion by peer counsellors in sub-Saharan Africa (PROMISE-EBF): A cluster-randomised trial. *Lancet*, 378(9789), 420–427. [https://doi.org/10.1016/S0140-6736\(11\)60738-1](https://doi.org/10.1016/S0140-6736(11)60738-1)
- UNICEF (2012). *Key messages booklet: The community infant and young child feeding counselling package.*, (pp. 1–45). UNICEF. [https://www.unicef.org/nutrition/files/Key\\_Message\\_Booklet\\_2012\\_small.pdf](https://www.unicef.org/nutrition/files/Key_Message_Booklet_2012_small.pdf)
- Vinther, T., & Helsing, E. (1997). *Breastfeeding, how to support success: A practical guide for health workers.* (p. 1–85). World Health Organization Regional Office For Europe Copenhagen.
- Wanjohi, M., Griffiths, P., Wekesah, F., Muriuki, P., Muhia, N., Musoke, R. N., Fouts, H. N., Madise, N. J., & Kimani-Murage, E. W. (2017). Sociocultural factors influencing breastfeeding practices in two slums in Nairobi. *Kenya. International Breastfeeding Journal*, 12, 5. <https://doi.org/10.1186/s13006-016-0092-7>
- Waswa, L. M., Jordan, I., Herrmann, J., Krawinkel, M. B., & Keding, G. B. (2015). Community-based educational intervention improved the diversity of complementary diets in western Kenya: Results from a randomized controlled trial. *Public Health Nutrition*, 18(18), 3406–3419. <https://doi.org/10.1017/S13688980015000920>
- WHO & UNICEF (1993). *Breastfeeding counselling a training course. Participants' manual: Part one, Sessions 1–9* (pp. 171). WHO/UNICEF. [https://www.who.int/maternal\\_child\\_adolescent/documents/pdfs/bc\\_participants\\_manual.pdf?ua=1](https://www.who.int/maternal_child_adolescent/documents/pdfs/bc_participants_manual.pdf?ua=1)
- WHO, Unicef, USAID, AED, UCDAVIS, IFPRI (2008). Indicators for assessing infant and young child feeding practices. Part I: Definitions (pp. 19). OMS and UNICEF. ISBN: 978 92 4 159666 4.
- Zuur, A. F., Hilbe, J. M., & Ieno, E. N. (2013). *Beginner's Guide to GLM and GLMM with R. Ed* (pp. 1–256). Highland Statistics Ltd. ISBN: 9780957174139.

**How to cite this article:** Bodjrenou FSU, Amoussa Hounkpatin W, Termote C, Dato G, Savy M. Determining factors associated with breastfeeding and complementary feeding practices in rural Southern Benin. *Food Sci Nutr*. 2020;00:1–10. <https://doi.org/10.1002/fsn3.1971>