

## P49

Maternal and Perinatal Complications of Vascular Pathologies Associated with Pregnancy:  
Frequency and Associated Factors in a Third Level Maternity in Benin

Tognifode Veronique\*, Aboubakar Moufalilou, **Ogoudjobi Mathieu**, Dakin Mohamed,  
Dangbemey Patrice, Gayito René, Hounkponou Fanny, Tshabu Christiane, Hounkpatin Benjamin,  
Denakpo Jutsin Lewis

Journal of Gynecology and Obstetrics 2022; 10(2): 60-68

doi: 10.11648/j.jgo.20221002.12

<https://www.sciencepublishinggroup.com/journal/paperinfo?journalid=255&doi=10.11648/j.jgo.20221002.12>

---

|                          |   |
|--------------------------|---|
| Nom de la revue          | : Journal of Gynecology and Obstetrics  |
| Portée de la revue       | : Internationale  |
| Type (papier / en ligne) | : Papier / en ligne   |
| Comité de lecture        | : Oui   |
| ISBN – ISSN              | : 2376-7812 / 2376-7820   |
| Editeur                  | : Science Publishing Group<br><a href="http://www.sciencepublishinggroup.com/j/jgo">http://www.sciencepublishinggroup.com/j/jgo</a>   |
| Impact factor            | : 0.807   |
| Indexation               | : Academickeys; CNKI SCHOLAR; ResearchBib; Universal Impact Factor; WorldCat<br><a href="http://www.sciencepublishinggroup.com/journal/indexing?journalid=255">http://www.sciencepublishinggroup.com/journal/indexing?journalid=255</a> |

---



SciencePG

Science Publishing Group



Journal of  
**Gynecology  
and Obstetrics**



Science Publishing Group

## Journal of Gynecology and Obstetrics

[Home](#) | [Archive](#) | [Special Issues](#) | [Indexing](#) | [Editorial Board](#) | [Reviewers](#) | [Submission Guidelines](#) | [Article Processing](#)

[Home](#) / [Journals](#) / [Medicine, Health & Food](#) / [Journal of Gynecology and Obstetrics](#)



Journal of  
**Gynecology  
and Obstetrics**



Science Publishing Group

Vol. 10, Issue 3, May 2022

ISSN Print: 2376-7812

Frequency: Bimonthly

ISSN Online: 2376-7820

[Recommend to Library](#)

<https://www.sciencepublishinggroup.com/j/jgo>

[Latest Articles](#)

[About This Journal](#)

- ▶ **Clinical Value of the N-S-P Scheme for Diagnosing Vulvar Dermatitis (DATRIV Study, Part 2)**  
Vesna Harni, Damir Babic, Suzana Ljubojevic-Hadzavdic, Dubravko Barisic

## Journal of Gynecology and Obstetrics

[Home](#) | [Archive](#) | [Special Issues](#) | [Indexing](#) | [Editorial Board](#) | [Reviewers](#) | [Submission Guidelines](#) | [Article Processing Charges](#) | [Publication Ethics](#)

### Archive

2022, Volume 10

2021, Volume 9

Vol. 9, Issue 6, Nov.

Vol. 9, Issue 5, Sep.

Vol. 9, Issue 4, Jul.

Vol. 9, Issue 3, May

Vol. 9, Issue 2, Mar.

Vol. 9, Issue 1, Jan.

[Home](#) / [Journals](#) / [Medicine, Health & Food](#) / [Journal of Gynecology and Obstetrics](#) / [Indexing](#)

## Indexing

*Journal of Gynecology and Obstetrics* has been included by the following Abstracting and Indexing databases:

[Academickeys](#)

[CrossRef](#)

[Eurasian Scientific Journal Index](#)

[MIAR](#)

[ResearchBib](#)

[Wissenschaftszentrum Berlin](#)

[CNKI SCHOLAR](#)

[Directory of Research Journals Indexing](#)

[EZB](#)

[Polish Scholarly Bibliography](#)

[Universal Impact Factor](#)

[WorldCat](#)

# Maternal and Perinatal Complications of Vascular Pathologies Associated with Pregnancy: Frequency and Associated Factors in a Third Level Maternity in Benin

Tognifode Veronique<sup>1,\*</sup>, Aboubakar Moufalilou<sup>1</sup>, **Ogoudjobi Mathieu<sup>1</sup>**, Dakin Mohamed<sup>1</sup>, Dangbemey Patrice<sup>1</sup>, Gayito René<sup>2</sup>, Hounkponou Fanny<sup>2</sup>, Tshabu Christiane<sup>1</sup>, Hounkpatin Benjamin<sup>1</sup>, Denakpo Jutsin Lewis<sup>1</sup>

<sup>1</sup>Mother and Child Department, Faculty of Health Sciences, University of Abomey-Calavi, Cotonou, Benin

<sup>2</sup>Department of Obstetrics Gynecology, Tanguiéta Zone Hospital, Tanguiéta, Benin

## Email address:

verofode@yahoo.fr (T. Veronique)

\*Corresponding author

## To cite this article:

Tognifode Veronique, Aboubakar Moufalilou, Ogoudjobi Mathieu, Dakin Mohamed, Dangbemey Patrice, Gayito René, Hounkponou Fanny, Tshabu Christiane, Hounkpatin Benjamin, Denakpo Jutsin Lewis. Maternal and Perinatal Complications of Vascular Pathologies Associated with Pregnancy: Frequency and Associated Factors in a Third Level Maternity in Benin. *Journal of Gynecology and Obstetrics*. Vol. 10, No. 2, 2022, pp. 60-68. doi: 10.11648/j.jgo.20221002.12

**Received:** January 28, 2022; **Accepted:** February 21, 2022; **Published:** March 9, 2022

---

**Abstract:** Background and aims: Vascular pathologies (hypertensive disorders and diabetes) associated with pregnancy are important public health problem in Benin because of their frequencies and their complications. This study aims to analyze maternal and perinatal complications of vascular pathologies associated with pregnancy. Methods: This analytical cross-sectional study used retrospective data collected over a three-year period. All mothers who presented a vascular pathology and their newborns during the study period at the Clinique Universitaire de Gynécologie obstétrique (CUGO) of Centre National Hospitalier Universitaire Hubert Koutoukou Maga (CNHU-HKM) of Cotonou were included. Maternal complications associated neurological, hematological, cardio-pulmonary and renal complications whereas perinatal complications included intrauterine growth restriction, fetal asphyxia, prematurity, neonatal resuscitation, low birthweight and perinatal death. Factors associated to the complications were identified by using chi-square test and a multivariate binary logistic regression model. Analyzes were performed with SAS software version 9.4. Results: The frequency of vascular pathologies associated with pregnancy was 14.95%. Diabetes was associated in 4.57%. Their evolutions were marked by the occurrence of numerous maternal complications dominated by retroplacental hematoma (12.08%), eclampsia (08.37%) which passed through signs of eclampsism (21.07%); and hellp syndrome. Perinatal complications were dominated by prematurity (43.07%), low birthweight (29.12%), fetal asphyxia (12.32%) and perinatal death (18.59%). These complications were associated with the precocity of the diagnosis, admission mode and the importance of proteinuria. Conclusion: Maternal and perinatal complications of vascular pathologies associated with pregnancy are frequent in current obstetric practice. It is therefore urgent to implement health promotion interventions targeting the factors associated with them in order to reverse the trends.

**Keywords:** Complications, Maternal, Perinatal, Vascular Pathologies, Pregnancy

---

## 1. Background

Hypertensive disorders and diabetes associated with pregnancy are major public health problem not only because of their high prevalence but also because of their maternal

and fetal consequences. According to the World Health Organization (WHO), hypertensive disorders of pregnancy (HDP) are the second leading cause of maternal death and the first cause of perinatal morbidity and mortality in the world [1]. Their incidence varies from 10 to 15% [2]. This incidence differs according to the level of development of

the countries with a rate between 10.6% in the United States, 7% in Canada, 9% in China [3-5]. In developing countries, the frequencies are higher, 17% in Guinea, 12.3% in Lomé, 12.5% in Benin [5-7]. These vascular pathologies are a major cause of maternal and perinatal morbidity and mortality. They are the cause of 18% of all maternal deaths [1] after immediate post-partum hemorrhage. However, there are disparities between countries. United Nations statistics report a rate of 16% of maternal deaths attributable to hypertensive disorders in high level income countries [7]. In Asia and Africa, nearly 10% of maternal deaths are associated with HDP. The complications which lead to this death are: eclampsia, HELLP syndrome, retroplacental hematoma (RPH), acute pulmonary edema (APO), and strokes.

The complications of vascular pathologies can hold back on the fetus and the newborn. These complications are dominated by prematurity, perinatal asphyxia, intrauterine growth restriction (IUGR) and fetal death in utero. In France, the neonatal mortality rate due to vascular pathologies is 12.5% [8]. In Africa, a case-control study carried out in Niamey (Niger) showed a fetal mortality rate four times higher in the group of hypertensive pregnant women compared to the group of normotensive pregnant women [8].

In addition to complications, several factors are associated with poor perinatal prognosis of vascular pathologies in low-income countries. These are young maternal age, existence of a pathological obstetrical history, absence of antenatal care visits (ANC), a number of ANC less than 4, poor surveillance, low gestational age at the time of the diagnosis of the disease, delay in the decision to terminate the pregnancy, delay in referral to level II or III maternity hospitals [9].

Besides HDP, diabetes, whether pre-existing or not, can complicate pregnancies. The prevalence of gestational diabetes is increasing worldwide with rates ranging from 1.8% to 23% depending on the study population and diagnostic criteria [10]. A 2017 WHO study showed that the incidence of gestational diabetes was estimated at 16.2% in pregnant women aged 20-40 years [11]. In the same study, it was shown that 21.4 million children born to diabetic mothers with more than 90% from low-income countries had neonatal complications. The most common fetal complications were macrosomia, IUGR, prematurity, congenital malformations, especially cardiovascular and renal, and chromosomal anomalies. Macrosomia is the most frequent complication [12]. Its risk is doubled in cases of gestational diabetes compared to a normal pregnancy. Its incidence in newborns of diabetic mothers varies between 17 and 30%, compared to only 10% in the general population. Newborns from diabetic mothers are more able to have fetal trauma (skull and clavicle fractures) and metabolic disorders such as hypoglycemia in the first hours of life and hypocalcemia [13, 14].

In a precarious health environment, such as that of BENIN, where the medical devices are insufficient, it is essential to screen the population of pregnant women at risk of

hypertension or diabetes in order to improve the prognosis of the pregnancy with a multidisciplinary approach. Several studies have been done on hypertensive disorders and diabetes associated with pregnancy; however, very few have linked patient characteristics to maternal and perinatal outcome. The objective of this study is to analyze maternal and perinatal complications of vascular pathologies associated with pregnancy in a third level maternity hospital in Cotonou (Benin) by determining 1) their frequency and 2) their associated factors.

## 2. Methodology

### 2.1. Data

This analytic cross-sectional study with retrospective data collection over a 3-year period, from January 1, 2016, to December 31, 2018 included the records of all mothers with vascular pathology and their newborns during the study period at the Clinique Universitaire de Gynécologie obstétrique (CUGO) of Centre National Hospitalier Universitaire Hubert Koutoukou Maga (CNHU-HKM) of Cotonou which is a third level maternity. The records of patients who had given birth but had not developed vascular pathologies during pregnancy, delivery, or postpartum were ruled out. The same applies to newborns of mothers who had a vascular pathology associated with pregnancy but did not deliver at CUGO (outborn). Permission from the heads of the CUGO and pediatric departments of the CNHU-HKM were obtained. Data were collected with respect to confidentiality and anonymity, both for the babies and their parents.

### 2.2. Variables

Dependent variable was the maternal and perinatal complications of vascular pathologies. Independent variables were related to sociodemographic characteristics (age, occupation, education level, residence); clinical characteristics (obstetric, medical, surgical, and family history, mode and reason for admission, type of delivery); physical examination data (blood pressure, weight, height, body mass index, level of consciousness using GLASGOW scale, neurosensory signs, lower limb edema, uterine height, fetal heart sounds, albuminuria); paraclinical data. For children, variables were related to clinical characteristics (birth term, weight, sex, notion of resuscitation, fetal asphyxia, prematurity, low birthweight, respiratory distress, hypothermia). Maternal complications included: neurological (eclampsia, stroke), hematological (retroplacental hematoma, HELLP syndrome), cardiopulmonary and renal complications. Perinatal complications included IUGR, fetal asphyxia, prematurity, neonatal resuscitation, hypotrophy and perinatal death. The different maternal and perinatal complications were used as binary variables.

### 2.3. Statistical Analyses


Study sample was described in terms of sociodemographic

characteristics (age, occupation, level of education, residence), mode and reason for admission, obstetrical and medical-surgical history. Obstetrical examination, paraclinical check-up, delivery modalities as well as the maternal and perinatal complications were also described. Quantitative variables were expressed as mean with standard deviation and qualitative variables as proportions with confidence intervals. Factors associated with maternal and perinatal complications were identified by using bivariate analyses to select candidate variables (chi-2 test for qualitative variables), and then a multivariate model was performed by binary logistic regression model. Analyses were performed with SAS software 9.4 (SAS Institute, Inc, Cary, NC). A p value less than 5% was significant.

### 3. Results

#### *Sociodemographic and clinical characteristics of patients*

This study included 1291 cases of vascular pathologies associated with pregnancy out of a total of 8634 deliveries, i.e. a hospital frequency of 14.95%. Pre-eclampsia and gestational hypertension were the most frequent clinical types (Figure 1) with overall frequencies of 63.5% and 30.36% respectively. Diabetes was associated in 4.57% of cases.



**Figure 1.** Distribution of hypertensive disorders of pregnancy at CUGO of CNHU-HKM from 2016 to 2018 (n=1291).

The average age of the patients was 28.32 years, with the 18 to 35 age group being the most represented. As shown in Table 1, the majority of patients were not educated (43.99%) and lived in urban areas in more than half of the cases. The main reasons for referral were pre-eclampsia, lumbopelvic pain and hypertension. The majority of patients were pauciparous with a short spacing interval between births. Pregnancy was well attended in 45% of cases with 59.41% of patients who delivered by caesarean section.

**Table 1.** Socio-demographic and clinical characteristics of patients.

|                  | Numbers | Percentages (%) |
|------------------|---------|-----------------|
| Age groups       |         |                 |
| Youngs           | 34      | 02.62           |
| Middle           | 1040    | 80.56           |
| Olds             | 192     | 14.87           |
| Missing          | 25      | 01.94           |
| Education        |         |                 |
| No education     | 568     | 43.99           |
| Primary level    | 285     | 22.08           |
| Secondary level  | 266     | 20.60           |
| University level | 172     | 13.33           |
| Residence        |         |                 |
| Urban            | 746     | 57.78           |
| Rural            | 545     | 42.22           |
| Admission mode   |         |                 |
| Herself          | 318     | 24.60           |
| Referred         | 973     | 75.40           |
| Admission reason |         |                 |
| Lumbopelvic pain | 325     | 25.17           |
| Hypertension     | 221     | 17.12           |
| Pre-eclampsia    | 356     | 27.57           |
| Eclampsia        | 118     | 09.14           |
| Loss of fluid    | 61      | 04.72           |
| Convulsions      | 54      | 04.18           |
| Delivery labor   | 161     | 12.47           |
| Metrorrhagia     | 108     | 08.36           |
| Fetal asphyxia   | 115     | 08.90           |
| Gestation        |         |                 |
| First gestation  | 401     | 31.00           |
| Second gestation | 422     | 32.70           |
| Multigestation   | 468     | 36.30           |
| Parity           |         |                 |
| No delivery      | 127     | 14.32           |
| One              | 224     | 25.22           |
| Two              | 361     | 40.65           |
| ≥ 3              | 176     | 19.81           |

|                                    | Numbers | Percentages (%) |
|------------------------------------|---------|-----------------|
| Spacing interval between births    |         |                 |
| Short                              | 38      | 07.27           |
| Normal                             | 485     | 92.73           |
| Antenatal care visits (ANC)        |         |                 |
| < 4 ANC                            | 374     | 54.28           |
| ≥ 4 ANC                            | 315     | 45.72           |
| Gestational week (GW) at diagnosis |         |                 |
| Before 34 GW                       | 537     | 41.60           |
| Between 34 et 37 GW                | 225     | 17.43           |
| More than 37 GW                    | 559     | 40.98           |
| Number of fetuses                  |         |                 |
| Singleton                          | 1176    | 91.10           |
| Twin or multiple                   | 115     | 08.90           |
| Type of delivery                   |         |                 |
| Vaginal                            | 524     | 40.59           |
| Cesarian section                   | 767     | 59.41           |

*Maternal and perinatal complications of vascular pathologies associated with pregnancy and associated factors*

Table 2 shows that the main maternal complications were retroplacental hematoma (12.08%) and eclampsia (08.37%)

between those who had signs of eclampsism (21.07%). Perinatal complications in this study were mostly prematurity (43.07%), low birthweight (29.12%), fetal asphyxia (12.32%) and perinatal death (18.59%).

**Table 2.** Maternal and perinatal complications of vascular pathologies associated with pregnancy at the CUGO of CNHU-HKM from 2016 to 2018.

|   | Numbers | Percentages (%) |
|---|---------|-----------------|
| Maternal complications                  |         |                 |
| <i>Eclampsism</i>                       |         |                 |
| Yes                                     | 272     | 21.07           |
| No                                      | 1019    | 78.93           |
| <i>Eclampsia</i>                        |         |                 |
| Yes                                     | 108     | 08.37           |
| No                                      | 1183    | 91.63           |
| <i>Stroke</i>                           |         |                 |
| Yes                                     | 08      | 00.62           |
| No                                      | 1283    | 99.38           |
| <i>Retroplacental hematoma</i>          |         |                 |
| Yes                                     | 156     | 12.08           |
| No                                      | 1135    | 87.92           |
| <i>HELLP syndrome</i>                   |         |                 |
| Yes                                     | 25      | 01.94           |
| No                                      | 1266    | 98.06           |
| <i>Acute pulmonary edema</i>            |         |                 |
| Yes                                     | 11      | 00.85           |
| No                                      | 1280    | 99.15           |
| <i>Acute kidney failure</i>             |         |                 |
| Yes                                     | 38      | 02.94           |
| No                                      | 206     | 15.95           |
| Missing                                 | 1047    | 81.11           |
| Perinatal complications                 |         |                 |
| <i>Prematurity</i>                      |         |                 |
| Yes                                     | 556     | 43.07           |
| No                                      | 553     | 42.83           |
| Missing                                 | 182     | 14.10           |
| <i>Neonatal resuscitation</i>           |         |                 |
| Yes                                     | 101     | 07.82           |
| No                                      | 881     | 68.24           |
| Missing                                 | 309     | 23.94           |
| <i>Low birthweight</i>                  |         |                 |
| Yes                                     | 376     | 29.12           |
| No                                      | 612     | 47.41           |
| Missing                                 | 303     | 23.47           |
| <i>Intra-uterine growth restriction</i> |         |                 |
| Yes                                     | 55      | 04.26           |
| No                                      | 1236    | 95.74           |

|                        | Numbers | Percentages (%) |
|------------------------|---------|-----------------|
| <i>Fetal asphyxia</i>  |         |                 |
| Yes                    | 159     | 12.32           |
| No                     | 1132    | 87.68           |
| <i>Perinatal death</i> |         |                 |
| Yes                    | 240     | 18.59           |
| No                     | 1051    | 81.41           |

As shown in Table 3 below, in bivariate analysis, the frequency of maternal complications was significantly higher in middle-aged patients, in cases of significant proteinuria, in patients referred, in patients with poor antenatal care and in those who had caesarean section. On the other hand, the occurrence of perinatal complications

decreased significantly with the term at diagnosis. In multivariate analysis, only the effects of significant proteinuria (OR=5.83; 95% CI [2.85-11.9]) and term at diagnosis persisted with a linear trend of significant reduction of the risk of maternal and perinatal complications in the latter (OR=0.49; 95% CI [0.34-0.70]).

**Table 3.** Factors associated with maternal complications of vascular pathologies associated with pregnancy at the CUGO of the CNHU-HKM.

|                                    | Maternal complications |              | p-value | OR*  | 95%CI*       |
|------------------------------------|------------------------|--------------|---------|------|--------------|
|                                    | Yes                    | No           |         |      |              |
| Age groups                         |                        |              | 0.04    |      |              |
| Youngs                             | 16 (03.14%)            | 18 (02.38%)  |         | 1.26 | [0.66-2.30]  |
| Middle                             | 431 (84.68%)           | 609 (80.45%) |         | 1    |              |
| Olds                               | 62 (12.18%)            | 130 (17.17%) |         | 0.77 | [0.38-1.55]  |
| Type of hypertensive disorders     |                        |              | <0.0001 |      |              |
| Without significant praterinuria   | 62 (12.04%)            | 360 (46.39%) |         | 1    |              |
| With significant praterinuria      | 453 (87.96%)           | 416 (53.61%) |         | 5.83 | [2.85-11.92] |
| Admission mode                     |                        |              | <0.0001 |      |              |
| Herself                            | 71 (13.79%)            | 247 (31.83%) |         | 1    |              |
| Referred                           | 444 (86.21%)           | 529 (68.17%) |         | 1.12 | [0.55-2.27]  |
| First delivery                     |                        |              | 0.12    |      |              |
| No                                 | 387 (75.15%)           | 553 (71.26%) |         | 1    |              |
| Yes                                | 128 (24.85%)           | 223 (28.74%) |         | 0.96 | [0.51-1.80]  |
| Gestational week (GW) at diagnosis |                        |              | <0.0001 |      |              |
| Before 34 GW                       | 128 (24.85%)           | 401 (51.68%) |         | 1    |              |
| Between 34 et 37 GW                | 92 (17.86%)            | 133 (17.14%) |         | 1.70 | [1.20-3.81]  |
| More than 37 GW                    | 295 (57.28%)           | 242 (31.19%) |         | 4.14 | [2.02-8.50]  |
| <i>Test for linear trend</i>       |                        |              |         | 0.49 | [0.34-0.70]  |
| Type of delivery                   |                        |              | 0.002   |      |              |
| Vaginal                            | 183 (35.53%)           | 435 (56.06%) |         | 1    |              |
| Cesarian section                   | 332 (64.47%)           | 341 (43.94%) |         | 1.67 | [0.88-3.18]  |
| Spacing interval between births    |                        |              | 0.55    |      |              |
| Short                              | 181 (91.98%)           | 304 (93.25%) |         | 1    |              |
| Normal                             | 16 (8.12%)             | 22 (6.75%)   |         | 0.95 | [0.35-2.58]  |
| Antenatal care visits (ANC)        |                        |              | 0.001   |      |              |
| < 4 ANC                            | 93 (37.65)             | 222 (50.23%) |         | 1    |              |
| ≥4 ANC                             | 154 (62.35%)           | 220 (49.77%) |         | 1.04 | [0.58-1.87]  |

Abbreviations:

OR: Odd ratio

CI: Confidence intervals

\*Multivariable logistic regression model

Bold indicates statistical significance

As shown in Table 4 below, the frequency of perinatal complications was significantly higher in patients with significant proteinuria, in referred patients, in patients with less antenatal care visits or short spacing interval between births, and in those who delivered by cesarean section. On the

other hand, their frequencies decreased significantly with the term at diagnosis. In multivariate analysis, the effects of mode of admission, significant proteinuria, and earliness of diagnosis persisted with a significant linear trend test for the latter (OR=0.06 IC95% [0.02-0.14]).

**Table 4.** Factors associated with perinatal complications of vascular pathologies associated with pregnancy at the CUGO of the CNHU-HKM.

|            | Perinatal complications |              | p-value | OR*  | CI*         |
|------------|-------------------------|--------------|---------|------|-------------|
|            | Yes                     | No           |         |      |             |
| Age groups |                         |              | 0.14    |      |             |
| Youngs     | 24 (02.85%)             | 10 (02.36%)  |         | 1.33 | [0.75-2.15] |
| Middle     | 702 (83.37%)            | 338 (79.72%) |         | 1    |             |

|                                    | Perinatal complications |              | p-value | OR*         | CI*                |
|------------------------------------|-------------------------|--------------|---------|-------------|--------------------|
|                                    | Yes                     | No           |         |             |                    |
| Olds                               | 116 (13.78%)            | 76 (17.92%)  |         | 0.71        | [0.28-1.80]        |
| Type of hypertensive disorders     |                         |              | <0.0001 |             |                    |
| Without significant praterinaria   | 182 (21.26%)            | 240 (55.17%) |         | 1           |                    |
| With significant praterinaria      | 674 (78.74%)            | 195 (44.83%) |         | 3.22        | [1.43-7.26]        |
| Admission mode                     |                         |              | <0.0001 |             |                    |
| Herself                            | 149 (17.41%)            | 169 (38.85%) |         | 1           |                    |
| Referred                           | 707 (82.59%)            | 266 (61.15%) |         | 2.49        | [1.12-5.53]        |
| First delivery                     |                         |              | 0.06    |             |                    |
| No                                 | 637 (74.42%)            | 303 (69.66%) |         | 1           |                    |
| Yes                                | 219 (25.58%)            | 132 (30.34%) |         | 0.70        | [0.31-1.61]        |
| Gestational week (GW) at diagnosis |                         |              | <0.0001 |             |                    |
| Before 34 GW                       | 175 (20.44%)            | 354 (81.38%) |         | 1           |                    |
| Between 34 et 37 GW                | 215 (25.12%)            | 10 (02.30%)  |         | 20.70       | [10.2-30.81]       |
| More than 37 GW                    | 466 (54.44%)            | 71 (16.32%)  |         | 44.18       | [12.6-154.8]       |
| <i>Test for linear trend</i>       |                         |              |         | <i>0.06</i> | <i>[0.02-0.14]</i> |
| Type of delivery                   |                         |              | 0.25    |             |                    |
| Vaginal                            | 338 (39.49%)            | 186 (42.76%) |         | 1           |                    |
| Cesarian section                   | 518 (60.51%)            | 249 (57.24%) |         | 0.94        | [0.43-2.06]        |
| Spacing interval between births    |                         |              | 0.03    |             |                    |
| Short                              | 298 (90.85%)            | 187 (95.90%) |         | 1           |                    |
| Normal                             | 30 (09.15%)             | 08 (04.10%)  |         | 1.24        | [0.28-5.32]        |
| Antenatal care visits (ANC)        |                         |              | 0.02    |             |                    |
| < 4 ANC                            | 189 (42.57%)            | 126 (51.43%) |         | 1           |                    |
| ≥4 ANC                             | 225 (57.43%)            | 119 (48.57%) |         | 1.24        | [0.59-2.63]        |

Abbreviations:

OR: Odd ratio

CI: Confidence intervals

\*Multivariable logistic regression model

Bold indicates statistical significance.

## 4. Discussion

Hypertensive disorders of pregnancy and diabetes are frequent in the world, regardless of the level of development of the countries, but with an increasing trend in low-income countries. The frequency of HDP in Africa varies from 12.3% [15] in Togo to 17.5% in Guinea [16]. Our frequency of 14.95% falls within this range. In general, the rates reported in high-income countries are low, ranging from 7 to 10% in Canada and the United States [4, 17, 18]. However, there is little data on all HPD with pregnancy, which makes comparisons difficult. The available data are mostly related to pre-eclampsia, which is the main clinical form of hypertension associated with pregnancy (63.5% of all forms of hypertension in our study). Like the association between hypertension and pregnancy, pre-eclampsia is more frequent in low-income countries than in high-income ones, with rates varying from 3 to 7%, particularly in the United States, the United Kingdom and France [18-20]. However, there are limitations to this comparison, as the rates calculated in high-income countries are related to the total number of pregnancies, whereas in low-income countries they are expressed in relation to the number of deliveries, which is the case in our study.

Besides HDP, the association of diabetes and pregnancy represents a potential risk for both mother and child. The frequency of diabetes associated with pregnancy is very diversely assessed in the literature; it varies from 1% to 17% according to Western and Asian studies and from 5% to 38% according to African studies [20]. Our frequency of 4.57%

takes into account gestational diabetes and type 2 diabetes. The tendency of Africans to adopt a westernized diet combined with a sedentary lifestyle explains the increase in diabetes rates.

The average age of the patients was  $28 \pm 6.36$  years. According to the authors, these ages correspond to the period of maximum fertility [21]. In Africa, the age of childbearing is lower because of early marriage and unschooling of young girls. However, this age tends to increase, as is the case in northern countries where the age of childbirth is decreasing. The patients did not attend school in 43.99% of cases, corresponding to an unfavorable socioeconomic profile. Multigestation and pauciparous women were the most numerous in our study with respective rates of 36.3% and 40.6%. The same result was found in other African samples [22, 23].

HDP discovering in this study was late in almost half of the cases. In fact, it was the term at which the diagnosis was made, explained by the poor monitoring of the pregnancy and late referral of the patients. In fact, three quarters of the patients in this study were referred from local hospitals, health centers or private clinics when they presented a hypertensive attack before or during labor. There were no ANV or this one was irregular in 54.3% of cases. The situation is similar for most low-income countries. Lower rates of ANC were found in two studies from Senegal (65.4%) and Morocco (60.4%) [22, 23]. Studies have already identified the factors that explain the low use of prenatal services. These are economic and socio-cultural factors and poor health coverage in low-income countries [24, 25].

Eclampsia is the first complication of HDP. It is also the main complication in most African and Western samples, 77.8% in Côte d'Ivoire [26], 32.35% in Morocco [09] and 10.75% in Japan [27]. The differences between the series reported in the literature are related to the level of health development of the countries with a much lower frequency of eclampsia in high-income countries. In addition to eclampsia, retroplacental hematoma (RPH) appears to be the second most common complication of HDP, 12.08% in this study. This rate is close to that reported in Mali (11.6%). These two main complications associated with hellp syndrome, which is still under-diagnosed in our working conditions, explain the lethality rate of HDP, which is 1.54% in this study. Obstetrically, cesarean section was the main delivery method (59.4%). Indeed, most of the patients presented a severe clinical picture, endangering the vital prognosis of the mother and the fetus, which indicated that a vaginal delivery was not appropriate.

HDP has also fetal and neonatal consequences. Fetal consequences include IUGR, acute fetal asphyxia, in utero death, and prematurity. IUGR is the result of chronic fetal suffering and can lead to very low birthweight. In this study, the rate was 4.26%. Higher rates have been found in other African series, 18.1% in Nigeria [28], 19.4% in Congo [29]. In general, IUGR rates are lower in series from high-income countries, thus the rate of IUGR from vascular causes was 7.2% in the United States [30], and 9% in Switzerland [31]. When considering the different rates reported in the literature, the IUGR rate in our study would be comparable to those of Western series. However, this rate is underestimated in our study firstly due to selection bias and secondly to the diagnostic criteria for IUGR which lack precision in most African series. Besides IUGR, one of the frequent complications of vascular pathologies associated with pregnancy is prematurity (43.07% in this study). This is a prematurity caused by maternal and/or fetal complications such as eclampsia, hellp syndrome, IUGR, hyperuricemia greater than 80 mg/L, abnormalities in the fetal heartbeat and pathological umbilical and cerebral dopplers. In-utero death occurs at the end of a severe IUGR or during an acute event (eclampsia, retroplacental hematoma, hellp syndrome). In our series, perinatal mortality was 18.59%, higher than in some African studies 11.3% in Morocco [09], 13.2% in Congo [29].

Significant proteinuria was associated with maternal and perinatal complications. Indeed, it increases the risk of eclampsia, retroplacental hematoma, and hellp syndrome and leads to induced prematurity with a high morbidity and mortality. Other studies had already reached the same conclusions [32, 33]. Moreover, early diagnosis was associated with the occurrence of maternal and perinatal complications. Indeed, it has been shown that early forms of vascular pathologies associated with pregnancy are more serious [34]. This severity associated in our working conditions with the diagnostic delay linked to the late recourse to care explains the unfavorable maternal and perinatal

outcome. Perinatal complications were significantly more frequent in referred patients. This is related to the status of the CNHU of Cotonou which is the center of last resort in BENIN health pyramid. It was not possible in this study to specify whether there were dysfunctions in the referral that could have an impact on maternal and perinatal prognosis. However, studies have already shown that referrals in health systems in developing countries are characterized by the first and second delays [35].

This retrospective study has inherent shortcomings in this type of data collection, particularly in most developing countries. Out of 1533 records expected from the delivery registers, only 1400 records were found, including 109 records that could not be used because of the absence of important data such as maternal and perinatal prognosis. These deficiencies are related to poor archiving of data and lack of computerization of records. This may result in an underestimation of the true frequency of vascular pathologies associated with pregnancy, and of the proportions of the different variables studied. Ideally, this selection bias should have been explored by comparing those excluded with those included. However, as the number of files retained was comparable to what is observed in similar studies, this exclusion was not likely to significantly influence the trend of our results. Despite these limitations, the strengths of this study include the large sample size and the analysis methodology (logistic regression) to control for potential confounders.

## 5. Conclusion

At the end of this hospital study, it was found that the frequency of vascular pathologies associated with pregnancy was 14.95%. Their evolution was marked by the occurrence of numerous maternal complications dominated by retroplacental hematoma, eclampsia which was characterized by signs of eclampsism, and hellp syndrome. Perinatal complications were mostly prematurity, low birthweight, fetal asphyxia and perinatal death. These different complications were associated with the earliness of the diagnosis, the mode of admission and the importance of proteinuria. Thus, early management of these pathologies will improve the maternal and perinatal prognosis. Prenatal consultations are therefore essential to detect the first evocative clinical signs in order to limit maternal and perinatal morbidity.

## References

- [1] Khan KS, Wojdyla D, Say L, Gülmezoglu A. M., Van Look P. F. WHO analysis of causes of maternal death: a systematic review. *The Lancet*. 2006; 367 (9516): 1066-74. DOI: 10.1016/s0140-6736(06)68397-9.
- [2] Abalos E, Cuesta C, Grosso AL, Chou D, Say L. Global and regional estimates of preeclampsia and eclampsia: a systematic review. *EJOG*. 2013; 170 (1): 1-7. DOI: 10.1016/j.ejogrb.2013.05.005.

- [3] Mogos MF, Salemi JL, Spooner KK, McFarlin BL, Salihu HH. Hypertensive disorders of pregnancy and postpartum readmission in the United States: national surveillance of the revolving door. *Journal of Hypertension*. 2018; 36 (3): 608-18. DOI: 10.1097/HJH.0000000000001594.
- [4] Public Health Agency of Canada. Canadian Chronic Disease Indicators. [En ligne]. 2016 [cité le 26 févr 2021]. Disponible: <https://health-infobase.canada.ca/ccdi/>.
- [5] Bah AO, Diallo AAS, Diallo MH, Keita N, Diallo MS. Hypertension artérielle et grossesse: aspects épidémiologiques et facteurs de risque. *MedAfrNoire*. [En ligne]. 2000 [cité 26 Mar 2021]; (47): 422-25. Disponible: <http://www.santetropicale.com/Resume/104704.pdf>.
- [6] Baragou S, Goeh-Akue E, Pio M, Afassinou YM, Atta B. Hypertension artérielle et grossesse à Lomé (Afrique sub-saharienne): aspects épidémiologiques, diagnostiques et facteurs de risque. *Annales de Cardiologie et d'Angéologie*. 2014; 63 (3): 145-50. DOI: 10.1016/j.ancard.2014.05.006.
- [7] Mounier-Vehier C, Amar J, Boivin J-M, Denolle T, Fauvel J-P, Plu-Bureau G, et al. Hypertension artérielle et grossesse: Consensus d'experts de la Société française d'hypertension artérielle, filiale de la Société française de cardiologie. *Presse médicale*. 2016; 45 (7-8P1): 682-99. DOI: 10.1016/j.lpm.2016.05.012.
- [8] Minani PG. HTA (hypertension artérielle) et grossesse au CHU de Kamenge. *Mémoire Online*. [En ligne]. 2011 [cité le 26 févr 2021]. Disponible: <https://www.memoireonline.com/04/12/5666/HTA-hypertension-artérielle-et-grossesse-au-CHU-de-Kamenge.html>.
- [9] El-M'Zah S, Aboufalah A. Pronostic maternel et périnatal de l'hypertension artérielle gravidique [Doctorat en médecine en ligne]. Marrakech: Université de CADI AYYAD, Faculté de médecine et de pharmacie; 2010 [Cité le 26 févr 2020]. Disponible: <http://wd.fmpm.uca.ma/biblio/theses/annee-hm/FT/2010/these112-10.pdf>.
- [10] Pirson N, Maiter D, Alexopoulou O. Prise en charge du diabète gestationnel en 2016: une revue de la littérature. *Endocrinologie et Nutrition*. [En ligne]. 2016 [cité le 26 févr 2021]. Disponible: <https://www.louvainmedical.be/sites/default/files/content/article/pdf/lmed-12-2016-01-pirson.pdf>.
- [11] Fédération internationale du diabète. Atlas du diabète de la FID. *International Diabetes Federation*. 7e édition [En ligne]. 2017 [cité le 08 Jan 2021]. Disponible: [https://www.federationdesdiabetiques.org/sites/default/files/fid/documents/idf\\_atlas\\_8e\\_fr.pdf](https://www.federationdesdiabetiques.org/sites/default/files/fid/documents/idf_atlas_8e_fr.pdf).
- [12] Mitanchez D. Complications fœtales et néonatales du diagnostic gestationnel: mortalité périnatale, malformations congénitales, macrosomie, dystocie des épaules, traumatisme obstétrical, complications néonatales. *Journal de Gynécologie Obstétrique et Biologie de la Reproduction*. 2010; 39 (8S2): 189-99. DOI: 10.1016/S0368-2315(10)70046-6.
- [13] Billionnet C, Mitanchez D, Weill A, Nizard J, Alla F, Hartemann A, et al. Gestational diabetes and adverse perinatal outcomes from 716, 152 births in France in 2012. *Diabetologia*. 2017; 60 (4): 636-44. DOI: 1007/s00125-017-4206-6.
- [14] Varzos MS. Incidence du diabète sur la grossesse. *Diabetologia*. [En ligne] 2018 [cité le 12 Jan 2021] 61 (8): 6-7. Disponible: [http://www.uclmontgodinne.be/files/2018-05-31\\_colloque\\_maternite\\_incidence\\_m\\_s.varzos](http://www.uclmontgodinne.be/files/2018-05-31_colloque_maternite_incidence_m_s.varzos).
- [15] Campbell DM, MacGillivray I, Carr-Hill R. Pre-eclampsia in second pregnancy. *Br J Obstet Gynaecol*. 1985; 92 (2): 131 40. DOI: 10.1111/j.1471-0528.1985.tb01064.x.
- [16] Sibai BM, el-Nazer A, Gonzalez-Ruiz A. Severe preeclampsia-eclampsia in young primigravid women: subsequent pregnancy outcome and remote prognosis. *Am J Obstet Gynecol*. 1986; 155 (5): 1011 6. DOI: 10.1016/0002-9378(86)90336-4.
- [17] Makkonen N, Heinonen S, Kirkinen P. Obstetric prognosis in second pregnancy after preeclampsia in first pregnancy. *Hypertens Pregnancy*. 2000; 19 (2): 173 81. DOI: 10.1081/prg-100100133.
- [18] Dukler D, Porath A, Bashiri A, Erez O, Mazor M. Remote prognosis of primiparous women with preeclampsia. *Eur J Obstet Gynecol Reprod Biol*. 2001; 96 (1): 69 74. DOI: 10.1016/s0301-2115(00)00392-4.
- [19] Davies AM, Czaczkes JW, Sadovsky E, Prywes R, Weiskopf P, Sterk VV. Toxemia of pregnancy in Jerusalem. I. Epidemiological studies of a total community. *Isr J Med Sci*. [En ligne]. 1970; 6 (2): 253 66. Disponible: <https://www.ncbi.nlm.nih.gov/pubmed/5506239>.
- [20] Baniyas BB, Devoe LD, Nolan TE. Severe preeclampsia in preterm pregnancy between 26 and 32 weeks' gestation. *Am J Perinatol*. 1992; 9 (5 6): 357 60. DOI: 10.1055/s-2007-999264.
- [21] Arngrimsson R, Björnsson S, Geirsson RT, Björnsson H, Walker JJ, Snaedal G. Genetic and familial predisposition to eclampsia and pre-eclampsia in a defined population. *Br J Obstet Gynaecol*. 2012; 97 (9): 762 9. DOI: 10.1111/j.1471-0528.1990.tb02569.x.
- [22] Maxwell CV, Lieberman E, Norton M, Cohen A, Seely EW, Lee-Parritz A. Relationship of twin zygosity and risk of preeclampsia. *Am J Obstet Gynecol*. 2001; 185 (4): 819 21. DOI: 10.1067/mob.2001.117352.
- [23] Savvidou MD, Karanastasi E, Skentou C, Geerts L, Nicolaides KH. Twin chorionicity and pre-eclampsia. *Ultrasound Obstet Gynecol*. 2001; 18 (3): 228 31. DOI: 10.1046/j.0960-7692.2001.00470.x.
- [24] Garner PR, D'Alton ME, Dudley DK, Huard P, Hardie M. Preeclampsia in diabetic pregnancies. *Am J Obstet Gynecol*. 2011; 163 (2): 505 8. DOI: 10.1016/0002-9378(90)91184-e.
- [25] McCowan LM, Buist RG, North RA, Gamble G. Perinatal morbidity in chronic hypertension. *Br J Obstet Gynaecol*. 2006; 103 (2): 123 9. DOI: 10.1111/j.1471-0528.1996.tb09662.x.
- [26] Pattison NS, Chamley LW, McKay EJ, Liggins GC, Butler WS. Antiphospholipid antibodies in pregnancy: prevalence and clinical associations. *Br J Obstet Gynaecol*. 2013; 100 (10): 909 13. DOI: 10.1111/j.1471-0528.1993.tb15105.x.
- [27] Yasuda M, Takakuwa K, Tokunaga A, Tanaka K. Prospective studies of the association between anticardiolipin antibody and outcome of pregnancy. *Obstet Gynecol*. 2005; 86 (4 Pt 1): 555 9. DOI: 10.1016/0029-7844(95)00247-0.
- [28] Olusanya BO, Solanke OA. Perinatal Outcomes Associated with Maternal Hypertensive Disorders of Pregnancy in a Developing Country. *Hypertension in Pregnancy*. 2012; 31 (1): 120 30. DOI: 10.3109/10641955.2010.525280.

- [29] Mabaga JM, Mubinda PK, Mavuta CZ, Mukuku O, Kakudji PL, Luboya ON. Pronostic périnatal de l'association hypertension artérielle et grossesse aux Cliniques Universitaires de Lubumbashi. ResearchGate. [En ligne]. 2017 [cité 11 janv 2021]. Disponible: <https://www.researchgate.net/publication/323417054>.
- [30] Netgen. Retard de croissance intra-utérin: impact sur le développement et la fonction cérébrale [Internet]. Rev Méd Suisse. [cité 10 Jan 2021]. Disponible: <https://www.revmed.ch/RMS/2008/RMS-146/Retard-de-croissance-intra-uterin-impact-sur-le-developpement-et-la-fonction-cerebrale>.
- [31] Retard de croissance intra-utérin: recommandations pour la pratique clinique – Texte court. Journal de Gynécologie Obstétrique et Biologie de la Reproduction. 2013; 42 (8): 1018-25. DOI: 10.1016/j.jgyn.2013.09.023.
- [32] Witlin AG, Saade GR, Mattar F, Sibai BM. Risk factors for abruptio placentae and eclampsia: analysis of 445 consecutively managed women with severe preeclampsia and eclampsia. Am J Obstet Gynecol. 1999; 180 (6 Pt 1): 1322-9. DOI: 10.1016/s0002-9378(99)70014-1.
- [33] Tikkanen M. Placental abruption: epidemiology, risk factors and consequences. Acta Obstet Gynecol Scand. 2011; 90 (2): 140-9. DOI: 0.1111/j.1600-0412.2010.01030.
- [34] Céline Guillemot. Pré-éclampsie sévère avant 34 SA: diagnostic, prise en charge et pronostic maternofoetal. État des lieux au CHRU de Brest de janvier 2011 à décembre 2014. Sciences du Vivant [q-bio]. 2016. ffdumas-01560524f.
- [35] Sombié I, Méda ZC, Savadogo LBG, Somé DT, Bamouni SF, Dadjoari M, et al. La lutte contre la mortalité maternelle au Burkina Faso est-elle adaptée pour réduire les trois retards? Sante Pub.