



GSJ: Volume 6, Issue 9, September 2018, Online: ISSN 2320-9186

[www.globalscientificjournal.com](http://www.globalscientificjournal.com)

## SOME TRADITIONAL FERMENTED FOOD FROM AFRICAN LOCUST BEANS (*PARKIA BIGLOBOSA*): PRODUCTION AND VOLATILE COMPOUNDS

Oscar ZANNOU<sup>1\*</sup>, Ifagbémi Bienvenue CHABI<sup>2</sup>, Ikay KOCA<sup>1</sup>

<sup>1</sup>Department of Food Engineering, Ondokuz Mayıs University, 55139, Samsun, Turkey

<sup>2</sup>School of Nutrition, Sciences and Food Technologies, University of Abomey-Calavi, P.O. Box 526, Cotonou 01, Republic of Benin

\*Corresponding author: [zannouoscar@gmail.com](mailto:zannouoscar@gmail.com)

### Abstract

African locust bean is used for both medicinal and domestic purposes in West Africa. The seeds are processed to nutritive and flavoring fermented condiments such as *netetu*, *iru*, *sonru*, *afitin*, *soumbala*, and *dawadawa*. The production process and the volatile compounds of these fermented condiments were investigated in the present study. The findings showed that dried and cleaned seeds were used to be produced traditional condiments. The processing operations implying mainly and successively a first long boiling (6-24h), dehulling, a second short cooking (1-2h) and fermentation (24-72h). The variations of the operations involving in the production affect physicochemical, biochemical and microbiological characteristics of the final products. More than 160 volatile compounds namely pyrazines, ketones, aldehydes, alcohols, esters, alkanes, alkenes, benzene derivatives, pyridines, furan, volatile phenols, sulfur compounds, and terpenes have been reported in fermented African locust beans. Among these, pyrazines, ketones, aldehydes, and alcohols were found to be the most abundant groups with 2,5-Dimethyl pyrazine, tetramethyl pyrazine, trimethyl pyrazine, 3-methyl butanal, benzene acetaldehyde, 2-nonadecanone, 2-decanone, 3,5-Dimethyl phenylmethanol and 3-methyl-1-butanol as the most important aroma compounds found in the condiments. Production steps mainly fermentation (type and duration), boiling and use of additives conferred to each fermented seed its unique aroma characteristics. Use of pure starter culture enhances overall aroma profile of the finished products.

**Keywords:** Fermentation, *Parkia biglobosa*, fermented condiments, aroma, West Africa.

## 1. Introduction

African traditional culinary background is greatly bound by an enormous variety of fermented foods. These concern mainly cereals, tubers, dairy products and some non-timber forest products such as African locust beans. Fermentation is one of the most important steps intervening during the processing of traditional foods in Africa. It is considered as a process which involves microorganisms growth and their metabolic activities [1]. When fermentation occurring, microorganisms transform the available hydrocarbons into organic acids, alcohols, carbon dioxide as well as bacteriocins [2,3]. Bacteriocin designates the small, heat-stable cationic peptides produced by lactic acid bacteria, which displays a strong inhibition power as it kills or inhibits pathogen microorganism, thus assuring food stability, shelf-life, safety [2,4], as well as overall quality. To producers, the process of fermentation is cultural, cheaper, less energy consumption and also needs a short time to be achieved. Fermentation ensures not only the preservation of the processed foods by increasing their shelf life but also increases their flavor quality and nutritional values. Preservative effect of fermentation is usually linked with the generation of the antimicrobials. According to the definition used by the Commission of the European Communities (EU Directive 95/2/EC), preservatives are substances that extend the shelf life of foodstuffs by protecting them against deterioration caused by microorganisms [5]. Similarly, the U.S. Food and Drug Administration (FDA) defines preservatives as any chemical that when added to food tends to prevent or retard deterioration [5]. Natural antimicrobials can be defined as substances produced by living organisms in their fight with other organisms for space and their competition for nutrients and the main sources of these compounds are plants (secondary metabolites in essential oils and phytoalexins), microorganisms (bacteriocins and organic acids), and animals (lysozyme from eggs, lactoferrins from milk). In one hand, the fermentation renders more digestible the raw products [6] and does not request a specific expertise or expensive materials. In another hand, it increases the bioavailability of nutrients such as proteins, lipids, hydrocarbons, minerals, and vitamins. According to [7], the fermentation contributes to the removing of non-useful substances. The fermented foods are associated with numerous health profits as they can not only prevent osteoporosis, allergies, hypertension, carcinogenesis, obesity, intestinal troubles, diabetes and atherosclerosis but also can decrease blood cholesterol and enhance human immunity [8,9,10]. Furthermore, anti-oxidant, anti-microbial, anti-fungal, anti-inflammatory, anti-diabetic and antiatherosclerotic effects are conferred to the fermented products [7].

The genus *Parkia* from the subfamily of *Mimosoideae* which, in return, belongs to the large family of *Fabaceae*, is composed of thirty-four plant species [11]. African locust beans (*Parkia biglobosa*) appertains to the genus of *Parkia*. For West African natives the whole tree of *Parkia biglobosa*, from roots to leaves, is useful and can be utilized either as foodstuffs or in traditional medicine. For instance, a survey performed in the Republic of Benin revealed that the fruits particularly seeds and pulps are the most used, followed by barks, branches, leaves, roots and flowers [12]. Domestical use of this tree has a limitless importance for Africa Savanna's populations. Many studies reported the use of *P. biglobosa* in endogenous arenas. To Kwon-Ndung et al [13], beside the medicinal use, *P. biglobosa* is used as firewood, forage, varnish and for charcoal production, whereas, Irvine [14] and Hagos [15] mentioned *P. biglobosa* wood drifted products which encompass bows, hoe handles, mortarboards, seats, and drumsticks. Its yellow pulps are used as flour for the preparation of enriched porridges and local meals. Bark infusions are used to relieve diseases like febrifuge, toothache, bronchitis, diarrhea, tracheitis, venereal diseases, pneumonia as well as "Cure salée" [16]. Some communities in Ghana use roots, leaf, barks and fruits to recover for bums, fever, boil, diarrhea, and stomach disorders [17]. Furthermore, in the Republic of Benin, *P. biglobosa* is mostly utilized for folk medicine and foods but it is also employed for its medico-magic and cultural characteristics [12]. Leaves, roots, pods and stems of this plant contain important bioactive compounds essentially saponins, tannins, steroids, flavones, glycosides, alkaloids and flavonoids [18,19,20,21,22]. For the same authors, extracts of different parts such as roots, leaves, pod, stem and barks of *P. biglobosa* have been successfully tested on an important number of pathogenic microorganisms including *Enterobacter aerogenes*, *Streptococcus aureus*, *Staphylococcus aureus*, *Salmonella typhi*, *Escherichia coli*, *Salmonella typhimurium*, *Pseudomonas aeruginosa*, *Proteus mirabilis*, *Shigella spp*, *Rhizopus spp*, *Klebsiella sp* and *Mucor spp*. The presence of bioactive compounds and strong antimicrobial properties detected in *Parkia biglobosa* witnessed its remedy value and importance of human health.

In Africa, the most of fermented foods or beverages are obtained through a spontaneous or natural fermentation. Technologies for the production of fermented foods or beverages are mastered by producers, however, they vary depending on cultural background, commodities and geographical situation. Thus, *mawè* [23], *Tchoukoutou* [24], *lafoun*, *koko* [25], *afitin*, *iru* and *sonru* [26], *kisra* [27], *dawadawa* [28], *dèguè* [29], *soumbala* [30], *fura*, *lanhouin*, and so forth, are some fermented foods and beverages encountered in Africa. Foods

can be subjected to three types of fermentation namely alkali, lactic acid and alcoholic. These have been combined in two major classes such as aerobic fermentation which includes alkaline and fungal, and anaerobic fermentation which comprises lactic and alcoholic [2]. According to [31], alcoholic fermentation generates ethanol and yeasts as main metabolites and includes mostly wines and beers, while, lactic acid fermentation is usually implemented with lactic acid bacteria and implies mainly cereals, milk and somehow tubers. For the same author, alkali fermentation occurs often in seeds and fishes condiments. Seeds of *Parkia biglobosa* are the most promoted and used to produce both human and animal alimentation. Seeds were detected as a valuable source of nutrients especially proteins for poussins [32,33] and fishes [34]. For human consumption, the seeds are absolutely cooked and fermented in order not only to eliminate antinutritional factor but to improve nutritional, preservative and organoleptic qualities. Thus, the fermented seeds are recognized as a pleasant and nutritious condiment used in various African dishes. They often represent significant substitutes of general proteins sources such as fishes, eggs and animal proteins for the populations of rural, suburban as well as urban zones. The name, cooking and fermentation conditions of the resulted fermented foods, vary extensively according to production regions or countries. For instance, they are known as *iru*, *sonru* and *afitin* in the Republic of Benin [26], *dawadawa* or *daddawa*, *iru* and *ogiri-igala* in Nigeria [35,36], *dawadawa*, *iru* and *kpalugu* in Ghana [35,36], *kinda* in Sierra leone, *netelou* or *soumbara* in Gambia [36], *soumbala* in Burkina Faso [37], *netetu* in Senegal [38], *soumbara* in Ivory Coast and *dawadawa* in Chad. These names are accompanied by the variability of production procedure which confers to each product a characteristic flavor.

The flavor which is defined as a mixture of taste and odor (aroma), is a determinant characteristic that discriminates a food and influences the consumers buying or consumer's choice. According to [39], sensory properties of a food count the most of consumers' decision. Although in smallest amounts as ppm or ppb, aroma compounds are issued from complex biochemical reactions which include fermentation, lipid oxidation, enzymes, amino acids, degradation proteins and hydrocarbons, and Maillard reaction. Since then, numerous techniques have been developed for their extraction from different matrices and sensitive methods such as gas chromatography (GC), gas chromatography-mass spectrometry (GC-MS), and gas chromatographic-mass spectrometry-olfactometry (GC-MS-O) have been worked out for their quantification, identification and determination of key odorants, respectively [40]. Extant literature indicated that overall aroma profile of condiments obtained

from *P. biglobosa* seeds, is generated mainly during and cooking and fermentation under the metabolic activities of *Bacillus spp* [37,41,42]. Moreover, *Bacillus spp* have displayed great degradation activities of lipids and proteins [43,44,45]. Hence, in this review, an account of the reported production processes of *iru*, *sonru*, *afitin*, *soumbala*, *dawadawa*, *soumbara* and *netetu* and their respective aroma profile was investigated.

## 2. Production

Dried *P. biglobosa* seeds are used to produce different condiments. Seeds are washed and sorted in order to eliminate spoiled grains and foreigner particles. Processing of different seeds condiments such as *iru*, *sonru*, *soumbala*, *soumbara*, *dawadawa*, *afitin* and *netetu*, takes place through four common major operations. These include a first long cooking, dehulling, a second short cooking, and fermentation. Nonetheless, some intermediate operations like washing, cooling and addition of additives intervene during manufacturing of the condiments. The technique of production is ancestrally transferred and depend on cultural background, regions, and countries. This transformation activity is mainly performed by women for both house livelihoods and commercial purpose as it represents an important source of income, especially for rural populations. Generally, men are charged to harvest and collect the fruits while women transform the seeds into various condiments [17].

The first cooking is the first important operation that seeds undergo during condiments production and it consists of boiling seed in water for 6 to 24 hours. This operation allows not only a loss of astringency or bitterness of the seed but also the softening of the seminal integument [38]. For this cooking, the parameter time is very important and varies highly depending on the type of condiments and country or culture. Thus, during processing of fermented condiments of the Republic of Benin (*iru*, *afitin* and *sonru*), seeds are cooked for about 12 hours [26] while for *iru* and *dawadawa* of Nigeria, seeds undergo a cooking of 6-12 and 12 hours, respectively [46,47,48]. Likewise, to obtain *soumbala* of Burkina Faso, seeds are boiled for 24 hours [37] while for *netetu* of Senegal, seeds are cooked for 15-24 hours [38,49]. After this step, the cooked seeds are let to cool and wash in order to take off any remaining impurity and astringency. Seeds are then ready for dehulling.

Dehulling is the second major operation involving in the production of the fermented condiments and consists removing seed coats. Traditionally, seeds are marched by feet in a

large timbered mortar and further seed coats are taken off by pressing between palms of hands. In the Republic of Benin, fine sand or ash is added [26] with a goal to render the dehulling easier as this operation is tedious and needs more physical force. Afterward, seeds are minutely washed and sorted, and the cleaned cotyledons are sent to the second cooking.

The second cooking is a short 1 to 2 hours boiling of cotyledons in water. This operation is also important as it is associated with a kind of bleaching that helps to minimize secondary contamination which may result from the various manipulations during dehulling [38]. During the production of *iru* and *sonru*, additives *iku-iru* and *yanyanku* are appended towards the end of the second cooking [26]. *Yanyanku* is a ground dried seeds of *Hibiscus sabdariffa* while *iku-iru* is a mixture of ground dried seeds of *Hibiscus sabdariffa* L. and pepper as shown in Fig 2. *H. sabdariffa* also named Roselle, hibiscus, Jamaica sorrel or red sorrel (English) and karkadeh (Arabic) is a herbaceous plant largely planted in tropical and subtropical areas [50]. It appertains to *Malvaceae* family and is recognized as an annual, bushy plant with a height of up to 2.5 m, characterized by smooth, cylindrical red stems, reddish veins and long, green leaves [51]. *H. sabdariffa* seeds are revealed to contain potential nutrients as they displayed important values of proteins, fat, carbohydrate, fiber, minerals and, vitamins [52,53,54]. The introduction of additives from *H. sabdariffa* seeds may enhance not only the nutritional value and flavor of the condiments but also may improve their medicinal properties. Previous studies showed that additives impacted microbiologic characteristics of condiments [26]. In Nigeria, this second cooking can be extended from 45 minutes to 6 hours [46,47] while in the most of the countries it lasts 1 to 2 hours [26,37,38,49]. After the cooking, cotyledons are well drained.

The fermentation represents the last most important operation involved in the production of seeds condiments. Although above-cited operations prepare cotyledons to render the fermentation most successful, this step is the most sensitive amongst all operations. It gives to cotyledons all their nutritional capacities, microbiologic properties, and organoleptic characteristics. Cotyledons are spread in basket trays, calabash trays or containers, wrapped with heavy cloths, jutes backs and/ or polyethylene bags and then left for fermentation which lasts from 24 hours to days at room temperature. In the Republic of Benin, cotyledons are fermented for 24 hours to obtain *afitin*, whereas, for *iru* and *sonru*, they are subjected to 48 hours of fermentation [26]. As for *netetu* produced in Senegal, the fermentation takes 3 days [38,49] while in Burkina Faso, *soumbala* is obtained after 36 to 48 hours of fermentation [37,55]. Thus, time is a discriminative factor which characterizes the

condiments. Furthermore, long fermentation leads to more softening of cotyledons [26]. Temperature is another important factor which influences the fermentation. Generally, fermentation occurs at room temperature, about 30°C, as this temperature represents the average of the most condiments producers countries. [38] suggested 28-40°C and this was supported by the study of [48], in which 40°C was detected as the optimum temperature to yield more proteins. After the fermentation, the condiment is either dried, ground or dumped.

The Fermentation of *P. biglobosa* seeds is a spontaneous and lactic type carried out in alkaline condition [38,49]. Microbiologic studies revealed that *Bacillus* species are dominant microflora implying in the fermentation of *P. biglobosa* seeds [26,35, 38,43,49,56,57,58]. These microorganisms showed high metabolic activities during seeds fermentation with degradation of proteins, lipids and carbohydrates [43,44,45,57]. Furthermore, they showed antifungal activity in condiments by secreting mainly Iturin A and Surfactin [38]. The fermented condiments are safe for consumption, however, a few *Staphylococcus* germ was detected in some condiments. The presence of *Staphylococcus* in condiments is probably due to bad handling during manufacturing and commercialization [26,49,56]. Therefore, hygienic conditions have to be set up and respected during and after production. In addition, salting was suggested to reduce the microbial load of the fermented seeds condiments [59].

The raw African locust beans are too hard and almost unedible, however, they become soft, edible with strong appreciated flavor since they are processed to the fermented condiments. These are incorporated in traditional soups and stews as flavorful, protein and fat-rich foodstuffs. They are also eaten together cereal-based dumplings and porridges. Furthermore, the condiments are introduced in traditional vegetable sauces and can be reduced in powder. In addition, they replace animal and fish proteins in some dishes. The variations of the operations involving in the production affect absolutely physicochemical, biochemical and microbiological characteristics of the final products.

### **3. Aroma profile of the fermented condiments of *P. Biglobosa***

Quality of the fermented condiments of *P. biglobosa* is important for their commercial value and is consisted of color, flavor, and freshness. The flavor is defined as taste, which consists of non-volatile compounds, and aroma, which consists of volatile compounds. likewise, phenolic compounds are accountable for the color. Aroma compounds are important

not only for the distinction of the condiments but also for the evaluation of their quality. Nowadays, more than 160 volatile compounds namely pyrazines, ketones, aldehydes, alcohols, esters, alkanes, alkenes, benzene derivatives, pyridines, furan, volatile phenols, sulfur compounds, and terpenes have been reported in fermented African locust beans. Most of these compounds are generated during the manufacturing process mainly during fermenta-




Fig 1: Flow diagram of some African locust beans condiments




Fig 2: Flow diagram of yanyanku and Iku-iru [26]

Tion, and heating [37,41,60]. The main precursors of the volatile compounds in the fermented condiments are peptides, fatty acids and amino-acids [41]. Some variations have been quantitatively and qualitatively noticed in the overall aroma profile of *iru*, *sonru*, *afitin* and *dawadawa* (Tab. 1). This may probably due to the origin of the raw material, extraction technique, solvent as well as processing conditions notably heating and fermentation parameters.

Pyrazines represent the most important class detected in the investigated *P. biglobosa* condiments [37,42,60]. Also known as 1,4-diazine, pyrazines are six-membered aromatic heterocycles containing nitrogen atoms. They are naturally encountered in plants, animals and many foodstuffs as odorous and pharmacological agents. Pyrazines have been reported with roasted, toasted, nut-and peanut-like odor in roasted soybeans oils, yogurt, coffee, teas, roasted sorghum, fermented soybeans, roasted almond, wines and seeds of *B. Officinalis* [43,61,62,64,65,66]. They are increasingly used for medicinal purposes due to their anti-inflammatory, diuretics, antineoplastics, anti-infective, antidepressant, antiviral and neuroprotective properties [67]. The release of pyrazines is stimulated by heating. Furthermore, bacteria and fungi can produce and use pyrazines [68]. Although pyrazines are found in numerous products, they are known to be aroma compounds of heated products. A total of 19 aroma compound pyrazines (Tab. 1) have been quantified and identified *soumbala*, *afitin*, *sonru*, *iru*, and *daddawa*. These aroma compounds have been generated during heating and fermentation by metabolic reactions of the *Bacillus spp* [37,42,60]. The metabolic reactions of the *Bacillus spp* occurring during fermentation *P. biglobosa* cotyledons concerned degradation of lipids, proteins and poly and oligosaccharide [43,44,45,69]. Among all detected pyrazine, 2,5-dimethyl pyrazine is most the abundant aroma compounds figured out in the fermented condiments, followed by tetramethylpyrazine and trimethyl pyrazine. These aroma compounds have been reported in the similar fermented soybeans condiments [61]. 2,5-Dimethyl pyrazine (earthy, potato sweet) is one of the most important aroma compounds detected in palm sugars [70] and cooked *Digitaria exilis* Stapf (Lasekan et al., 2001), while tetramethylpyrazine (Nutty) and trimethyl pyrazine (nutty) have been detected as key odorants in the fermented Chinese Baijiu [71].

Aldehydes revealed to be the second most significant volatile class identified quantitatively in the *P. biglobosa* condiments [26,42,60,72]. Aldehydes are synthesized from the oxidative reaction of fatty acid mainly  $\alpha$ -linolenic acid and linoleic [73]. They can be also liberated from Strecker degradation, non-enzymatic and heat-induced of amino acids [74]. Aldehydes displayed green and citrus notes and have been reported in various food including hazelnuts, wines, beer, saffron, virgin olive oil, olive oil, fermented soybeans and canola oil [61,75,76,77,78]. The presence of aldehydes in the fermented condiments of *P. biglobosa* is associated with Maillard reaction occurring during the fermentation [26,42]. A total of 33 individual aldehydes have been found in the data regarding volatile compounds of *P. biglobosa* (Tab. 1). Among these, 3-methyl butanal, benzene acetaldehyde, benzaldehyde, and

2,4-decadienal (E, E) are the most significant as they displayed higher concentrations in the condiments. Previously, benzaldehyde (sweet, almond) and benzene acetaldehyde (sweet, honey-like) have detected to contribute to the active odorants of kinds of honey [79], whereas, 2,4-decadienal (E, E) (fryer oil) and 3-methyl butanal (fruity, malty) have been found in liquid whey and black sapote, respectively, as aroma active compounds [80,81].

Ketones are products of complex biosynthesis reactions which imply the degradation of fatty and amino acids [82,83] and Maillard reaction [84]. In the case of fermented seeds like fermented *P. biglobosa* seeds or fermented soybeans, it believed that ketones are originated from the degradation of lipids, amino acids and saccharides [26,42,61,85]. They contribute highly to the aroma profile of the fermented condiments of *P. biglobosa* as 21 individual aromatic ketones have been recorded in all the studied spices (Tab. 1). The distribution of ketones in the investigated products varied and depended on the type of the condiments, type of fermentation (spontaneous or controlled) and type of starter culture. For instance, the most important ketones discovered the in traditional and controlled fermentation soumbala are 2-nonadecanone, acetophenone, 2-heptanone and 2-heptadecanone [37], while, the most abundant aromatic ketones in the controlled fermentation *afitin*, *iru* and *sonru* are 2-decanone, 1-butanone, 3-hydroxy-2-butanone and 2-pentadecanone [60]. Most these volatile compounds have been reported in the fermented soybean condiments [61]. Acetophenone has been reported in the essential oil of *Camellia sinensis* L. [86] while 2-Nonadecanone has been indicated in both essential oils of *Salix aegyptiaca* L. [87] and *Camellia sinensis* L. [86]. On one hand, *Salix aegyptiaca* L. is used to treat a headache and also showed cardiovascular protective, sedative and analgesic effects, and on another hand, *Camellia sinensis* L. in various forms is used as anti-inflammatory, antiviral, immunoprotective and anti-histaminic [88]. Furthermore, 2-decanone, 3-hydroxy-2-butanone and 2-nonadecanone have been detected to contribute the scent of the fermented baobab (*Adansonia digitata* L.) seeds [89].

A total of 27 alcohols have been reported in the condiments. Generally, lipoxygenase in the presence of oxygen, acted on unsaturated fatty acids to produce volatile alcohols [90]. Additionally, they could be released during fermentation of *P. biglobosa* cotyledons under microorganisms action [26,45]. In spite of the similarities between traditional and starter culture condiments, the highest concentration of volatile alcohols have been recorded in the controlled fermentation products. 3,5-Dimethyl phenylmethanol, 2-phenyl butanol, 3-methyl-1-butanol, and 2-hexanol represented the major alcohols in the fermented condiments of the Republic of Benin [42,60] while 3-methyl-1-butanol, 2-pentanol, 1-octen-3-ol and 3,7,11-

trimethyl-2,6,10-dodecatrien-1-ol were found in *soumbala* [37]. Previously, 1-octan-3-ol has been found in the fermented soybeans [61]. Likewise, 2-hexanol and 1-octen-3-ol have been detected as aroma contributors of essential oils of *Camellia oleifera* Abel. seeds [91]. It is well-known that seeds oils of *Camellia oleifera* seeds are comparable to olive oils with health benefits including antioxidant properties, reduction of cholesterol and blood pressure [92,93,94]. Moreover, 3-methyl-1-butanol (cheese-like, ripe onion) and 1-octen-3-ol (earthy, dusty, spicy, mushroom) have figured out as aroma active compounds of *Tuber melanosporum* and *Tuber aestivum* [95] in one hand and of essence and puree of kiwi [96] in another hand.

Table 1: Volatile compounds of the fermented seeds of *P. biglobosa*

No	Compounds	Afitin	Sonru	Iru	Dawadawa/ Daddawa	Soumbal a
<b>Pyrazines</b>						
1	2,5-Dimethyl pyrazine	✓	✓	✓	✓	✓
2	Methyl pyrazine	✓	✓	✓		✓
3	2,6-Dimethyl pyrazine	✓	✓	✓	✓	✓
4	Ethyl pyrazine					✓
5	2,3-Dimethyl pyrazine	✓	✓	✓		✓
6	2-Ethyl-6-methyl pyrazine	✓	✓	✓	✓	✓
7	2-Ethyl-5-methyl pyrazine				✓	✓
8	Trimethyl pyrazine	✓	✓	✓	✓	✓
9	2,6-Diethyl pyrazine					✓
10	3-Ethyl-2,5-dimethyl pyrazine				✓	✓
11	2-Ethyl-3,5-dimethyl pyrazine	✓	✓	✓		✓
12	Tetramethyl pyrazine	✓	✓	✓	✓	✓
13	2,5-Dimethyl-3-(2-methyl) pyrazine					✓
14	2,3-Dimethyl-5- propylpyrazine	✓	✓	✓		✓

15	2,3-Dimethyl-5-propenyl pyrazine (E,E)	✓	✓	✓		✓
16	2,5-Dimethyl-3-(3-methyl) pyrazine					✓
17	2-Methyl pyrazine	✓	✓	✓		
18	3-Ethyl-3,5-dimethyl pyrazine					
19	2,3,5-Trimethyl-6-ethyl pyrazine	✓	✓	✓		
<b>Aldehydes</b>						
20	3-Methyl butanal	✓	✓	✓		✓
21	Pentanal	✓	✓	✓	✓	✓
22	2-Butenal	✓	✓	✓		✓
23	Hexanal				✓	✓
24	Heptanal	✓	✓	✓	✓	✓
25	3-Methyl-2-butenal	✓	✓	✓		✓
26	Octanal	✓	✓	✓		✓
27	Nonanal				✓	✓
28	2-Octenal (E)	✓	✓	✓		✓
29	3-Methylthio propanal					✓
30	3-Furancarboxaldehyde					✓
31	2,4-Heptadienal (E,E)	✓	✓	✓		✓
32	Benzaldehyde	✓	✓	✓	✓	✓
33	Nonenal	✓	✓	✓		
34	2-Nonenal	✓	✓	✓	✓	
35	2-Nonenal (E)	✓	✓	✓		✓
36	Benzeneacetaldehyde	✓	✓	✓	✓	✓
37	2,4-Decadienal				✓	✓
38	2,4-Decadienal (E,E)	✓	✓	✓		✓
39	2-Phenylpropenal					✓
40	2-Phenylbutanal	✓	✓	✓		✓
41	5-Methyl-2-phenyl-2- hexanal	✓	✓	✓		✓

42	3-Phenyl-2-propenal	✓	✓	✓	
43	3-Phenyl butanal				
44	1-Decanal	✓	✓	✓	✓
45	1-Tetradecanal	✓	✓	✓	
46	2-Butyl-octenal				✓
47	Dodecanal				✓
48	2,4-Nonadienal				✓
49	Hendecanal				✓
50	2-Nonanal				✓
51	2,4-Undecadienal				✓
52	2,4,6-Dodecatrienal				✓
	<b>Ketones</b>				
53	2-Methyl-3-pentanone	✓	✓	✓	✓
54	5-Methyl-3-hexanone	✓	✓	✓	✓
55	2-Heptanone	✓	✓	✓	✓
56	2-Octanone	✓	✓	✓	✓
57	2-Decanone	✓	✓	✓	✓
58	2-Undecanone				✓
59	Acetophone				✓
60	Acetophenone				✓
61	2-Tetradecanone	✓	✓	✓	✓
62	2-Pentadecanone	✓	✓	✓	✓
63	2-Heptadecanone	✓	✓	✓	✓
64	2-Nonadecanone	✓	✓	✓	✓
65	2-Pentacosanone	✓	✓	✓	✓
66	3-Hydroxy-2-butanone	✓	✓	✓	
67	1-Butanone	✓	✓	✓	
68	6-Methyl-2-heptanone	✓	✓	✓	
69	3-Ethyl hexanone				✓
70	3-Ethyl heptanone				✓
71	3-Ethyl octanone				✓
72	Ethyl nonanone				✓
73	2,5-Hexadione				✓

<b>Alcohols</b>					
74	2-Methyl-1-propanol	✓	✓	✓	✓
75	3-Pentanol	✓	✓	✓	✓
76	2-Pentanol				✓
77	3-Methyl-1-butanol	✓	✓	✓	✓
78	1-Pentanol	✓	✓	✓	✓
79	2-Methyl-2-buten-1-ol				✓
80	1-Octen-3-ol				✓
81	2-Ethyl-1-hexanol	✓	✓	✓	✓
82	3,7-Dimethyl-2,6-octadien-1-ol				✓
83	3,5-Dimethyl benzenemethanol	✓	✓	✓	✓
84	Benzyl alcohol	✓	✓	✓	✓
85	Octadecanol	✓	✓	✓	✓
86	3,7,11-Triethyl-2,6,10-dodecatrien-1-ol				✓
87	2-Ethyl cyclobutanol				✓
88	2-Hexanol	✓	✓	✓	
89	1-Hexadecanol	✓	✓	✓	
90	Decanol	✓	✓	✓	
91	2-Phenyl ethanol	✓	✓	✓	
92	2-Butanol	✓	✓	✓	
93	1-Hexanol	✓	✓	✓	
94	Ethanol				✓
95	Propanol				✓
96	1,2-Ethandiol				✓
97	Octanol				✓
98	Dodecanol				✓
99	Undecanol				✓
<b>Esters</b>					
100	2-Methylpropyl butanoate	✓	✓	✓	
101	Methyl benzeneacetate				✓

102	Ethyl benzeneacetate				✓
103	Methyl pentadecanoate				✓
104	Ethyl 4-ethoxy-benzoate				✓
105	Methyl hexadecanoate	✓	✓	✓	✓
106	Ethyl hexadecanoate				✓
107	Methyl octadecanoate				✓
108	3-Hexenyl acetate, (E)	✓		✓	
109	9-Methyl oleate, (Z)	✓	✓	✓	
110	Ethyl oleate		✓		
111	9,12-Methyl octadecadienoate, (Z, Z)	✓	✓	✓	
112	Ethyl linoleate	✓	✓	✓	✓
113	2-Methyl-ethyl-butanoate	✓	✓	✓	
114	Ethyl acetate				✓
115	Ethyl hexanoate				✓
116	Octadecyl acetate				✓
117	Octyl formate				✓
	<b>Phenols</b>				
118	2-Methoxy phenol				✓
119	2-Methoxy-4-methyl phenol				✓
120	Phenol	✓	✓	✓	✓
121	4-Ethyl-2-methoxy phenol				✓
122	2,4-Diisopropyl phenol				✓
	<b>Acids</b>				
123	9,12-Octadecadienoic acid (Z, Z)	✓	✓	✓	✓
124	Hexadecanoic acid				✓
125	9-Hexadecanoic acid (Z)	✓	✓	✓	✓
126	2,3-Dimethoxy butanedioic acid	✓			✓
127	2-Methyl butanoic acid	✓	✓	✓	
128	1,2-Benzendicarboxylic	✓	✓	✓	✓

---

	aid				
129	Tetradecanoic acid	✓	✓	✓	
130	9,12-Methyl octadecadienoic acid (Z, Z)	✓	✓	✓	
131	2-Methyl-2-pentanoic acid				✓
132	Cyclopropanenonanoic acid				✓
133	Decanoic acid			✓	
	<b>Alkanes</b>				
134	Tridecane				✓
135	1-Heptyl-2-methyl cyclopropane				✓
136	Cyclododecane			✓	✓
137	Cyclotetradecane			✓	
138	Cyclohexadecane			✓	
139	2-Methyl undecane				✓
140	Docosane	✓	✓	✓	✓
141	Tetracosane	✓	✓		✓
142	Octacosane				✓
143	Heptacosane	✓	✓	✓	
144	Eicosane	✓	✓	✓	
145	9-Octyldocosane	✓	✓	✓	
146	Heneicosane	✓	✓	✓	
147	Hexacosane			✓	
	<b>Alkenes</b>				
148	3-Ethyl-2-methyl-1,3-hexadiene				✓
149	1,5-Cyclodecadiene (E, Z)				✓
150	1-Octadecene				✓
151	1-Dodecene	✓	✓		
152	1-Decene	✓		✓	
153	1-Heptadecene		✓	✓	

---

154	Tetradecene					✓
	<b>Benzene derivatives</b>					
155	1-Ethyl-2,4-dimethyl benzene					✓
156	1-Ethyl-3,5-dimethyl benzene					✓
157	1,2,3,5-Tetramethyl benzene					✓
158	1,2-Dichloro benzene					✓
159	Benzeneacetoitrile					✓
160	1,3-Dimethyl benzene	✓	✓	✓		
161	1,2-Dimethyl benzene					✓
162	Toluene					✓
163	Chlorobenzene	✓	✓	✓		
	<b>Pyridines</b>					
164	Pyridine					✓
165	2-Propyl pyridine					✓
166	2-Diphenylmethyl pyridine					✓
167	2,6-Diphenyl pyridine					✓
	<b>Furans</b>					
168	2-Ethyl furan	✓	✓	✓		✓
169	2-Pentyl furan	✓	✓	✓		
	<b>Sulfur compounds</b>					
170	Dimethyl disulfide	✓	✓	✓	✓	✓
171	Dimethyl trisulfide	✓	✓	✓		✓
172	1-(Methylthio) propane					✓
173	1,2,4-Trithiolane	✓		✓		✓
	<b>Terpenes</b>					
174	Limonene					✓
	<b>Others</b>					
175	Indole					✓
176	Trimethyl oxazole					✓

---

<b>177</b>	3,4,5-Trimethyl pyrazole	✓
<b>178</b>	5-Methyl isothiazole	✓
<b>179</b>	4,5-Dihydro-2-methyl thiazole	✓

---

✓ : Presence; Source: [37,42,60,72]

Esters, Phenols, benzenes derivatives, alkanes, alkenes, acids, sulfur compounds, pyridines, furans, and others are contributors to the overall sensorial characteristic of the fermented seeds of *P. biglobosa*. Similarly, alkanes, alkenes, benzenes, sulfur compounds, and furans have detected in a small amount in the fermented seeds of baobab and soybeans [61,89,97]. Volatile compounds of the fermented seeds of *P. biglobosa* increased quantitatively and qualitatively during fermentation. This observation is linked with the degradation of primary and secondary metabolites by microorganisms notably *Bacillus spp* [26,37,42,43,44,45]. A similar increase of volatile compounds after fermentation has been registered in the fermented seeds of baobab and soybeans [61,89]. Duration of fermentation affects aroma compounds of the fermented seeds as a long fermentation provided a higher amount of aroma compounds. It has been noticed that the product obtained with 48h of fermentation are richer in volatile compounds than the product obtained at 24h of fermentation [42,60]. Alike, results have been noted in Japanese fermented soy (*natto*) [97]. Likewise, type of fermentation as it is spontaneous or controlled, influence scent of the condiment. For instance, the condiments resulted from the controlled fermentation (use of *Bacillus spp* as starter culture) contained more aroma compounds than the condiments obtained with spontaneous fermentation [37,60]. Other factors which influence sensorial properties of the fermented seeds are processing conditions mainly boiling or roasting, additives and post-fermentation operations like drying. The extant literature showed that both boiling and roasting before fermentation, affected the global chemical and biochemical constituents of the fermented condiments of African locust beans [46] and this consequently affects aroma profiles of the finished products. In the same way, boiling and roasting revealed to affect the overall aroma of the fermented soybean condiments [98]. Moreover, the use of additives such as *Iku-iru* and *Yanyanku* increased microorganisms counts [26] together with aroma compounds [42]. Additionally, drying after fermentation has been revealed to decrease volatile compounds of the fermented seeds condiments [89,97].

#### 4. Conclusion

The present study described the production process and volatile profiles of different fermented condiments of African locust bean seeds encompassing *afitin*, *soumbala*, *sonru*, *iru*, *dawadawa* and *netetu*. Four mainly operations including successively a first long cooking, dehulling, a second short cooking and fermentation were applied during the production of all studied condiments. However, some production parameters such as time, temperature and additives rendered each condiment unique. About 130 volatile compounds have identified and quantified in these products. Pyrazines, aldehydes, ketones, and alcohols have been detected to contribute mainly to the overall aroma profile of the products. Further experiments should be performed for characterizing aroma-active compounds of these fermented condiments.

#### References

- [1] Wilburn JR, Ryan EP. Fermented foods in health promotion and disease prevention: An overview A2-Frias, Juana. In Fermented foods in health and disease prevention. eds. Martinez-Villaluenga C, Penas E. Boston: Academic Press; 2017. 3-19 pp.
- [2] Nout MJR. Food technologies: Fermentation A2-Motarjemi, Yasmine Encyclopedia of food safety. Waltham: Academic Press; 2014. 168-77 pp.
- [3] Kim JH, Kim Y, Kim YJ, Park Y. Conjugated Linoleic acid: Potential health benefits as a functional food ingredient. Ann Rev Food Sci Technol 2016; 7:221-44.
- [4] Dincer E, Kivanc M. Characterization of lactic acid bacteria from Turkish Pastirma. Ann Microbiol 2012; 62:1155-63.
- [5] Desobry S, Debeaufort F. Edible coatings and films to improve food quality: Encapsulation of flavors, nutraceuticals, and antibacterials. Eds. Baldwin EA, Hagenmaier R, Bai J. Boca Raton, London, New York: CRC press, Taylor & Francis Group; 2012. 362 p.
- [6] Olasupo, N.A., Odunfa, S.A., Obayori, O.S., 2010. Fermented Foods and Beverages of the World: Ethnic African fermented foods. Eds. Tamang JP, Kailasapathy K. London: press; 2010. 323-52 pp.

- [7] Şanlıer, N., Gökçen, B.B., Sezgin, A.C. (2017). Health benefits of fermented foods. *Crit Rev Food Sci Nutr* 2017; 0:1-22.
- [8] Mathara JM, Schillinger U, Kutima PM, Mbugua SK, Holzappel WH. Isolation, identification and characterisation of dominant microorganisms of *Kule naoto*: The *Maasai* traditional fermented milk in Kenya. *Inte J Food Microbiol* 2004; 64:269-78.
- [9] Mathara JM, Schillinger U, Guigas C, Franz C, Kutima PM, Mbugua SK, Shin H-K, Holzappel WH. Functional characteristics of *Lactobacillus spp* from traditional *Maasai* fermented milk products in Kenya. *Int J Food Microbiol* 2008; 126:57-64.
- [10] Tamang J, Samuel D. Fermented Foods and Beverages of the World: Dietary cultures and antiquity of fermented foods and beverages. Eds. Tamang JP, Kailasapathy K. London: CRC press; 2010. 1-40 pp.
- [11] Amusa O, Adesoye A, Ogunkanmi A, Omoche O, Olowe O, Akinyosoye S, Omodele T. Genetic diversity of *Parkia biglobosa* from different agroecological zones of Nigeria using RAPD markers. *Int J Biodiv* 2014; 1-6.
- [12] Koura K, Ganglo JC, Assogbadjo AE, Agbangla C. Ethnic differences in use values and use patterns of *Parkia biglobosa* in Northern Benin. *J Ethnobiol Ethnomed* 2011; 7:1-12.
- [13] Kwon-Ndung, EHC, Chup CD, Binbol NA, Ismaila A. Ecological implications of climate change on the genetic diversity and distribution of African locust bean *Parkia biglobosa* in Central Nigeria. *IOP Conference Series: Earth Environ Sci* 2009; 6: Article ID372026.
- [14] Irvine FR. Wood plants of Ghana. London: Oxford University Press; 1961. 104 p.
- [15] Hagos TH. A revision of the genus *Parkia* R. Br. (Mim) in Africa. *Acta Botanica Neerlandica* 1962; 11:231-265.
- [16] Tringali C, Spatafora C, Longo OD. Bioactive constituents of the bark of *Parkia biglobosa*. *Fitoterapia* 2000; 71:118-25.
- [17] Nyadanu D, Amoah RA, Obeng B, Kwarteng AO, Akromah R, Aboagye LM, Adu-Dapaah H. Ethnobotany and analysis of food components of African locust bean (*Parkia biglobosa* (Jacq.) Benth.) in the transitional zone of Ghana: Implications for domestication, conservation and breeding of improved varieties. *Genet Resour Crop Evol* 2016; 64: 1231-40.

- [18] Ajaiyeoba EO. Phytochemical and antibacterial properties of *Parkia biglobosa* and *Parkia bicolor* leaf extracts. *Afr J Biomed Res* 2002; 5:125-29.
- [19] El-Mahmood AM, Ameh JM. *In vitro* antibacterial activity of *Parkia biglobosa* (Jacq.) root bark extract against some microorganisms associated with urinary tract infections. *Afr J Biotechnol* 2007; 6:1272-75.
- [20] Udobi CE, Onaolapo JA. Phytochemical analysis and antibacterial evaluation of the leaf stem bark and root of the African locust bean (*Parkia biglobosa*). *J Med Plant Res* Vol 2009; 3:338-44.
- [21] Bukar A, Uba A, Oyeyi TI. Phytochemical Analysis and Antimicrobial Activity of *Parkia Biglobosa* (Jacq.) Benth. Extracts against some food-borne microorganisms. *Adv Environ Biol* 2010; 4:74-79.
- [22] Ibekwe VI, Ezeji EU. Antimicrobial activity of leaf and root of *Parkia biglobosa* (African locust bean). *The Afr J Pant Sci Biotechnol* 2011; 5:70-72.
- [23] Hounhouigan DJ, Jansen JMM, Nout, MJR, Nago MC, Rombouts FM. Traditional African foods quality and nutrition: Production of quality maize-based fermented dough in Benin urban area. Eds. Westby A, Reilly PJA. Dar es Salaam, Tanzania: Proceedings of Regional Workshop held by the International Foundation for Science; 1991. 9-18 pp.
- [24] Kayodé APP, Hounhouigan JD, Nout MJR. Impact of brewing process operations on phytate, phenolic compounds and *in vitro* solubility of iron and zinc in opaque sorghum beer. *LWT-Food Sci Technol* 2007; 40:834-41.
- [25] Whitby, P. Foods of Ghana. *Food Res Inst Bulletin* 1968; 1:1-31.
- [26] Azokpota P, Hounhouigan DJ, Nago MC. Microbiological and chemical changes during the fermentation of African locust bean (*Parkia biglobosa*) to produce *afitin*, *iru* and *sonru*, three traditional condiments produced in Benin. *Int J Food Microbiol* 2006; 107:304-309.
- [27] Ali AA and Mustafa MM. Use of starter cultures of lactic acid bacteria and yeasts in the preparation of *Kisra*, a Sudanese fermented food. *Pak J Nutr* 2009; 8:349-53.
- [28] Antai SP, Ibrahim MH. Microorganisms associated with African locust bean (*Parkia filicoidea* Welw) fermentation for *dawadawa* production. *J Appl Bacteriol* 1986; 61:145-48.

- [29] Tchekessi CKC, Bokossa A, Agbangla C, Azokpota P, Daube G, Scippo M-L, Korsak N, Angelov A, Bokossa YI. (2014). Production and microbiological evaluation of three types of "Dèguè", a local fermented drink made from milk in Benin. *Int J Multidiscipl Cur Res* 2014; 2:714-20.
- [30] Diawara B, Sawadogo L, Kabore IZ. Contribution a l'étude des procedes traditionnels de fabrication du 'sombala' au Burkina Faso. Aspects biochimiques, microbiologiques et technologiques. *Sci Techn* 1992; 20:5-14.
- [31] Mensah, P. Fermentation - the key to food safety assurance in Africa? *Food Control* 1997; 8:271-78.
- [32] Obun CO. Performance and some heamatological response of finisher broilers fed graded levels of fermented locust bean (*Parkia biglobosa*) seeds meal. *Asi J Agri Res* 2007; 1:125-30.
- [33] Maidala A, Doma UD, Egbo LM. Nutritional evaluation of local processing methods of African locust bean seeds (*Parkia biglobosa*) on proximate composition and growth performance of broiler chickens. *J Environ Technol Sustain Agri* 2013; 2 (1):1-9.
- [34] Audu BS, Adamu KM, Binga SA. Effect of substituting fishmeal diets with varying quantities of ensiled parboiled beniseeds (*Sesamum indicum*) and raw African locust bean (*Parkia biglobosa*) on the growth responses and food utilization of the Nile tilapia (*Oreochromis niloticus*). *Int J Zool Res* 2008; 4:42-7.
- [35] Odunfa SA, Oyewole OB. Identification of *Bacillus* species from 'iru', a fermented African locust bean product. *J. Basic Microbiol* 1986; 26(2): 101-08.
- [36] Adeyeye EI, Ipinmorotin KO, Oguntokun MO. Chemical composition and functional properties of African locust bean (*Parkia biglobosa*) seeds. *Pak J Sci Indus Res* 2002; 45:29-33.
- [37] Ouoba LII, Diawara B, Annan NT, Poll L, Jakobsen M. Volatile compounds of *Sombala*, a fermented African locust bean (*Parkia biglobosa*) food condiment. *J Appl Microbiol* 2005; 99:1413-21.
- [38] N'dir B, Hbid CL, Cornelius C, Roblain D, Jacques P, Vanhentenryck F, Diop M, Thonard P. Propriétés antifongiques de la microflora sporulée du *netétu*. *Cahiers Agric* 1994; 3:23-30.

- [39] Furst T, Connors M, Bisogni CA, Sobal J, Winter Falk L. Food choice: A conceptual model of the process. *Appetite* 1996; 26:247-66.
- [40] Reineccius G. (2006). *Flavor chemistry technology*. 2 ed. Taylor & Francis Group; 2006. 53p.
- [41] Beaumont M. Flavouring composition prepared by fermentation with *Bacillus spp.* *Int J Food Microbiol* 2002; 75:189-96.
- [42] Azokpota P, Hounhouigan DJ, Annan NT, Nago MC, Jakobsen M. Diversity of volatile compounds of *afitin*, *iru* and *sonru*, three fermented food condiments from Benin. *World J Microbiol Biotechnol* 2008; 24:879-85.
- [43] Ouoba LII, Rechinger KB, Barkholt V, Diawara B, Traore AS, Jakobsen M. Degradation of proteins during the fermentation of African locust bean (*Parkia biglobosa*) by strains of *Bacillus subtilis* and *Bacillus pumilus* for production of *Soumbala*. *J Appl Microbiol* 2003; 94:396-402.
- [44] Ouoba LII, Cantor MD, Diawara B, Traoré AS, Jakobsen M. Degradation of African locust bean oil by *Bacillus subtilis* and *Bacillus pumilus* isolated from *Soumbala*, a fermented African locust bean condiment. *J Appl Microbiol* 2003; 95:868-73.
- [45] Azokpota P, Hounhouigan DJ, Nago MC, Jakobsen M. Esterase and protease activities of *Bacillus spp* from *afitin*, *iru* and *sonru*, three African locust bean (*Parkia biglobosa*) condiments from Benin. *Afr J Biotechnol* 2006; 5:265-72.
- [46] Omafuvbe BO, Falade OS, Osuntogun BA, Adewusi SRA. Chemical and biochemical changes in African locust bean (*Parkia biglobosa*) and Melon (*Citrullus vulgaris*) Seeds during fermentation to condiments. *Pak J Nutr* 2004; 3(3):140-45.
- [47] Ajala CA, Etejere EO, Olatunji TL. Determination of protein content in the two types of *Parkia biglobosa* (fermented and unfermented) seeds and seedlings. *Alb J Agric Sci* 2013; 12(4):711-13.
- [48] Ojewumi ME, Omoleye AJ, Ajayi AA. (2016) *Optimum Fermentation Temperature for the Protein Yield of Parkia biglobosa Seeds (Iyere)*. In: 3rd International Conference On African Development Issues (CU-ICADI), May 9- May 11.

- [49] N'dir B, Gningue RD, Keita NDG, Souane M, Laurent L, Cornelius C, Thonard P. Caracteristiques microbiologiques et organoleptiques du netetu du commerce. Cahiers Agric 1997; 6:299-304.
- [50] Cisse M, Dornier M, Sakho M, Ndiaye A, Reynes M, Sock O. Le bissap (*Hibiscus sabdariffa* L.): Composition et principales utilisations. Fruits 2009; 64:179-93.
- [51] Mohamed R, Fernandez J, Pineda M, Aguilar M. Roselle (*Hibiscus sabdariffa*) seed oil is a rich source of  $\gamma$ -tocopherol. J Food Sci 2007; 72(3):207-11.
- [52] Samy M.S. Chemical and nutritional studies on roselle seeds (*Hibiscus sabdariffa* L.). Z Ernahrungswiss 1980; 19:47-9.
- [53] Rao P U. Nutrient composition and biological evaluation of mesta (*Hibiscus sabdariffa*) seeds. Plant Food Human Nutr 1996; 49:27-34.
- [54] Hainida E, Ismail A, Hashim N, Mohd.-Esa N, Zakiah A. Effects of defatted dried roselle (*Hibiscus sabdariffa* L.) seed powder on lipid profiles of hypercholesterolemia rats. J Sci Food Agri 2008; 88:1043-50.
- [55] Diawara B, Sawadogo L, Kabore IZ. Contribution a l'etude des procedes traditionnels de fabrication du 'sombala' au Burkina Faso. Aspects biochimiques, microbiologiques et technologiques. Sci Techn 1992; 20:5-14.
- [56] Jideani IAO, Okeke CR. Comparative study of microorganisms and sensory attributes of condiments from the fermentation different seeds. Plant Food Human Nutr 1991; 41:27-34.
- [57] Savadogo A, Ilboudo AJ, Gnankiné O, Traoré AS. Numeration and Identification of thermotolerant endospore-forming *Bacillus* from two fermented condiments *Bikalga* and *Sombala*. Adv Environ Biol 2011; 5(9):2960-66.
- [58] Fatoumata C, Soronikpoho S, Souleymane T, Kouakou B, Marcellin DK. Caractéristiques biochimiques et microbiologiques de moutardes africaines produites à base de graines fermentées de *Parkia biglobosa* et de *Glycine max*, vendues en Côte d'Ivoire. Int J Biol Chem Sci 2016; 10(2):506-18.
- [59] Ademola IT, Baiyewu RA, Adekunle EA, Awe AB, Adewumi OJ, Ayodele OO, Oluwatoke FJ. Microbial load of processed *Parkia biglobosa* seeds: Towards enhanced shelf life. Afr J Agric Res 2013; 8(1):102-05.

- [60] Azokpota P, Hounhouigan DJ, Annan NT, Odjo T, Nago, M.C, Jakobsen M. Volatile compounds profile and sensory evaluation of Beninese condiments produced by inocula of *Bacillus subtilis*. J Sci Food Agric 2010; 90: 438-44.
- [61] Owens JD, Allagheny N, Kippinga G, Amesa JM. Formation of Volatile Compounds During *Bacillus subtilis* Fermentation of Soya Beans. J Sci Food Agric 1997; 74:132-40.
- [62] Cheng H. Volatile Flavor Compounds in Yogurt: A Review. Crit Rev Food Sci Nutr 2010; 50:938-50.
- [63] Lee J, Chambers DH, Chambers IVE, Adhikari K, Yoon Y. Volatile aroma compounds in various brewed green teas. Molecule 2013; 18:10024-041.
- [64] Amanpour A, Selli S. (2016). Differentiation of volatile profiles and odor activity values of turkish coffee and french press coffee. J Food Process Preserv 2016; 40:1116-24.
- [65] Sheibani E, Duncan SE, Kuhn DD, Dietrich AM, Newkirk JJ, O'Keefe SF. Changes in flavor volatile composition of oolong tea after panning during tea processing. Food Sci Nutr 2015; 4(3):456-68.
- [66] Wu S, Xu T, Huang, D. Chemical compositions of the volatile extracts from seeds of *Dendranthema nankingense* and *Borago officinalis*. J Food Drug Analysis 2015; 23:253-59.
- [67] Dolezal M, Zitko J. Pyrazine derivatives: a patent review. Expert Opin Ther Patent 2015; 25(1):33-47.
- [68] Rajini RS, Aparna P, Sasikala C, Ramana CV. Microbial metabolism of pyrazines. Crit Rev Microbiol 2011; 37:99-112.
- [69] Ouoba LII, Diawara B, Amoa-Awua WK, Traoré AS Lange Moller P. Genotyping of starter cultures of *Bacillus subtilis* and *Bacillus pumilus* for fermentation of African locust bean (*Parkia biglobosa*) to produce Soumbala. Int J Food Microbiol 2004; 90: 197-205.
- [70] Wan Aida WM, Ho CW, Maskat MY, Osman H. Relating descriptive sensory analysis to gas chromatography/mass spectrometry of palm sugars using partial least squares regression. ASEAN Food J 2008; 15(1):35-45.
- [71] Zhao D, Shi D, Sun J, Li A, Sun B, Zhao M, Chen F, Sun X, Li H, Huang M, Zheng F. Characterization of key aroma compounds in Gujingong Chinese Baijiu by gas

chromatography–olfactometry, quantitative measurements, and sensory evaluation. *Food Res Int* 2018; 105:616-27.

[72] Onyenekwel PC, Odeh C, Nweze CC. Volatile constituents of ogiri, soybean daddawa and locust bean daddawa three fermented Nigerian food flavour enhancers. *Electron J Environ Agri Food Chem* 2012; 11(1):115-21.

[73] Hatanaka A. The biogenesis of green odour by green leaves. *Phytochem* 1993; 34:1201-18.

[74] Strecker A. On a peculiar oxidation by alloxan (in German). *Justus Liebigs Ann Chem* 1862; 123:363.

[75] Kinderlerer JL, Johnson S. Rancidity in Hazelnuts due to Volatile Aliphatic Aldehydes. *J Sci Food Agric* 1992; 58:89-93.

[76] Fullana A, Carbonell-Barrachina AA, Sidhu S. Comparison of volatile aldehydes present in the cooking fumes of extra virgin olive, olive, and canola oils. *J Agric Food Chem* 2004; 52:5207-14.

[77] Culleré L, San-Juan, F., Cacho, J. Characterisation of aroma active compounds of Spanish saffron by gas chromatography–olfactometry: Quantitative evaluation of the most relevant aromatic compounds. *Food Chemistry* 2011; 127:1866-71.

[78] Amanpour A, Sonmezdag AS, Kelebek H, Selli S. GC-MS-olfactometric characterization of the most aroma-active components in a representative aromatic extract from Iranian saffron (*Crocus sativus* L.). *Food Chem* 2015; 182:251-56.

[79] Castro-Varquez L, Diaz-Maroto MC, Gonzalez-Vinas MA, Pérez-Coello MS. Differentiation of monofloral citrus, rosemary, eucalyptus, lavender, thyme and heather honeys based on volatile composition and sensory descriptive analysis. *Food Chem* 2009; 112:1022-30.

[80] Karagül-Yüceer Y, Drake MA, Cadwallader KR. Aroma-active components of liquid cheddar whey. *J Food Sci* 2003; 68(4):1215-19.

[81] Pino JA, Ortiz-Vazquez E, Sauri-Duch E, L. Cuevas-Glory E. Characterization of aroma-active compounds in black sapote (*Diospyros digyna* JACQ.). *Acta Alimentaria* 2014; 43(4):547-52.

- [82] Josephson DB, Lindsay RC. Enzymic generation of volatile aroma compounds from fresh fish. In Biogenesis of Aroma. Eds. Parliament TH, Croteau R. Washington, DC: American Chemical Society; 1986. 201-219 pp.
- [83] Chung HY, Cadwallader KR. Aroma extract dilution analysis of blue crab claw meat volatiles. *J Agri Food Chem* 1994; 42:2867-70.
- [84] Chung HY, Yung IKS, Ma WCJ, Kim JS. (2002). Analysis of volatile components in frozen and dried scallops (*Patinopecten yessoensis*) by gas chromatography/mass spectrometry. *Food Res Int* 2002; 35:43-53.
- [85] Janssens L, de Pooter HL, Vandamme EJ, Schamp NM. Production of flavours by microorganisms. *Process Biochem* 1992; 27:195-215.
- [86] Chen Y-J, Kuo P-C, Yang M-L, Li F-Y, Tzen JTC. Effects of baking and aging on the changes of phenolic and volatile compounds in the preparation of old Tieguanyin oolong teas. *Food Res Int* 2013; 53:732-43.
- [87] Shampur T, Sheikhshoae I, Afzali D, Mostafavi A, Mirtadzadini SM. Chemical compositions of *Salix aegyptiaca* L. obtained by simultaneous hydrodistillation and extraction. *J Essential Oil Bearing Plant* 2011; 14(5):543-48.
- [88] Dufresne CJ, Farnworth ER, A review of latest research findings on the health promotion properties of tea. *J Nutr Biochem* 2001; 12:404-21.
- [89] Parkouda C, Diawara B, Lowor S, Diako C, Saalia FK, Annan NT, Jensen JS, Tano-Debrah K, Jakobsen M. Volatile compounds of maari, a fermented product from baobab (*Adansonia digitata* L.) seeds. *Afr J Biotechnol* 2011; 10(20):4197-206.
- [90] Schlotzhauer WS, Pair SD, Horvat RJ. Volatile constituents from the flowers of Japanese honeysuckle (*Lonicera japonica*). *J Agri Food Chem* 1996; 44:206-09.
- [91] Cao W, Lin L, Niu Y, Xiao Z, Fang X. Characterization of aroma volatiles in camellia seed oils (*Camellia oleifera* Abel.) by HS-SPME/GC/MS and electronic nose combined with multivariate analysis. *Food Sci Technol Res* 2016; 22 (4):497-505.
- [92] Ali Sahari M, Ataii D, Hamedi M. Characteristics of tea seed oil in comparison with sunflower and olive oils and its effect as a natural antioxidant. *J Am Oil Chem Soc* 2004; 81:585-88.

- [93] He L, Guo-ying Z, Huai-yuan Z, Jun-ang L. Research progress on the health function of tea oil. *J Med Plants Res* 2010; 5:485-89.
- [94] Zeb A. Triacylglycerols composition, oxidation and oxidation compounds in *Camellia* oil using liquid chromatography-mass spectrometry. *Chem Phys Lipids* 2012; 165:608-14.
- [95] Culleré L, Ferreira V, Chevret B, Venturini ME, Sánchez-Gimeno AC, Blanco D. Characterisation of aroma active compounds in black truffles (*Tuber melanosporum*) and summer truffles (*Tuber aestivum*) by gas chromatography-olfactometry. *Food Chem* 2010; 122:300-06.
- [96] Jordan MJ, Margaria CA, Shaw PE, Goodner KL. Aroma active components in aqueous kiwi fruit essence and kiwi fruit puree by GC-MS and Multidimensional GC/GC-O. *J Agri Food Chem* 2002; 50:5386-90.
- [97] Leejeerajumnean A, Duckham SC, Owens JD, Ames JM. Volatile compounds in *Bacillus*-fermented soybeans. *J Sci Food Agri* 2001; 81:525-59.
- [98] Dakwa S, Sakyi-Dawson E, Diako C, Annan NT, Amoa-Awua WK. Effect of boiling and roasting on the fermentation of soybeans into dawadawa (soy-dawadawa). *Int J Food Microbiol* 2005; 104:69-82.