



Home Paper Archives Journal Indexing Research Conference Research Position

Main Menu

My Profile
My Journal
Add Journal

Searching By

Search More ...

PARTNERS

Like 6.9K

Tweet

Advertisement

Journal of Animal Science and Veterinary Medicine (ISSN: 2536-7099)

Publisher: Integrity Research Journals

ISSN-L: 2536-7099

E-ISSN: 2536-7099

IF(Impact Factor): 2019 Evaluation Pending

Website: <http://integrityresjournals.org/journal/JASVM>

- [1. Current topics in veterinary medicine and animal science](#)
- [2. Bulletin of University of Agricultural Sciences and Veterinary Medicine Cluj-Napoca - Animal Science and Biotechnologies](#)
- [3. Veterinary Medicine and Animal Sciences](#)
- [4. Veterinary medicine/small animal clinician](#)
- [5. Journal of Veterinary Medicine and Animal Health](#)

Description

Journal of Animal Science and veterinary medicine (JASVM) (ISSN: 2536-7099) (DOI: 10.31248/JASVM) is an open access, peer-review journal that publishes articles in all areas of Animal Science and Health such as Veterinary

**Advertise
Your
Business
Here**

**Text Link
AdBanner**

Contact Us ?

science, Animal husbandry, Animal nutrition, Anatomy, Biological science, Animal Pathology, Animal microbiology, Infectious diseases, Animal parasitology, Animal physiology, Animal breeding Animal biotechnology, Animal Genetics, Transgenic animal production, Veterinary medicine, Equine, Animal reproduction, Animal ethology, Animal products technology, Animal management and economics, Poultry farming etc.

The Journal is positioned to provide the most rapid system for reviewing and publishing. The articles published in JASVM will be of tremendous contribution to the field of medicine and Agriculture. Journal of Animal Science and Veterinary Medicine publishes per article and one issue every two months. All published articles are freely accessible and hard copies of the published articles will be made available to the interested authors on request.

Last modified: 2018-08-28 22:14:35

Volumes

[Vol.2 No.6](#)

[Vol.2 No.5](#)

[Vol.2 No.4](#)

[Vol.2 No.3](#)

[Vol.2 No.2](#)

[Vol.2 No.1](#)

[Vol.1 No.4](#)

[Vol.1 No.3](#)

[Vol.1 No.1](#)

[Vol.1 No.2](#)

About
[ResearchBib](#)
[Careers](#)
[Contact us](#)

Legal
[Terms of Service](#)
[Privacy Policy](#)
[Ad Choices](#)
[Cookies](#)

© 2019 ResearchBib. All Rights Reserved.

ICI World of Journals (/search/form) / **Search**



Search for journals adapted to your scientific interests. The Journal's Passports include information on editors, field of activity, publisher and indexation in the ICI Journals Master List database and evaluation by the Polish Ministry of Science and Higher Education (if applicable). Editorial offices that take care of scientific content distribution and apply principles of transparency also include the published issues, their contents and full texts of scientific articles (optional) in their Passports. Passports of scientific journals are updated and completed by their representatives.

Search

Journal title

 With citations only

 MSHE
points ⓘ

0

200

0

20

40

70

100

140

200

ICV points ICV ye ▼
0 200
?
0 50 100 150 200

Sorting ▼
Clear
Search

Search Results



Journal title: Journal of Animal Science and Veterinary Medicine (/search/details?id=50152)

ISSN: 2536-7099

GICID: n/d

Country / Language: NG / EN

Publisher: Integrity Research Journals

Citation: **1** ICV 2018: **ON**

MNISW 2019: **N/D** ICV 2017: **N/I**

Deposited publications: 18 > Full text: 95% | Abstract: 100% | Keywords: 100% | References: 100%



Journal title: Journal of Animal Science and Veterinary Medicine (/search/details?id=50152)

ISSN: 2536-7099

GICID: *n/d*

Country / Language: NG / EN

Publisher: Integrity Research Journals

Citation:	1	ICV 2018:	ON
MNISW 2019:	N/D	ICV 2017:	N/I

Deposited publications: 20 > Full text: 95% | Abstract: 100% | Keywords: 100% | References: 100%

Main page (<http://jml.indexcopernicus.com>)


© Index Copernicus 2017





Search by Journal Title

[Advanced search](#)



Open Academic Journals Index

Home

Library OAJI

Journals List

Statistics

Our Journals

TOP 50 Most Downloaded Articles

For Publishers

Apply for Evaluation / Free Service

Journal Search

OAJI Logo

Contact Us

Journal of Animal Science and Veterinary Medicine

Published by

[Integrity Research Journals](#)

Year publication 2016

Frequency 6

Article Publishing Frequency 2

0.101

CGIJ OAJI



Recently Added Articles

[Mehran University Research Journal of Engineering and Technology.](#)

[International Journal of Ayurveda and Pharmaceutical Chemistry.](#)

[University Economic Bulletin. Collection of scientific articles of scientists and post-graduate stu](#)

[Tecnura](#)

Register Journal**Visitors Counter**

 Today	931
 Yesterday	2106
 This week	17255
 Last week	13714
 This month	22427
 Last month	63140
 All days	3038044

Abbreviation	JASVM	Country	Nigeria		
ISSN (print)	-	ISSN (online)	2536-7099		
Journal Website	http://www.integrityresjournals.org				
Editor in Chief	Prof. SMITH, Olusiji Frederick				
ISI	—	All articles	<u>35</u>	Date added to OAJI	24 Apr 2018
Scopus	—	All issues	<u>17</u>	Free access	2016-2019
DOAJ	—	Full text language	English		

Journal discipline

Agricultural Sciences

Journals**208****Journal description**

Journal of Animal Science and Veterinary Medicine (JASVM) (ISSN: 2536-7099) (DOI: 10.31248/JASVM) is an Open Access, Peer-Reviewed Journal that publishes original articles in all areas of animal science and health. The Journal is positioned to provide the most rapid system for reviewing and publishing. Articles published in JASVM will be of tremendous contribution to the field of Medicine, Animal Science and Animal Health. JASVM publishes per article as soon as the manuscript is accepted. The Journals also publish e-books every quarter. All published articles and e-books are freely accessible on our website. Hard copies of the published articles and e-books will be made available to the interested authors on request.

Journal is indexed by

—



Menu

- [Home \(https://integrityresjournals.org/\)](https://integrityresjournals.org/)
- [About us \(https://integrityresjournals.org/about-us\)](https://integrityresjournals.org/about-us)
- [Our Journals \(https://integrityresjournals.org/journal\)](https://integrityresjournals.org/journal)
- [Publication Ethics \(https://integrityresjournals.org/publication-ethics\)](https://integrityresjournals.org/publication-ethics)
- [Copyright \(https://integrityresjournals.org/copyright\)](https://integrityresjournals.org/copyright)
- [Contact Us \(https://integrityresjournals.org/contact\)](https://integrityresjournals.org/contact)

Other links

- [Submit Manuscript \(https://integrityresjournals.org/submit-manuscript\)](https://integrityresjournals.org/submit-manuscript)
- [Track Manuscript \(https://integrityresjournals.org/track-manuscript\)](https://integrityresjournals.org/track-manuscript)
- [Processing Fees \(https://integrityresjournals.org/processing-fees\)](https://integrityresjournals.org/processing-fees)
- [Make Payment \(https://integrityresjournals.org/track-manuscript\)](https://integrityresjournals.org/track-manuscript)

Socia media

f () g+ () t ()


Journal of Animal Science and Veterinary Medicine (/journal/JASVM)



JOURNAL OF ANIMAL SCIENCE AND VETERINARY MEDICINE

Integrity Research Journals

ISSN: 2536-7099

Model:  Open Access/Peer Reviewed

DOI: 10.31248/JASVM

Start Year: 2016

Email: jasvm@integrityresjournals.org

[Submit Manuscript \(/submit-manuscript\)](/submit-manuscript)

Journal of Animal Science and Veterinary Medicine (JASVM) (ISSN: 2536-7099) is an Open Access, Peer-Reviewed Journal that publishes original articles in all areas of animal science and health. The Journal is positioned to provide the most rapid system for reviewing and publishing. Articles published in JASVM will be of tremendous contribution to the field of Medicine,... Continue here (/journal/JASVM/about_jasvm)

CURRENT ISSUE (Vol.4 (4) August 2019)

[View Complete Issue \(/journal/JASVM/edition/30_Aug_2019\)](/journal/JASVM/edition/30_Aug_2019)

Laying Hen Strains And Breeding System: A Review Of Hen Behavior And Welfare, Hen Production Performance And Eggs Quality Traits In Africa And Around The World (</journal/JASVM/article-abstract/9BA0F91A1>)

Authors: Tossou M. L., Houndonougbo P. V., Akouedegni, C. G., Houndonougbo M. F., Chrysostome C. A. A. M. and Abiola F. A.

DOI: 10.31248/JASVM2019.130 | **Article Number:** 9BA0F91A1

The breeding system for laying hens has been the focus of scientific research for many years. During the last decade, new laying hen systems were rapidly introduced with the aim of improving the health of poultry and the welfare of consumers, producers and industries on the one hand and meeting the requirements on the other hand. The choice of the genetic type... Continue here (<https://integrityresjournals.org/journal/JASVM/article-abstract/9BA0F91A1>)

Squamous Cell Carcinoma On The Eyelid Of Nigerian Albino Horse (</journal/JASVM/article-abstract/FD3A3C2F2>)

Authors: Garba, U. M.* Nwokike, E. C., Mbuk, E. U., Ali, A., Zamdayu, J. M., Yunusa, H. G., Amaje, J. and Auwal, U.

DOI: 10.31248/JASVM2019.136 | **Article Number:** FD3A3C2F2

Squamous cell carcinoma is the most common neoplasia of the skin in horses. An adult albino stallion of Nigerian indigenous (Arewa) breed used for military ceremonial parade was presented with the chief complaints of growth on left side lower eyelid and unilateral epiphora. Tentatively diagnosis of eyelid squamous cell carcinoma was made and managed successfully.... Continue here (<https://integrityresjournals.org/journal/JASVM/article-abstract/FD3A3C2F2>)

Artificial Insemination In Botswana: Challenges And Opportunities - A Review (</journal/JASVM/article-abstract/49071F933>)

Authors: John Cassius Moreki, Tiyapo Daisy Pelaelo-grand and Aubrey Ranko

DOI: 10.31248/JASVM2019.150 | **Article Number:** 49071F933

Artificial insemination (AI) was introduced in Botswana in 1966 to provide smallholder cattle farmers with access to superior bulls for cross-breeding to improve their herd quality and performance. This review endeavours to highlight farmers' uptake of AI service, challenges and the opportunities in the use of AI technology in Botswana. The 15 AI centres... Continue here (<https://integrityresjournals.org/journal/JASVM/article-abstract/49071F933>)

Effect Of Roselle (hibiscus Sabdariffa Linn) Calyx Residue On Growth Performance, Haematological And Serum Biochemical Indices Of Weaner Rabbits (</journal/JASVM/article-abstract/A68938F74>)

Authors: Omolade O. Awodola-peters,, Mariam O. Yahaya and Oluwafunmilayo O. Adetola

DOI: 10.31248/JASVM2019.149 | **Article Number:** A68938F74

The study was carried out to determine the growth performance, haematological and serum biochemical indices of weaner rabbits fed roselle (Hibiscus sabdariffa) calyx residue. Thirty (30) rabbits were allotted into five treatments with six rabbits per treatment. Each treatment was replicated three times with two

rabbits per replicate. The rabbits were fed... Continue here
<https://integrityresjournals.org/journal/JASVM/article-abstract/A68938F74>

Indigenous Pig Farming In Rural Areas Of Sierra Leone: Practices, Constraints And Impact On Livelihood (/journal/JASVM/article-abstract/DD95C02C5)

Authors: Abdulai Mahmood Conteh* and Brima Gogra

DOI: 10.31248/JASVM2019.145 | **Article Number:** DD95C02C5

Poverty and inequality between urban and rural areas remained major social issues in Sierra Leone. These are drivers forcing rural people to engage in feasible livelihood options such as animal husbandry. The current study, that lasted for seven (7) months attempt to investigate pig farming in rural areas of Moyamba district, Sierra Leone. The overarching... Continue here
<https://integrityresjournals.org/journal/JASVM/article-abstract/DD95C02C5>

[View Complete Issue \(/journal/JASVM/edition/30_Aug_2019\)](/journal/JASVM/edition/30_Aug_2019)

Integrity Research Journals

a research publishing organization owned and managed by Integrity Mega Research Publishers, an international, open access, peer-reviewed publisher committed to advancing knowledge across the globe by facilitating access to high quality scholarly articles of broad interest through online publication.

Search by Title, Article Num



Company



- About Us (<https://integrityresjournals.org/about>)
- Open Access (<https://integrityresjournals.org/open-access>)
- Journals (<https://integrityresjournals.org/journal>)
- Conferences (<https://integrityresjournals.org/conferences>)
- Contact (<https://integrityresjournals.org/contact>)

Our Policies

- Waiver Policy (<https://integrityresjournals.org/waiver-policy>)
- Authors Guidelines (<https://integrityresjournals.org/author-guidelines>)
- Publication Ethics (<https://integrityresjournals.org/publication-ethics>)
- Privacy Policy (<https://integrityresjournals.org/privacy>)
- Terms of Use (<https://integrityresjournals.org/terms>)


Stay updated

Sign up if you would like to receive the newest updates and other free goods.

Enter your Email

Go

Copyright © 2019 Integrity Research Journals. All rights reserved.



Laying hen strains and breeding system: A review of hen behavior and welfare, hen production performance and eggs quality traits in Africa and around the world

Tossou M. L., Houndonougbo P. V., Akouedegni, C. G.*, Houndonougbo M. F., Chrysostome C. A. A. M. and Abiola F.A.

UAC-FSA – Laboratoire de Recherches Avicoles et de Zoo Economie 03 BP 2819 Cotonou-Benin.

*Corresponding author. Email: akouedegniguenole@gmail.com; Tel: (+229) 95177790.

Copyright © 2019 Tossou et al. This article remains permanently open access under the terms of the [Creative Commons Attribution License 4.0](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Received 16th January, 2019; Accepted 18th February, 2019

ABSTRACT: The breeding system for laying hens has been the focus of scientific research for many years. During the last decade, new laying hen systems were rapidly introduced with the aim of improving the health of poultry and the welfare of consumers, producers and industries on the one hand and meeting the requirements on the other hand. The choice of the genetic type usually depends on the production performance and the color of the eggs. To this end, the aim of this work was to make a comparative review of the results obtained by different authors on effect of strain and farming system on production performance (egg production and mortality) and the characteristics of egg quality (egg weight, proportion of different parts of the egg, Haugh units, egg yolk color and carotenoid) of laying hens. Although the productive performance in ground systems is often low compared to the cage farming system, the eggs of the ground-based system have been proven by many studies to have the best nutritional properties. This review of literature on the use of different strains and different table egg farming systems could give a new direction to research.

Keywords: Egg quality, farming systems, laying hen, traits.

INTRODUCTION

Commercial egg production in industrial systems starts to increase rapidly in the United States after the second world war and much later in Western Europe and Africa. Its main features include containment facilities, illumination and artificial ventilation, a large number of highly productive battery-powered hybrid laying hens which provide a small space, the use of a complete ration (which may contain antibiotics, stimulants, hormones, artificial colors, etc.), the use of several hygiene conditions and cleaning products. This type of production provides ready-to-eat eggs in large quantities each year at low prices (Pavlovski et al., 2010), because the battery-based farming system has become the most popular egg-producing system in the world (Matt et al., 2011).

However, in recent decades, new laying hen systems were rapidly introduced to improve the health of poultry

and the well-being of consumers, producers, industries and environmental requirements (Matković et al., 2007). It is also known that genetic origin affects egg production and determines the type of feed to be used (Reiter and Kutritz, 2001). Several breeders in the US are using improved soil fertility and healthy food production as benefits from this type of production (Hilimire, 2012). Henry (2002) reported that the most common breeding systems for laying hens are free-range and organic farming. Most of these farming systems include hens of different ages, subjected to a natural environment, fresh air, inorganic rocks (inorganic stone) and organic ration (plant and animal bases) of the degree that can allow increase in their daily nutritional requirements by 20% (Henry, 2002; Oberholtzer et al., 2006). In addition, birds are exposed to environmental climatic factors for both pest

and predator attacks.

Some countries have already achieved a significant percentage of eggs from alternative livestock systems. Sales of organic eggs in the US have grown at a 20% annual rate since the 1990s, suggesting that the organic egg market is the fastest growing part of the organic agriculture sector (Oberholtzer et al., 2006). Among European countries, Italy has the largest market for organic products, followed by Spain (Mesías et al., 2011).

Three types of systems may involve the use of hybrid or local livestock adapted to existing environmental conditions. The genetic type used often depends on the color of the eggs. White leghorns are among the most popular and productive layers, while Rhode Island Red comes in second and produces most brown eggs in the US (Burbaugh et al., 2010). Hilimire (2012) reported that Plymouth Rock and Ameraucanas are primarily used for the production of red-shelled eggs. The main problems associated with rearing commercial hybrids are strictly controlled rearing conditions in three types of systems including weak eggs, excessive heart attack, high incidence of congestive heart failure (ascites), poor foraging capacity and a low heat tolerance and also, producers are often discouraged from breeding these hybrids in alternative production systems.

EFFECT OF STRAIN AND BREEDING SYSTEM ON EGG PRODUCTION

Egg production is one of the most important production characteristics of laying hens. It is equivalent to the number of eggs laid per year expressed as a percentage. A large number of researchers have studied this parameter through different farming systems. Anderson et al. (2007) revealed that there are statistically significant differences between egg production for brown Hy line layers reared in cage and ground respectively. Castellini et al. (2006) compared white caged Leghorn individuals in an organic and inorganic system. Birds in organic systems showed a low rate of egg production as controls (organic 63.4%, organic-plus 61.4%, control 74.1%). This difference is probably due to the high motor activity and simultaneously to the low energy and protein intake that is diluted by grass consumption mainly in the batch of organic-plus subjects. In addition, compared to the control, the spawning rate of organic birds is much more affected by seasonal conditions (temperature and photoperiod). A similar experiment using Ancona laying hens was conducted by Mugnai et al. (2009) which led to similar results (spawning rate: organic 60.6%, organic plus 59.6%, control 70.9%). Differences in productive performance between the cage system (80.82%) and the free-range system (72.88%) were obtained by Senčić and Butko (2006) with brown Lohman breeding hens as well by Yakubu et al. (2007) who found better production in Brown and Brown Lohman caged (74%) than in litter (69%) in

terms of spawning performance. Similar results were published by Suto et al. (1997) and by Flock et al. (2005).

A study conducted by Ferrante et al. (2009) revealed a similarity in nesting rates of Hy-line birds raised under organic and cage systems. The spawning peak was reached at 25 weeks and was higher in organic layers than in the cage layers (94.5% vs 93%), while 25 to 58 weeks, the lay rate was high among the organic layers caged subjects with a final result, however, being identical (86.40% vs. 86.35%). These results are different from those of Tauson and Holm (2001), who found 3% fewer eggs in chickens compared to caged subjects. This difference may be due to strong management in both fields of study. Barbosa et al. (2005) found no difference in spawning performance of Hy-line W36 and Brown Hy-line on litter and cage. Similarly, Basmacioğlu and Ergul (2005) found no difference in Babcock-300 and Isa Brown, both in litter and cage systems. Gerzilov et al. (2012) found no significant difference in conventional and litter-reared Isa Brown hens in terms of oviposition rate. Hetland et al. (2004) reported that enriched cage hens consume more feed compared to hens raised in conventional cages. Meanwhile, Elson and Croxall (2006) demonstrated that feeding is lower in hens maintained in an enriched cage compared to conventional cages. However, Valkonen et al. (2008) found no significant difference in enriched caged Lohmann and Leghorn on productivity and feed conversion. A study by Van Horne and Van Niekerk (1998) showed that feed conversion is less effective in the aviary and free range system compared to conventional system. In the alternative system, hens need to spend extra energy on producing heat and moving, which is often due to storage density and low temperature (Preisinger, 2000). Other studies conducted in several European countries indicate that the production of eggs in furnished cages is comparable to that obtained in conventional cages (Abrahamsson and Tauson, 1997). In contrast, Pohle and Cheng (2009a) reported that hens reared in furnished cages laid more eggs at 40 weeks compared to conventional caged hens because of significant improvements in feeding levels behavior of well-being. These results found that some genotypes may perform better in conventional cage culture systems, while other genotypes may respond positively to litter-free soil.

EFFECTS OF STRAIN AND REARING SYSTEM LAYING HEN MORTALITY

When analyzing mortality in chickens reared under ambient conditions, several adverse circumstances and conditions can cause birds to die (Bogosavljević-Bošković et al., 2012). In a free cage production system, the system itself, is a problem. Sossidou et al. (2011) showed high mortality in these production systems due to the broad spectrum of diseases and infections due to ambient conditions. Newcastle disease is reputed to be one of the

most important diseases in the cage culture system. During the first manifestations of disease, nearly 80% of the subjects may die (Permin and Pedersen., 2002). Several main reasons for the movement of boxed poultry controlled the facilities included in diseases, parasites and predator attacks (Fanatico, 2000).

In all livestock systems, birds can develop abnormal behaviors such as stinging and cannibalism, which are the major causes of mortality with laying hens (Rakonjack et al., 2013). Blokhuis et al. (2007) studied different layer rearing systems and found that one-third of mortalities are caused by stinging and cannibalism, while Weitzenburger et al. (2005) found that cannibalism matters for more than 65.5% of the mortality of laying hens raised in various cages. Oden et al. (2002) found that rearing white feather genotypes can lead to reduced mortality of laying hens due to their susceptibility to stinging and cannibalism as primary causes of mortality. However, Van de Weerd et al. (2009) have recommended the opposite by proposing the breeding of non-white layers in a free rearing system in order to reduce the action of infectious agents responsible for mortalities. Hegelund et al. (2006) in his method showed that the system of organic production can be a potential cause of health problems, especially when the surroundings of the outdoor farm are not cleaned properly. Mortality is also high (11 to 18%) in large, organically raised flocks of poultry but can be reduced to 7 to 9% in smaller organic flocks (Berg 2002). When mortality exceeds 20% in the organic system, it is very often due to cannibalism. Given the threats associated with different housing systems, organic production with small numbers of birds is recommended (Bestman, 2004). A common problem encountered in poultry production in organic production is salmonellosis. The comparison of the battery system and the organic production system shows that battery-reared hens experience high mortality and are threatened by zoonoses and salmonellosis (Fiks-Van Niekerk et al., 2003). However, Van Overbeke et al. (2006) maintain that organic poultry production is more at risk from salmonellosis as in many other production systems.

EFFECTS OF STRAIN AND REARING SYSTEM ON EXTERNAL EGG QUALITIES

Egg weight is mainly affected by chick genotypes (Alkan et al., 2008). In addition, egg-laying performance is an important factor influencing egg weight. Castellini et al. (2006) reported that egg weight is negatively correlated with egg deposition rate. The most productive birds regardless of group and season produced lighter eggs. Other important factors are the age of the hens (Rizzi and Chiericato, 2010; Rizzi and Cassandro, 2009; Akyurek and Okur, 2009; Škrbić et al., 2011) and the energy of the food and its nutritional value, particularly the essential amino acid content and the level of protein (Elwinger et al., 2002; Krawczyk, 2009). Eggs produced in free-range systems have higher average weights than those raised in battery

and conventional cages (Hughes et al., 1985; Hidalgo et al., 2008). Hughes et al. (1985) found that differences in egg weights may be due to changes in environmental temperatures in the box or in cage systems. However, Senčić and Butko (2006) obtained higher weights with layer-grown lohman brown hens (62.40 g) compared to those raised in cages (60.50 g). On the contrary, several authors have found that the average weight is low for layers reared in hut. Krawczyk (2009) attributes this result to the fact that animals can not satisfy their protein requirement in box breeding. Pištekova et al. (2006) found that Isa Brown layers raised on litter lay heavier eggs (62.02 g) than those raised in cages (60.63 g). Yakubu et al. (2007) found high weights eggs with Brown Bovans and Lohman Brown layers raised in cage systems compared to litter grown. Minelli et al. (2007) reported that the farming system itself, has a crucial impact on the weight of the egg; moreover, the effect of the ration is much more significant. Different farming systems have considerable effects on egg quality, including their physicochemical properties (Trziszka et al., 2004; Giannenas et al., 2009; Matt et al., 2009). Hidalgo et al. (2008) compared eggs purchased from supermarkets from different farming systems and found differences in shell percentage.

Another issue is the dirt of the eggshell and the microbiological condition of the shell. Dirty eggs have been found in litter systems compared to conventional cage systems (Leyendecker et al., 2001a), which may be due to nests or litter. Comparison of conventional cages and furnished cages shows that dirty eggs are less common in the furnished cage system (Abrahamsson and Tauson, 1997). However, De Reu et al. (2009) do not observe significant differences between the percentage of dirty eggs from furnished cages and conventional cages. Tauson et al. (1999), who compared three production systems (conventional cages, aviary and soil) and two bird genotypes (Lohman Selected Leghorn and Lohman Brown), found no differences in soil contamination. Eggs produced in different housing systems of Lohman Brown hens, but the eggs produced by the Lohman Select leghorn hens on the ground are less dirty than those of Lohman Selected Leghorn produced by the hens raised in the cage and in the aviary system. Protais et al. (2003) and De Reu et al. (2005) have shown that aviary egg shells are more contaminated with aerobic bacteria compared to egg shells in the cage system (furnished and conventional cages). De Reu et al. (2009) also counted fewer bacteria on furnished cage egg shells compared to egg shells obtained in the alternative system (4.75 vs. 4.98 cfu/shell).

However, the same authors did not find significant differences with respect to the percentage of enterobacteria on eggshells between cages and alternative systems. Wall et al. (2002) compared conventional cages and furnished cages, found significant differences between the percentages of enterobacterial contamination counted on eggshells (12.3% in furnished cages versus 5.8% in conventional cages). Schwarz et al. (1999) reported that outdoor hen eggs are characterized by higher aerobic bacterial

contamination compared to conventional cage eggs. However, De Reu et al. (2005) found no differences between negative grams of bacteria found on egg shells produced in conventional cages, furnished cages and aviaries. De Reu et al. (2009) reported that this bacterial contamination of eggs is determined not only by the system of production but also by the organization and management of the farm.

Wall et al. (2002) and Guesdon and Faure (2004) found that the number of cracked eggs was greater in furnished cages than in traditional cages. The eggs in the furnished cages is small and the eggs can go into one or the other. Meanwhile, Leyendecker (2003) showed a lower proportion of cracked eggs in furnished cages compared to conventional cages. In addition, the different inclinations of the cage floor can influence the number of cracked eggs (Valkonen et al., 2008). Similarly, De Reu et al. (2009) found a higher percentage of cracked eggs ($p < 0.01$) in furnished cages (7.8%) compared to an alternative production system. Guesdon et al. (2006) showed in their study of different hen housing systems that the proportion of cracked and broken eggs was higher in furnished cages (15.4, 19.6%) compared to standard cages (8.1, 12.2%). Hidalgo et al. (2008) found no significant differences in the quality of the shell of eggs produced in cages, in the open air and in organic systems. According to Pohle and Cheng (2009a), the Leghorn Blanc W-36 housing system reared in conventional, 19-week-old cages has no effect on shell thickness. Evidence previously observed shows that the production system has no conclusive effect on the number of dirty and cracked eggs, while the highest bacterial contamination of shells is characteristic of alternative systems.

EFFECTS OF STRAIN AND REARING SYSTEM ON INTERNAL EGG QUALITIES

Haugh unit is the objective measure of the quality of the egg based on the logarithm function of the albumen's height and the weight of the egg. Several factors may influence the Haugh unit: storage time and temperature, age of the hen, breeding system, strain and diet (dietary protein, essential amino acid composition e.g. lysine, methionine, food enzymes, protein sources), disease, supplements (ascorbic acid and vitamin E) and artificial exposure to ammonia (Roberts, 2004).

Carotenoids are natural pigments found in egg yolks. They give it the yellow color that can range from very pale yellow to bright dark or orange. Carotenoids represent less than 1% of egg yolk lipids mainly composed of carotene and xanthophyll (lutein, cryptoxanthin and zeaxanthin). The total concentration of lutein and zeaxanthin is ten times higher than that of combined cryptoxanthin and carotene (Shenstone, 1968). Lutein xanthophyll and zeaxanthin have been of great interest since it is known that an increase in the nutritional intake of both components can prevent adult macula degeneration (AMD) and cataract formation in

adulthood (Schlatterer and Breithaupt, 2006).

Economically, color is very important because it is a quality criterion for the consumer (Matt et al., 2011). The intensity of egg yolk is measured using the Roche scale. A dark color is much appreciated by consumers in different countries, for example in Australia, the preferred yellow color is 11 on the Roche scale (Roberts, 2004).

EFFECT OF STRAIN AND SYSTEM ON HEN BEHAVIOR

Among the factors that influence hen behavior are the housing system (Oden et al., 2002, Anderson et al., 2004, Whay et al., 2007), the density of occupancy (Albentosa et al. 2007), and the microclimate (Herbut et al., 2002, Prescott and Wathes, 2002). According to Weeks and Nicol (2006), the well-being of hens raised in conventional cages is compromised by the lack of nests. On the other side, enriched cages have nests and swims that allow birds to express their natural behavior such as nesting and swimming (Tauson, 2002). Yue and Duncan (2003) observed frustration and stereotypy of birds unable to use nests. Appleby et al. (2004) reported that birds have a strong instinct for finding a nest to lay eggs. Pohle and Cheng (2009b) observed that 25 to 41% of the birds analyzed used perches during the day. Appleby (1998) reported that approximately 80% of birds used perch at night. Also, Duncan et al. (1992) reported that 99% of chickens used approximately poles during the night. Pohle and Cheng (2009b) and Appleby et al., (2002) also found that birds prefer litter for pecking, resting and feathering by sandbaths. However, a sandbox can positively affect other forms of behavior such as pecking, feathering by sandbath and resting. Bird activity increases with large numbers per group, when the total size of the cage increases (Carey et al., 1995). However, cages with perches make birds less active (Matsui et al., 2004). Johnsen et al. (1998) reported that caged birds take longer to eat compared to aviary birds.

The systems of outdoor poultry production are suitable at present and are becoming more and more popular. This aviary housing system has a potentially positive impact on wellbeing because it allows them to perform natural behaviors such as moving, scraping, pecking, foraging and feeding (Mahboub et al., 2004, Tuytens et al., 2005). Freire et al., (2003) found that for a large number of raised hens, the outdoor system can have a positive effect on bird welfare by reducing their aggressive behavior. However, according to Hegelund et al. (2006), this production system can cause health problems, especially when it is not used properly. Green et al. (2000) reported that the frequency of birds that eat in the outdoor system increases during hot, sunny days. Also, Weitzenbürger et al. (2005) reported that cannibalism is a major problem in birds raised in both conventional and alternative production systems. They also reported that the protection of the outdoor system from predators is inadequate, which often

causes strong stress responses and makes birds less passionate about using them. Shimmura et al., (2008) did not observe the increase in the number of birds eating and cannibalism in birds raised outdoors. On the contrary, they believe that this type of farming system traps birds and prevents them from expressing their natural behavior. Mahboub et al. (2004) and Bilçik et al. (1999) reported that an increase in the number of birds per flock makes them more aggressive. Väisänen et al. (2005) show that shown that chicken lines have a lower capacity and cope with disturbances within their social group compared to their ancestors. Meanwhile, Anderson et al. (2007) believe that, in the long run, the reproductive genetic selection of the egg-type will make firm to enhance production parameters but will have no impact on the behavior of the offspring in the next cycles of production.

Important factors affecting bird behavior are also their genetic origin (Nielsen et al., 2003; Mahboub et al., 2004) and herd numbers (Reiter and Bessei 2000; Zeltner and Hirt 2003). Mahboub et al. (2004) showed differences in the behavior of two commercial hen lines. LSL (Lohman Selected Leghorn) hens were characterized by more movement on grassland but spent less time compared to Lohman Traditional hens. According to Nielsen et al. (2003), breeds of breeds kept in the open-air system frequently use the area around the farm and show much greater movement compared to hens of commercial lines that spend more time in the field to lie down and to feed.

In summary, avian behavior depends largely on the housing system. Hens raised in enriched cages and with outdoor access easily express their natural behavior which has a favorable effect on their well-being. Bird behavior is also determined by their genetic origin.

EFFECT OF STRAIN AND SYSTEM ON THE WELFARE OF HENS

According to Appleby et al. (2004), conventional cages do not provide the opportunity for birds to experience all of their freedom that is at the root of animal welfare and with the major consequences of preventing birds from expressing their natural behavior. Kopka et al. (2003) and Leyendecker et al. (2005) stated that, furnished cages improve hen well-being by reducing stress, aggression and stitching, and improve bone mineralization. However, Guesdon and Faure (2004) did not show a salutary effect on the welfare of birds reared in furnished cages compared to birds reared in traditional cages. Comparing conventional cage enriched, Barnett et al. (2009) found that herd size and space have a small effect on bird welfare while cage material (perch, nest sand bath, nest) has no influence on the well-being of birds and on measured physiological parameters, although it has a positive effect on bone strength. The presence of leg deformities in hens housed in furnished cages is likely due to the use of excessive perch, which may be a problem in this type of cage (Vits et al., 2005).

The open-air production system has a positive effect on the well-being of birds as it allows them to express their natural behavior such as moving, scratching, pecking, foraging and eating (Mahboub et al., 2004; Tuytens et al., 2005). The use of the outdoor system not only boosts bird immunity, but helps them reduce their stress during rearing (Bestman et al., 2003). The production system also has an effect on leg diseases. The strength of bone fracture is less in conventional cages than in alternative cage systems (Leyendecker et al., 2001). By comparing conventional cages with furnished cages and aviaries, Leyendecker (2003) has shown that the force of the humerus is considerably higher in the furnished cages compared to conventional cages, but it was not greater than in the aviaries. In furnished cages, the increase in the number of birds raised in a cage can negatively affect the plumage (Appleby et al., 2002; Hetland et al., 2004; Weitzenbürger et al., 2005). Other research provides evidence that perch-equipped cages and litter bedding improve the plumage of hens (Abrahamsson and Tauson, 1997). The length of the claw is also dependent on the production system. In alternative production systems, birds have an opportunity to equalize their claws by moving and scratching the litter, which is different from hens raised in conventional cages where the length of the claws increases the risk of injury (Vits et al., 2005) and weakens the movement. This problem has been solved in partially furnished cages, although the views on this matter are divided. The system of production has a considerable effect on the well-being of the bird. It appears that free-range diaper systems are better at providing high levels of well-being because they reduce the stress of livestock birds while they are empowered to perform their natural behavior.

CONCLUSION

This review emphasis the need for additional research and observations has shown differences in the adaptability of different breeds of hens raised in different production systems. This will make it possible to optimize housing conditions in accordance with the principles of well-being and to make an appropriate choice of species or breeds in commercial lines to maximize economic and production outcomes; and to have hens resistant to the stress of production. The system of production of organic poultry is another matter and its principles are prescribed by separate regulations.

CONFLICT OF INTEREST

The authors declare that they have no conflict of interest.

REFERENCES

Abrahamsson, P., & Tauson, R. (1997). Effects of group size on

- performance, health and birds' use of facilities in furnished cages for laying hens. *Acta Agriculturae Scandinavica A—Animal Sciences*, 47(4), 254-260.
- Akyurek, H., & Okur, A. A. (2009). Effect of storage time, temperature and hen age on egg quality in free-range layer hens. *Journal of Animal Veterinary Advances*, 8(10), 1953-1958.
- Albentosa, M. J., Cooper, J. J., Luddem, T., Redgate, S. E., Elson, H. A., & Walker, A. W. (2007). Evaluation of the effects of cage height and stocking density on the behaviour of laying hens in furnished cages. *British Poultry Science*, 48(1), 1-11.
- Alkan, S., Karsli, T., Galiç, A., & Karabağ, K. (2013). Determination of phenotypic correlations between internal and external quality traits of Guinea Fowl eggs. *Kafkas Universitesi Veteriner Fakültesi Dergisi*, 19(19), 861-867.
- Anderson, K. E., Davis, G. S., Jenkins, P. K., & Carroll, A. S. (2004). Effects of bird age, density, and molt on behavioral profiles of two commercial layer strains in cages. *Poultry Science*, 83(1), 15-23.
- Anderson, K. E., Jones, D. R., Davis, G. S., & Jenkins, P. K. (2007). Effects of genetic selection on behavioral profiles of Single Comb White Leghorn hens through two production cycles. *Poultry Science*, 86(9), 1814-1820.
- Appleby M.C., Mench, J. A., Hughes, B. O. (2004). Poultry behaviour and welfare. *CABI Publishing, Wallingford, UK*.
- Appleby, M. C. (1998). Modification of laying hen cages to improve behavior. *Poultry Science*, 77(12), 1828-1832.
- Appleby, M. C., Walker, A. W., Nicol, C. J., Lindberg, A. C., Freire, R., Hughes, B. O., & Elson, H. A. (2002). Development of furnished cages for laying hens. *British Poultry Science*, 43(4), 489-500.
- Barbosa Filho, J. A. D., Silva, M. A. N. Silva, I. J. O., & Coelho, A. A. D. (2005) Egg quality in layers housed in different production systems and submitted to two environmental conditions. *Brazilian Journal of Poultry Science*, 8(1), 23-28.
- Basmacioğlu, H., & Ergül, M. (2005). Research on the factors affecting cholesterol content and some other characteristics of eggs in laying hens the effects of genotype and rearing system. *Turkish Journal of Veterinary and Animal Sciences*, 29(1), 157-164.
- Barnett, J. (2009). The prize of peace (is eternal vigilance): A cautionary editorial essay on climate geopolitics. *Climate Change*, 96(1), 1-6.
- Berg, C. (2002). Health and welfare in organic poultry production. *Acta Veterinaria Scandinavica*, 43(1), 37-45.
- Bestman, M., (2004). Health in organic laying hens – facts and fairy tales. *Proc. of the 2nd SAFO Workshop, Witzenhausen, Germany*, Pp. 145–152.
- Bestman, M. W. P., & Wagenaar, J. P. (2003). Farm level factors associated with feather pecking in organic laying hens. *Livestock Production Science*, 80(1-2), 133-140.
- Blokhuys, H. J., Fiks Van Niekerk, T., Bessei, W., Elson, A., Guémené, D., Kjaer, J. B. & Van De Weerd, H. A. (2007). The Lay Wel project: welfare implications of changes in production systems for laying hens. *World's Poultry Science Journal*, 63(1), 101-114.
- Bogosavljević-Bošković, S., Rakonjac, S., Dosković, V., & Petrović, M. D. (2012). Broiler rearing systems: a review of major fattening results and meat quality traits. *World's Poultry Science Journal*, 68(02), 217-228.
- Burbaugh, B., Toro, E., & Gernat, A. (2010). Introduction to Pasture-Raised Poultry: Getting Started AN232. *Animal Science Department, Florida Cooperative Extension Service, Institute of Food and Agricultural Sciences, University of Florida*.
- Carey, J. B., Kuo, F. L., & Anderson, K. E. (1995). Effects of cage population on the productive performance of layers. *Poultry Science*, 74(4), 633-637.
- Castellini, C., Perella, F., Mugnai, C., & Dal Bosco, A. (2006). Welfare, productivity and qualitative traits of egg in laying hens reared under different rearing systems. Poster at *XII: European poultry conference, Verona*, 10-14 September.
- De Reu, K., Grijspeerdt, K., Heyndrickx, M., Zoons, J., De Baere, K., Uyttendaele, M., Debevere, J., & Herman, L. (2005). Bacterial eggshell contamination in conventional cages, furnished cages and aviary housing systems for laying hens. *British Poultry Science*, 46(2), 149-155.
- De Reu, K., Rodenburg, T. B., Grijspeerdt, K., Messens, W., Heyndrickx, M., Tuytens, F. A. M., Sonck, B., Zoons, J., & Herman, L. (2009). Bacteriological contamination, dirt, and cracks of eggshells in furnished cages and noncage systems for laying hens: An international on-farm comparison. *Poultry Science*, 88(11), 2442-2448.
- Duncan, E. T., Appleby, M. C., & Hughes, B. O. (1992). Effect of perches in laying cages on welfare and production of hens. *British Poultry Science*, 33(1), 25-35.
- Elson, H. A., & Croxall, R. (2006). European study on the comparative welfare of laying hens in cage and non-cage systems. *Archiv Fur Geflügelkunde*, 70(5), 194-198.
- Elwinger, K., Tufvesson, M., & Hartmann, C. (2002). Genotype-environment interactions in layers housed in an organic feed environment. *Archiv für Geflügelkunde*, 66, 32.
- Fanatico, A. (2000). Pastured Poultry: An HI Case Study Booklet. *ATTRA publication*. National Center for Appropriate Technology Fayetteville, AR.
- Ferrante, V., Lolli, S., Vezzoli, G., & Cavalchini, L. G. (2009). Effects of two different rearing systems (organic and barn) on production performance, animal welfare traits and egg quality characteristics in laying hens. *Italian Journal of Animal Science*, 8(2), 165-174.
- Fiks-Van Niekerk, T. G. C. M., Van Emous, R. A., & Reuvekamp, B. J. F. (2003). Experiences with production and egg quality in alternative systems and large enriched cages for laying hens in the Netherlands. In *Proc. Xth European Symposium on the Quality of Eggs and Egg Products, Saint-Brieuc, France* (Vol. 111, pp. 210-216).
- Flock, D. K., Laughlin, K. F., & Bentley, J. (2005). Minimizing losses in poultry breeding and production: How breeding companies contribute to poultry welfare. *World's Poultry Science Journal*, 61(2), 227-237.
- Freire, R., Wilkins, L. J., Short, F., & Nicol, C. J. (2003). Behaviour and welfare of individual laying hens in a non-cage system. *British poultry Science*, 44(1), 22-29.
- Gerzilov, V., Datkova, V., Mihaylova, S., Bozakova, N. (2012). Effect of poultry housing systems on egg production, *Bulgarian Journal of Agricultural Science*, 18(6), 953-957.
- Giannenas, I., Nisianakis, P., Gavriil, A., Kontopidis, G., & Kyriazakis, I. (2009). Trace mineral content of conventional, organic and courtyard eggs analysed by inductively coupled plasma mass spectrometry (ICP-MS). *Food Chemistry*, 114(2), 706-711.
- Green, L. E., Lewis, K., Kimpton, A., & Nicol, C. J. (2000). Cross-sectional study of the prevalence of feather pecking in laying hens in alternative systems and its associations with management and disease. *Veterinary Record*, 147(9), 233-238.
- Guesdon, V., & Faure, J. M. (2004). Laying performance and egg

- quality in hens kept in standard or furnished cages. *Animal Research*, 53(1), 45-57.
- Guesdon, V., Ahmed, A. M. H., Mallet, S., Faure, J. M., & Nys, Y. (2006). Effects of beak trimming and cage design on laying hen performance and egg quality. *British Poultry Science*, 47(1), 1-12.
- Hegelund, L., Sørensen, J. T., & Hermansen, J. E. (2006). Welfare and productivity of laying hens in commercial organic egg production systems in Denmark. *NJAS-Wageningen Journal of Life Sciences*, 54(2), 147-155.
- Hegelund, L., Sørensen, J. T., Kjaer, J. B., & Kristensen, I. S. (2005). Use of the range area in organic egg production systems: effect of climatic factors, flock size, age and artificial cover. *British poultry science*, 46(1), 1-8.
- Henry R. (2002). Organic Poultry-Eggs. *Maritime Certified Organic Growers-Organic Profiles*. March 2002.
- Herbut, E., Sosnowka-Czajka, E., Rychlik, I., & Sokolowicz, Z. (2002). Welfare of chickens reared under different thermal conditions. *Annals of Animal Science. Supplement*, 1, 71-74.
- Hetland, H., Moe, R. O., Tauson, R., Lervik, S., & Svihus, B. (2004). Effect of including whole oats into pellets on performance and plumage condition in laying hens housed in conventional and furnished cages. *Acta Agriculturae Scandinavica, Section A—Animal Science*, 54(4), 206-212.
- Hidalgo, A., Rossi, M., Clerici, F., & Ratti, S. (2008). A market study on the quality characteristics of eggs from different housing systems. *Food Chemistry*, 106(3), 1031-1038.
- Hilimire, K. (2012). The grass is greener: Farmers' experiences with pastured poultry. *Renewable Agriculture and Food Systems*, 27(3), 173-179.
- Hughes, B. O., Dun, P., & McCorquodale, C. C. (1985). Shell strength of eggs from medium-bodied hybrid hens housed in cages or on range in outside pens. *British Poultry Science*, 26(1), 129-136.
- Johnsen, P. F., Vestergaard, K. S., & Nørgaard-Nielsen, G. (1998). Influence of early rearing conditions on the development of feather pecking and cannibalism in domestic fowl. *Applied Animal Behaviour Science*, 60(1), 25-41.
- Kopka M. N., Cheng, H. W., Hester, P. Y. (2003). Bone mineral density of laying hens housed in enriched versus conventional cages. *Poultry Science*, 82, 29.
- Krawczyk, J. (2009). Quality of eggs from Polish native Greenleg Partridge chicken-hens maintained in organic vs. backyard production systems. *Animal Science Paper and Report*, 27(3), 227-235.
- Leyendecker, M. (2003). Einfluss verschiedener Legehennenhaltungssysteme (konventionelle Käfige, ausgestaltete Käfige, intensive Auslauf- und Volierenhaltung) auf die Legeleistung, Eiqualität und Knochenfestigkeit von Legehennen. Ph. D. Diss. Univ. Osnabrück, Fachbereich Biologie, Osnabrück, Germany.
- Leyendecker, M., Hamann, H., Hartung, J., Kamphues, J., Neumann, U., Sürle, C., & Distl, O. (2005). Keeping laying hens in furnished cages and an aviary housing system enhances their bone stability. *British Poultry Science*, 46(5), 536-544.
- Leyendecker, M., Hamann, H., Hartung, J., Kamphues, J., Ring, C., Glünder, G., Ahlers, C., Sander, I., Neumann, U., & Distl, O. (2001a). Analyse von Genotyp-Umwelt-Interaktionen zwischen Legehennenhybriden und Haltungssystemen in der Legeleistung, Eiqualität und Knochenfestigkeit. 1. *Mitteilung: Knochenfestigkeit. Züchtungskunde*, 73, 387-398.
- Leyendecker, M., Hamann, H., Hartung, J., Kamphues, J., Ring, C., Glünder, G., Ahlers, C., Sander, I., Neumann, U., & Distl O. (2001). Analysis of genotype-environment interactions between layer lines and housing systems for performance traits, egg quality and bone strength. 2nd communication: Egg quality traits. *Züchtungskunde*, 73, 308-323.
- Mahboub, H. D. H., Müller, J., & Von Borell, E. (2004). Outdoor use, tonic immobility, heterophil/lymphocyte ratio and feather condition in free-range laying hens of different genotype. *British Poultry Science*, 45(6), 738-744.
- Matković K., Vučemilo, M., & Matković, S. (2007). Utjecaj alternativnog načina držanja nesilica na kvalitetu jaja. *Meso*, n.1
- Matsui, K., Khalil, A. M., & Takeda, K. I. (2004). Effects of perches on behavior, heart rate, body temperature and locomotor activity of caged hens. *The Journal of Poultry Science*, 41(2), 120-130.
- Matt, D., Veromann, E., & Luik, A. (2009). Effect of housing systems on biochemical composition of chicken eggs. *Agronomy Research*, 7(2), 662-667.
- Matt, D., Rembialska, E., Luik, A., Peetsmann, E., and Pehme, S. (2011). Quality of organic vs. conventional food and effects on health: Report. Estonian University of Life Sciences 2011.
- Mesías, F. J., Martínez-Carrasco, F., Martínez, J. M., & Gaspar, P. (2011). Functional and organic eggs as an alternative to conventional production: a conjoint analysis of consumers' preferences. *Journal of the Science of Food and Agriculture*, 91(3), 532-538.
- Minelli, G., Sirri, F., Folegatti, E., Meluzzi, A., & Franchini, A. (2007). Egg quality traits of laying hens reared in organic and conventional systems. *Italian Journal of Animal Science*, 6(sup1), 728-730.
- Mugnai, C., Dal Bosco, A., & Castellini, C. (2009). Effect of rearing system and season on the performance and egg characteristics of Ancona laying hens. *Italian Journal of Animal Science*, 8(2), 175-188.
- Nielsen, B. L., Thomsen, M. G., Sørensen, P., & Young, J. F. (2003). Feed and strain effects on the use of outdoor areas by broilers. *British Poultry Science*, 44(2), 161-169.
- Oberholtzer L., C. Greene and E. Lopez, (2006). Organic poultry and eggs capture high price premiums and growing share of specialty markets. Outlooks report from the Economic Research Service. United States Department of Agriculture LDP-M-150-01, December 2006.
- Oden, K., Keeling, L. J., Algers, B. (2002). The behaviour of laying hens in two types of aviary systems on twenty-five commercial farms in Sweden. *British Poultry Science*, 43, 169-181.
- Pavlovski, Z., Škrbić, Z., Lukić, M., Krnjaja, V., Bijelić, Z., Trenkovski, S. (2010). Technology of production of eggs with free discharge of special and guaranteed quality. *Biotechnology in Animal Husbandry*, 26(special issue), 55-66.
- Permin, A., & Pedersen, G. (2002). The need for a holistic view on disease problems in free-range chickens. *Network for Smallholder Poultry Development, The Royal Veterinary and Agricultural University Frederiksberg, Denmark*. Pp. 9-13.
- Pištekova, M., Hovorka, M., Večerek, V., Strakova, E. & Suchy, P. (2006). The quality comparison of eggs laid by laying hens kept in battery cages and in a deep litter system. *Czech Journal of Animal Science*, 51(7), 318-325.
- Pohle, K., & Cheng, H. W. (2009a). Comparative effects of furnished and battery cages on egg production and physiological parameters in White Leghorn hens. *Poultry Science*, 88(10), 2042-2051.

- Pohle, K., & Cheng, H. W. (2009b). Furnished cage system and hen well-being: comparative effects of furnished cages and battery cages on behavioral exhibitions in White Leghorn chickens. *Poultry Science*, 88(8), 1559-1564.
- Preisinger, R. (2000). Lohmann Tradition, Praxiserfahrung und Entwicklungsperspektiven. *Lohmann Inform.*, 3, 13-16.
- Prescott, N. B., & Wathes, C. M. (2002). Preference and motivation of laying hens to eat under different illuminances and the effect of illuminance on eating behaviour. *British Poultry Science*, 43(2), 190-195.
- Protais, J., Queguiner, S., Boscher, E., Piquet, J. C., Nagard, B., & Salvat, G. (2003). Effect of housing systems on the bacterial flora of egg shells. *British Poultry Science*, 44(5), 788-790.
- Reiter, K., & Bessei, W. (2000). The behavior of broilers depending on group size and stocking density. *Archive for Aviculture*, 64, 93-98.
- Rizzi, C., & Cassandro, M. (2009). Quality of eggs of hybrid and Italian hens reared under an organic production system. In *World Poultry Science Association, Proceedings of the 19th (XII) European Symposium on Quality of Poultry Meat, XIII European Symposium on the Quality of Eggs and Egg Products, 21-25 June 2009, Turku, Finland*: unpaginated
- Rizzi, C., & Chiericato, G. M. (2010). Organic farming production. Effect of age on the productive yield and egg quality of hens of two commercial hybrid lines and two local breeds. *Italian Journal of Animal Science*, 4 (suppl. 3), 160-162.
- Rakonjac, S., Bogosavljević-Bošković, S., Dosković, V., & Petrović, D. M. (2013, June). Mortality of laying hens and broilers under different rearing systems. In *23rd International symposium „New Technologies in Contemporary Animal Production* (pp. 19-21).
- Reiter, K., & Kutritz, B. (2001). Behaviour and leg weakness in different broiler breeds. *Archiv fur Geflugelkunde*, 65(3), 137-141.
- Roberts, J. R. (2004). Factors affecting egg internal quality and egg shell quality in laying hens. *The Journal of Poultry Science*, 41(3), 161-177.
- Schlatterer, J., & Breithaupt, D. E. (2006). Xanthophylls in commercial egg yolks: quantification and identification by HPLC and LC-(APCI) MS using a C30 phase. *Journal of Agricultural and Food Chemistry*, 54(6), 2267-2273.
- Schwarz, G., Kobe, A., & Fries, R. (1999). Microflora on egg shells from different housing systems. *Arch. Geflügelk.*, 5, 220-224.
- Shenstone, F. S. (1968). The gross composition, chemistry and physico-chemical basis of organization of the yolk and white. *Egg quality: A study of the hen's egg*. (Ed. TC Carter) Pp. 26-57.
- Senčić, Đ., & Butko, D. (2006). proizvodnja nesilica i kvalitetakokošjih jaja iz slobodnog i kaveznog sustava držanja. *Poljoprivreda*, 12(2), 1-5.
- Shimmura, T., Suzuki, T., Hirahara, S., Eguchi, Y., Uetake, K., & Tanaka, T. (2008). Pecking behaviour of laying hens in single-tiered aviaries with and without outdoor area. *British Poultry Science*, 49(4), 396-401.
- Škrbić, Z., Pavlovski, Z., Lukić, M., Vitorović, D., Petričević, V., & Stojanović, L. (2011). Changes of egg quality properties with the age of layer hens in traditional and conventional production. *Biotechnology in Animal Husbandry*, 27(3), 659-667.
- Sossidou, E. N., Dal Bosco, A., Elson, H. A., & Fontes, C. M. G. A. (2011). Pasture-based systems for poultry production: implications and perspectives. *World's Poultry Science Journal*, 67(1), 47-58.
- Suto, Z., Horn, P., & Ujvari, J. (1997). The effect of different housing systems on production and egg quality traits of brown and Leghorn type layers. *Acta Agraria Kaposvariensis*, 1(1), 29-35.
- Tauson, R. (2002). Furnished cages and aviaries: production and health. *World Poultry Sci. J.*, 58: 49-63.
- Tauson, R., & Holm, K. E. (2001). First furnished small group cages for laying hens in evaluation program on commercial farms in Sweden. In H.Oester and Wyss (eds.) *Proceeding 6th European Symposium on Poultry Welfare* Zollikofen, Switzerland. Pp. 26-32.
- Tauson, R., Wahlström, A., & Abrahamsson, P. (1999). Effect of two floor housing systems and cages on health, production, and fear response in layers. *Journal of Applied Poultry Research*, 8(2), 152-159.
- Trziszka, T., Dobrzański, Z., Oziembłowski, M., Jarmoluk, A., Krasnowska, G. (2004). An attempt to compare the quality of chicken eggs from cage system and ecological production. *Archiv fur Geflugelkunde*, 68(6), 269-274.
- Tuytens, F., Heyndrickx, M., Boeck, M. D., Moreels, A., Nuffel, A. V., Poucke, E. V., Coillie, E. V., Dongen, S. V., & Lens, L. (2005). Comparison of broiler chicken health and welfare in organic versus traditional production systems. *Animal Science Papers and Reports*, 23(Suppl. 1), 217-222.
- Väisänen, J., Håkansson, J., & Jensen, P. (2005). Social interactions in Red Junglefowl (*Gallus gallus*) and White Leghorn layers in stable groups and after re-grouping. *British Poultry Science*, 46(2), 156-168.
- Valkonen, E., Venäläinen, E., Rossow, L., & Valaja, J. (2008). Effects of dietary energy content on the performance of laying hens in furnished and conventional cages. *Poultry Science*, 87(5), 844-852.
- van de Weerd, H. A., & Day, J. E. (2009). A review of environmental enrichment for pigs housed in intensive housing systems. *Applied Animal Behaviour Science*, 116(1), 1-20.
- Van Horne, P. L. M., & Van Niekerk T. G. C. M. (1998). Volieren- und Käfighaltung im Vergleich. *Dtsch. Geflugelwirtsch. Schwieneprod.*, 6, 14-16.
- Van Overbeke, I., Duchateau, L., De Zutter, L., Albers, G., & Ducatelle, R. (2006). A comparison survey of organic and conventional broiler chickens for infectious agents affecting health and food safety. *Avian diseases*, 50(2), 196-201.
- Vits, A., Weitzenbürger, D., Hamann, H., & Distl, O. (2005). Production, egg quality, bone strength, claw length, and keel bone deformities of laying hens housed in furnished cages with different group sizes. *Poultry science*, 84(10), 1511-1519.
- Wall, H., Tauson, R., & Elwinger, K. (2002). Effect of nest design, passages, and hybrid on use of nest and production performance of layers in furnished cages. *Poultry Science*, 81(3), 333-339.
- Weeks, C. A., & Nicol, C. J. (2006). Behavioural needs, priorities and preferences of laying hens. *World's Poultry Science Journal*, 62(2), 296-307.
- Weitzenbürger, D., Vits, A., Hamann, H., & Distl, O. (2005). Effect of furnished small group housing systems and furnished cages on mortality and causes of death in two layer strains. *British Poultry Science*, 46(5), 553-559.
- Whay, H. R., Main, D. C. J., Green, L. E., Heaven, G., Howell, H., Morgan, M., Pearson, A., & Webster, A. J. F. (2007). Assessment of the behaviour and welfare of laying hens on free-range units. *Veterinary Record*, 161(4), 119-128.
- Yakubu, A., Salako, A. E., & Ige, A. O. (2007). Effects of

- Genotype and Housing System on the Laying Performance of Chickens in Different Seasons in the Semi-Humid Tropics. *International Journal of Poultry Science*, 6(6), 434-439.
- Yue, S., & Duncan, I. J. H. (2003). Frustrated nesting behaviour: relation to extra-cuticular shell calcium and bone strength in White Leghorn hens. *British Poultry Science*, 44(2), 175-181.
- Zeltner, E., & Hirt, H. (2003). Effect of artificial structuring on the use of laying hen runs in a free-range system. *British Poultry Science*, 44(4), 533-537.