

PUBLICATION 04

[Dieu donné GNONLONFOUN, Constant ADJEN, Mendinatou AGBETOU, Thierry ADOUKONOU, Jennifer MAPAGA, Gérard GOUDJINOU, Dismand HOUINATO]	[Post-stroke epilepsy within a teaching hospital in Cotonou, Benin]	[Neuroscience & Medicine]	[(8): 47-52]	[2017]
---	--	--	---------------------	---------------

Recherche scientifique

ACCES OUVERT

Journal ▼

Search Title,Keywords,Author,etc.

[Accueil](#)[Des articles](#)[Journaux](#)[Des](#)[Livres](#)[Nouvelles](#)[Sur](#)[Soumettre](#)[Accueil](#) > [Revue](#) > [Médecine & Soins de santé](#) > [NM](#)

Neuroscience et médecine

[Citations](#) [h5 - index et classement](#)

Soumission de

1 0

[Ouvrir des numéros](#)[spéciaux](#)[Abonnement NM](#)[Alerte Courriel](#)[Directive sur les questions](#)[NM Papiers les plus populaires](#)[Éthique de la publication et](#)[NM Nouvelles](#)[Questions fréquemment posées](#)[Recommander à ses pairs](#)[Recommander à la bibliothèque](#)[Contactez nous](#)
[Des articles](#) [Archiver](#) [Indexage](#) [Objectifs et portée](#) [Comité éditorial](#) [Pour les auteurs](#)
[Frais de publication](#)

Éditeur en chef

Prof. **Thomas Müller**

Hôpital St. Joseph Berlin-Weißensee, Allemagne

Comité éditorial

Dr. Craig Stephen Atwood	Université du Wisconsin, États-Unis
Prof. Ben A. Bahr	Université de Caroline du Nord, USA
Prof. Jackson C. Bittencourt	Université de Sao Paulo, Brésil
Prof. Gerson Chadi	Université de Sao Paulo, Brésil
Prof. Malay Chatterjee	Université de Jadavpur, Inde
Prof. Gamal HA El-Sokkary	Université d'Assiout, Egypte
Dr. Paula CRP Faria	Institut polytechnique de Leiria (IPL), Portugal
Prof. Claire Louise Gibson	Université de Leicester, Royaume-Uni
Prof. Deborah Ann Hall	Rush University, États-Unis
Prof. Hee Tae Kim	Université Hanyang, Corée du Sud
Dr. Athanasia Kotini	Université Démocrite de Thrace, Grèce
Prof. Nikolai E. Lazarov	Université médicale de Sofia, Bulgarie
Prof. KP Mohanakumar	Institut indien de biologie chimique, Inde
Prof. George Perry	Université du Texas, États-Unis
Prof. Matteo Antonio Russo	Université de Rome, Italie
Prof. Ayman I. Sayegh	Tuskegee University, États-Unis
Prof. John Raymond Smythies	Université de Californie, USA
Prof. Bing-Wen Soong	Université nationale Yang-Ming, Taipei chinois
Prof. Hermona Soreq	Université hébraïque de Jérusalem, Israël
Prof. Maria Giuliana Vannucchi	Université de Florence, Italie
Prof. David Vauzour	Université d'East Anglia, Royaume-Uni
Prof. Moo-Ho a gagné	Université de Hallym, Corée du Sud
Prof. Yukihiko Yasui	Université Shimane, Japon
Prof. Ken Kin Lam Yung	Université baptiste de Hong Kong, Chine

Suivez SCIRP


 Connexion au système de
soumission

Accueil

[Journaux AZ](#)[Livres](#)[Des conférences](#)[Emplois](#)[Blog](#)[Plan du site](#)

A propos de SCIRP

[Journaux par sujet](#)[Pour les auteurs](#)[Frais de publication](#)[Problèmes spéciaux](#)[Recherches d'évaluation par les pairs](#)[Nouvelles](#)

Un service

[Système de suivi des manuscrits](#)[Traduction et relecture](#)[Carrières](#)[Abonnement](#)[Questions fréquemment posées](#)[Contactez nous](#)

Politiques

[Open Access](#)[Éthique de la publication](#)[Préservation](#)[Réaction](#)[La publicité](#)[Politique de confidentialité](#)

Recherche scientifique

ACCES OUVERT

Journal

Search Title,Keywords,Author,etc.

- Accueil
- Des articles
- Journaux
- Des
- Livres
- Nouvelles
- Sur
- Soumettre

Accueil > Revues > Médecine & Soins de santé > NM

Neuroscience et médecine

Citations **h5 - Index et classement**

Soumission de

1 0

Ouvrir des numéros spéciaux

Abonnement NM

Alerte Courriel

Directive sur les questions NM Papiers les plus populaires

Éthique de la publication et

NM Nouvelles

Questions fréquemment posées

Recommander à ses pairs

Recommander à la bibliothèque

Contactez nous

Des articles Archiver Indexage Objectifs et portée Comité éditorial Pour les auteurs

Frais de publication

Éditeur en chef

Prof. **Thomas Müller** Hôpital St. Joseph Berlin-Weißensee, Allemagne

Comité éditorial

- Dr. **Craig Stephen Atwood** Université du Wisconsin, États-Unis
- Prof. **Ben A. Bahr** Université de Caroline du Nord, USA
- Prof. **Jackson C. Bittencourt** Université de Sao Paulo, Brésil
- Prof. **Gerson Chadi** Université de Sao Paulo, Brésil
- Prof. **Malay Chatterjee** Université de Jadavpur, Inde
- Prof. **Gamal HA El-Sokkary** Université d'Assiout, Egypte
- Dr. **Paula CRP Faria** Institut polytechnique de Leiria (IPL), Portugal
- Prof. **Claire Louise Gibson** Université de Leicester, Royaume-Uni
- Prof. **Deborah Ann Hall** Rush University, États-Unis
- Prof. **Hee Tae Klm** Université Hanyang, Corée du Sud
- Dr. **Athanasia Kotini** Université Démocrite de Thrace, Grèce
- Prof. **Nikolai E. Lazarov** Université médicale de Sofia, Bulgarie
- Prof. **KP Mohanakumar** Institut indien de biologie chimique, Inde
- Prof. **George Perry** Université du Texas, États-Unis
- Prof. **Matteo Antonio Russo** Université de Rome, Italie
- Prof. **Ayman I. Sayegh** Tuskegee University, États-Unis
- Prof. **John Raymond Smythies** Université de Californie, USA
- Prof. **Bing-Wen Soong** Université nationale Yang-Ming, Taipei chinois
- Prof. **Hermona Soreq** Université hébraïque de Jérusalem, Israël
- Prof. **Maria Giuliana Vannucchi** Université de Florence, Italie
- Prof. **David Vauzour** Université d'East Anglia, Royaume-Uni
- Prof. **Moo-Ho a gagné** Université de Hallym, Corée du Sud
- Prof. **Yukihiko Yasui** Université Shimane, Japon
- Prof. **Ken Kin Lam Yung** Université baptiste de Hong Kong, Chine

Suivez SCIRP



Connexion au système de soumission

Accueil

Journaux AZ

Livres

Des conférences

Emplois

Blog

Plan du site

A propos de SCIRP

Journaux par sujet

Pour les auteurs

Frais de publication

Problèmes spéciaux

Processus d'évaluation par les pairs

Nouvelles

Un service

Système de suivi des manuscrits

Traduction et relecture

Carières

Abonnement

Questions fréquemment posées

Contactez nous

Politiques

Open Access

Ethique de la publication

Préservation

Rétraction

La publicité

Politique de confidentialité

Post-Stroke Epilepsy within a Teaching Hospital in Cotonou, Benin

Dieu Donné Gnonlonfoun^{1*}, Constant Adjien¹, Mendinatou Agbetou², Thierry Adoukonou², Jennifer Mapaga¹, Gérard Goudjinou¹, Dismand Houinato¹

¹Department of Neurology, National University Hospital Centre, Cotonou, Bénin

²Department of Neurology, National University Hospital Centre, Parakou, Bénin

Email: *dieudonne.gnonlonfoun@gmail.com

How to cite this paper: Gnonlonfoun, D.D., Adjien, C., Agbetou, M., Adoukonou, T., Mapaga, J., Goudjinou, G. and Houinato, D. (2017) Post-Stroke Epilepsy within a Teaching Hospital in Cotonou, Benin. *Neuroscience & Medicine*, 8, 47-52.
<https://doi.org/10.4236/nm.2017.84007>

Received: September 1, 2017

Accepted: December 1, 2017

Published: December 4, 2017

Copyright © 2017 by authors and Scientific Research Publishing Inc. This work is licensed under the Creative Commons Attribution International License (CC BY 4.0).

<http://creativecommons.org/licenses/by/4.0/>



Open Access

Abstract

Seizures occurring after stroke are risk factors for the onset of ensuing epilepsy. However, the incidence of vascular epilepsy is low. The purpose of this study was to study vascular epilepsy features in a teaching university. It was a retrospective, descriptive and analytical study, conducted from 1st January 2015 to 31st December 2016 in Cotonou Teaching Hospital (CNHU-HKM), Benin. It involved 35 patients brought together through comprehensive sampling. Vascular epilepsy hospital attendance rate was 2%. There were simple partial seizures in 74.28% of cases while cerebral ischemia was recorded in 44.44% of patients. Neurological consultation after stroke occurrence was a protective factor against the occurrence of vascular epilepsy with significant p at 0.001.

Keywords

Post-Stroke Epilepsy, Vascular Epilepsy, Cotonou, Benin

1. Introduction

Epilepsy and stroke are two common pathologies which increase in frequency with age [1]. Epilepsy is a chronic disease with 0.5% to 1% prevalence. It affects over 150/100,000 patients beyond 80 years [2]. Stroke is the third cause of mortality and the leading source of disability particularly among individuals over 65 years. The association of these two pathologies is not haphazard. Indeed, epilepsy is one of the most common stroke comorbidities [3]. Stroke accounts for 11% of adult symptomatic epilepsies [4], and beyond the age of 60, it is the leading cause of epilepsy (33%) ahead of degenerative diseases including Alzheimer's disease (15%), brain tumors (5%) or brain trauma injury (1% to 2%) [5]. The

imaging carried out for patients over 60 years of age experiencing their first seizure reveals 21% of unknown infarction. The prevalence of epileptic seizures following stroke varies from 5% to 20% with an average of 10% [6], however, only few patients will actually develop real vascular epilepsy estimated at 2 to 4% [7]. This study was conducted in view of studying vascular epilepsy features within a teaching university in Cotonou, to take stock of the various aspects of this disease within our work environment.

2. Methods

It was a retrospective, descriptive and analytical study, conducted from 1st January 2015 to 31st December 2016 in CNHU-HKM, Cotonou, Benin. Benin is a French Speaking country in West Africa, sharing borders with Nigeria on the east. It boasts of 9 million inhabitants and an area of 112.622 km². The study population covered all patients admitted to hospital or attended Neurology during the study period. The inclusion Criteria was any patient hospitalized or followed in consultation of Neurology for a diagnosis of vascular epilepsy.

The diagnosis of stroke was based on the brutal occurrence of a neurological deficit associated with characteristic brain imaging. The diagnosis of epilepsy was based on the repetition of epileptic seizures associated with epileptic grapho-elements with electroencephalogram. The diagnosis of vascular epilepsy was based on:

- Absence of a personal or family history of epilepsy or other epileptogenic neurological disease;
- Repetition of seizures, stereotyped, unprovoked, at least three weeks (21 days) after the stroke and in connection with the stroke;
- In relation with the latter with or without EEG graphic element.

Sample size was calculated using Daniel Schwartz formula $n = Z\alpha^2pq/i^2$ with 2% as the frequency of vascular epilepsies [8] is 30. But the study finally reached on 35 patients. Data collection was conducted through a compilation form including clinical, paraclinical, scalable, and therapeutic data in addition to socio-demographic data. EPIDATA and STATA/IC 11.0 were used for data entry, quality control and analysis. Comparisons were made using Chi² and Fisher tests if expected values were below five. The study of associated factors was carried out using the logistic regression model in univariate and multivariate analysis. The multivariate analysis was conducted by introducing into the model, all variables whose p value in univariate analysis is $\leq 20\%$. Significance threshold was 5% and confidence intervals were calculated at the rate of 95%.

3. Results

On the period, we received 1703 patients. 35 had vascular epilepsy. The frequency of vascular epilepsy was 2%. The sex-ratio was 1.9. The mean age of patients was 53.49 ± 18.76 (42 - 85) years. Only 8.57% of the patients were jobless most of them were married (74.29%) and 60% lived in urban area. **Table 1** highlights

Table 1. Major socio-demographic characteristics of the study population, Cotonou 2016.

	Total	Percentage
Age		
[40 - 60 years]	17	48.57
[60 - 80 years]	14	40.00
[80 - 100 years]	4	11.43
Sex		
Male	23	66.00
Female	12	34.00
Profession		
Private sector worker	7	20.00
Laborer	7	20.00
Housewife	7	20.00
Retired	6	17.14
Unemployed Trader	3	8.57
Craftsman	2	5.71
Civil servant	2	5.71
Marital Status		
Married	26	74.29
Single	4	11.43
Widower	3	8.57
Divorced	2	5.71
Place of residence		
Urban areas	21	60
Rural areas	14	40

the demographic characteristics of the individuals included in the study.

Among all patients who provided information on their therapeutic itinerary, upon the onset of their stroke 5.70% consulted a general practitioner, 14.28% a tradipractitioner and 20% a neurologist.

Most of our patients had simple partial type of seizures (74.28%) followed by secondarily generalized seizures (14.28%). Generalized seizures were observed in 5.71% of cases. The average number of seizures recorded was 5.83 ± 3.37 (1 - 16) per month. Patients have been experiencing seizures for a median period of 21.44 ± 27 (9 - 48) months. Discontinuation of anti-epileptic treatment among the study population was the most triggering factor of seizure recorded (45.7%).

27 patients (77.1%) had cerebral CT-scan. Table 2 shows these results.

The EEG was performed in 13 cases. The abnormalities are summarized in Figure 1.

All patients were put under anticonvulsant treatment. Phenobarbital was the most used drug, followed by benzodiazepine derivative drugs respectively in proportion of 57.14% and 28.57%. A mono, dual and triple therapy was respectively used among 62.85%, 25.71% and 11.42% of patients.

Table 2. Distribution of patients according to cerebral CT anomalies.

	Effective	Percentage
Ischemia	12	44.4%
Hemorrhage	7	25.9%
Leucoaraiose	5	18.5%
Lacune	3	11.2%




Figure 1. Distribution of patients according to EEG abnormalities.

Upon univariate analysis, it appeared that no socio-demographic variable influence vascular epilepsy. The result of the multivariate analysis is that consulting the neurologist after stroke occurrence influenced vascular epilepsy and was even a protective factor. **Table 3** shows this result.

4. Discussion

The prevalence of epileptic seizures after stroke is quite low. Similarly, in 2004, Camilo noted a frequency of 2% to 4% [9]. However, it could vary from 5% to 20% according to Ferro [10]. On the one hand, this gap is explained by the differences in terms of methodology and design from one study to the other; and on the other hand, the variability in the concept of vascular epilepsy [11]. The average age of vascular epilepsy occurrence established at 53 years in our study population is similar to that observed in the literature, and it reflects the age pyramid in our country. In fact, cerebrovascular injuries account for 30% to 50% of epilepsies among elderly individuals [3]. Unlike the female predominance observed in Szaflarski series [12], we recorded no such prevalence in our population. Most of our patients were married, and this reflects their good social integration owing to the type of seizure. Partial seizures were less known by the population, and they did not entail social dislocation or stigmatization. There could be any type of partial seizures. Semiology of seizures will indeed depend on the topography of vascular injury [13]. Partial seizures including motor symptoms with or without generalization were prevalent [14]. Patients experienced seizures within a median period of 21.44 ± 27.48 months, furthermore, the peak period for developing vascular epilepsy is between the first and

Table 3. Multivariate analysis.

	Vascular epilepsy	Gross RC	Adjusted RC	p
Neurologist				
Yes	5 (9.26)	1	1	0.001*
No	6 (2.29)	1.65	0.08 [0.01 - 0.37]	
Indeterminate	2 (14.29)	9.77	1.16 [0.19 - 6.82]	

*Significant Result.

eighteenth month after stroke occurrence [15]. The average frequency of seizures estimated approximately at 6 seizures per month at the 21st month reflects the difficulties in providing care and support for these patients. The predominance of ischemic stroke found in the brain CT scan [16] is in relation with the general predominance of ischemic stroke in Benin hospital facilities [17]. Therapeutically, phenobarbital was the most used drug in the light of its financial accessibility and scarcity of new antiepileptic drugs in our work environment [18]. Upon stroke occurrence, neurological consultation was the only factor associated with the occurrence of vascular epilepsy. Furthermore, it is a protective factor. This is explained by good care and support for stroke by the neurologist, because of his specific and updated knowledge on this cerebrovascular pathology.

5. Conclusion

Clinical aspects of epileptic seizures during stroke are differently perceived according to the studies. Partial seizures are predominant and require early diagnosis as well as care and support from the neurologist as protective factor against the occurrence of vascular epilepsy.

References

- [1] Hamidou, B., Aboa-Eboulé, C., Durier, J., Jacquin, A., Lemesle-Martin, M., Giroud, M., *et al.* (2013) Prognostic Value of Early Epileptic Seizures on Mortality and Functional Disability in Acute Stroke: The Dijon Stroke Registry (1985-2010). *Journal of Neurology*, **260**, 1043-1051. <https://doi.org/10.1007/s00415-012-6756-3>
- [2] Brodie, M.J. and Kwan, P. (2005) Epilepsy in Elderly People. *BMJ*, **331**, 1317-1322. <https://doi.org/10.1136/bmj.331.7528.1317>
- [3] Lamy, C., Domigo, V., Semah, F., Arquizan, C., Trystram, D., Coste, J., *et al.* (2003) Early and Late Seizures after Cryptogenic Ischemic Stroke in Young Adults. *Neurology*, **60**, 400-404. <https://doi.org/10.1212/WNL.60.3.400>
- [4] Loiseau, J., Loiseau, P., Duché, B., Guyot, M., Dartigues, J.F. and Aublet, B. (1990) A Survey of Epileptic Disorders in Southwest France: Seizures in Elderly Patients. *Annals of Neurology*, **27**, 232-237. <https://doi.org/10.1002/ana.410270304>
- [5] Hauser, W.A., Annegers, J.F. and Kurland, L.T. (1996) Incidence of Epilepsy and Unprovoked Seizures in Rochester, Minnesota: 1935-1984. *Epilepsia*, **34**, 453-468. <https://doi.org/10.1111/j.1528-1157.1993.tb02586.x>
- [6] Leys, D. (2003) Ischemic Strokes in Young Adults. *La Revue de Médecine Interne*,

- 24, 585-593. [https://doi.org/10.1016/S0248-8663\(03\)00071-7](https://doi.org/10.1016/S0248-8663(03)00071-7)
- [7] Dupont, S. and Baulac, M. (2004) Contribution of MRI to the Exploration of Partial Refractory Epilepsy. *Revue Neurologique*, **160**, 5891-5897. [https://doi.org/10.1016/S0035-3787\(04\)71187-X](https://doi.org/10.1016/S0035-3787(04)71187-X)
- [8] Bladin, C.F., Alexandrov, A.V., Bellavance, A., Bornstein, N., Chambers, B., Coté, R., et al. (2000) Seizures after Stroke: A Prospective Multicenter Study. *Archives of Neurology*, **57**, 1617-1622. <https://doi.org/10.1001/archneur.57.11.1617>
- [9] Camilo, O. and Goldstein, L.B. (2004) Seizures and Epilepsy after Ischemic Stroke. *Stroke*, **35**, 1769-1775. <https://doi.org/10.1161/01.STR.0000130989.17100.96>
- [10] Ferro, J.M. and Pinto, F. (2004) Poststroke Epilepsy: Epidemiology, Pathophysiology and Management. *Drugs Aging*, **21**, 639-653. <https://doi.org/10.2165/00002512-200421100-00003>
- [11] Daniele, O., Mattaliano, A., Tassinari, C.A. and Natalè, E. (1989) Epileptic Seizures and Cerebrovascular Disease. *Acta Neurologica Scandinavica*, **80**, 17-22. <https://doi.org/10.1111/j.1600-0404.1989.tb03836.x>
- [12] Szaflarski, J.P., Kay, B., Gotman, J., Privitera, M.D. and Holland, S.K. (2013) The Relationship between the Localization of the Generalized Spike and Wave Discharge Generators and the Response to Valproate. *Epilepsia*, **54**, 471-480. <https://doi.org/10.1111/epi.12062>
- [13] Vercueil, L. (2007) Seizure and Epilepsy after Stroke. *Presse Medicale*, **36**, 176-181. <https://doi.org/10.1016/j.lpm.2006.11.001>
- [14] Kremer, S., Minotti, L., Thiriaux, A., Grand, S., Satre, V., Le Bas, J.-F., et al. (2003) Epilepsy and Hypothalamic Hamartoma: Look at the Hand Pallister-Hall Syndrome. *Epileptic Disorders*, **5**, 27-30.
- [15] Hildebrand, J., Van Bogaert, P., Tugendhaft, P. and Thiriaux, A. (2002) Drug Treatment of Epilepsy. *Revue Medicale De Bruxelles*, **23**, 500-503.
- [16] Gnonlonfoun, D., Adjien, C., Ossou-Nguet, P., Mapoure, Y., Houinato, D. and Avode, D. (2013) Frequency of Neurology Diseases in CNHU-HKM of ²Cotonou. *Journal De La Societe De Biologie*, **18**, 43-48.
- [17] Adoukonou, T., Kouna-Ndouongo, P., Codjia, J.-M., Covi, R., Tognon-Tchegnonsi, F., Preux, P.-M., et al. (2013) Direct Hospital Cost of Stroke in Parakou in Northern Benin. *The Pan African Medical Journal*, **16**, 121-125.
- [18] Houinato, D., Adjien, K.C., Gnonlonfoun, D., Adoukonou, T., Chabi Iranin, I. and Avode, D. (2007) Study of Epilepsy Prevalence in Dangbo (Benin). *Le Bénin Médical*, **14**, 14-17.