

# Toxicology of deoxynivalenol and its acetylated and modified forms

Delphine Payros<sup>1</sup> · Imourana Alassane-Kpembé<sup>1</sup> · Alix Pierron<sup>1,2</sup> · Nicolas Loiseau<sup>1</sup> · Philippe Pinton<sup>1</sup> · Isabelle P. Oswald<sup>1</sup> 

Received: 9 August 2016 / Accepted: 22 August 2016  
© Springer-Verlag Berlin Heidelberg 2016

**Abstract** Mycotoxins are the most frequently occurring natural contaminants in human and animal diet. Among them, deoxynivalenol (DON), produced by *Fusarium*, is one of the most prevalent and thus represents an important health risk. Recent detection methods revealed new mycotoxins and new molecules derived from the “native” mycotoxins. The main derivatives of DON are the acetylated forms produced by the fungi (3- and 15-acetyl-DON), the biologically “modified” forms produced by the plant (deoxynivalenol-3- $\beta$ -D-glucopyranoside), or after bacteria transformation (de-epoxy DON, 3-epi-DON and 3-keto-DON) as well as the chemically “modified” forms (norDON A-C and DON-sulfonates). High proportions of acetylated and modified forms of DON co-occur with DON, increasing the exposure and the health risk. DON and its acetylated and modified forms are rapidly absorbed following ingestion. At the molecular level, DON binds to the ribosome, induces a ribotoxic stress leading to the activation of MAP kinases, cellular cell-cycle arrest and apoptosis. The toxic effects of DON include emesis and anorexia, alteration of intestinal and immune functions, reduced absorption of the nutrients as well as increased susceptibility to infection and chronic diseases. In contrast to DON, very little information exists concerning the acetylated and modified forms; some can be converted back to DON, their ability to bind to the ribosome and to induce cellular effects varies

according to the toxin. Except for the acetylated forms, their toxicity and impact on human and animal health are poorly documented.

**Keywords** Deoxynivalenol · Acetylated forms · Modified forms · Toxicokinetics · Ribosome binding · Toxicity · Health effects

## Abbreviations

Acetyl-DON	Acetyl-deoxynivalenol
3-acetyl-DON	3-Acetyl deoxynivalenol
15-acetyl-DON	15-Acetyl deoxynivalenol
CAT	Catalase
CDK	Cyclin-dependent kinase
D3G	Deoxynivalenol-3- $\beta$ -D-glucopyranoside
DOM-1	De-epoxy DON
DON	Deoxynivalenol
DONS	DON sulfate
ERK1/2	Extracellular signal-regulated kinase 1 and 2
Hck	Hematopoietic cell kinase (Tyrosine protein kinase)
IEC	Intestinal epithelial cells
IPEC	Intestinal porcine epithelial cells
GLUT5	D-Fructose Transporter Glucose Transporter-5
Ig	Immunoglobulin
iNOS	Nitric oxide synthase
JNK	c-Jun N-terminal kinase
MAP Kinase	Mitogen-activated protein kinase
NF- $\kappa$ B	Nuclear factor- $\kappa$ B
NO	Nitric oxide
PKR	Protein kinase R
ROS	Reactive oxygen species
RNS	Reactive nitrogen species

✉ Isabelle P. Oswald  
isabelle.oswald@toulouse.inra.fr

<sup>1</sup> Toxalim (Research center in Food Toxicology), Université de Toulouse, INRA, ENVT, INP-Purpan, UPS, Toulouse, France

<sup>2</sup> BIOMIN Research Center, Technopark 1, 3430 Tulln, Austria

SGLT1	D-Glucose/D-galactose sodium-dependent transporter
SOD	Superoxidase dismutase
TEER	Trans-epithelial electrical resistance

## Introduction

Mycotoxins are secondary metabolites produced by fungi that can contaminate a large variety of dietary material consumed by humans and animals. The food contamination by these natural toxins is a worldwide problem and represents an important risk factor for human and animal health, as up to 70 % of the world's crop production may be contaminated (Streit et al. 2013). The main mycotoxin-producing fungi are *Aspergillus*, *Fusarium* and *Penicillium*, producing aflatoxins, zearalenone, trichothecenes especially deoxynivalenol (DON), fumonisins, ochratoxins and patulin.

Among the mycotoxins produced by *Fusarium*, the worldwide incidence of trichothecenes contamination and more precisely the incidence of DON has increased because of climate changes, use of no-till farming to prevent soil erosion, non-optimal crop rotation and inadequate fungicide application (Larsen et al. 2004; Rodriguez-Carrasco et al. 2014). DON is resistant to milling, processing and heating, this toxin is thus present in the food chain (Sugita-Konishi et al. 2006). Studies of the occurrence of DON in food and feed matrices reported that DON is the most prevalent food-associated mycotoxin. In the USA, 73 and 92 % of wheat and corn are contaminated with DON, respectively (Canady et al. 2001). More than 40 countries, mainly in North America and Europe, have introduced regulatory or guideline levels for DON in food and feed. The Food and Drug Administration (FDA) and the European Commission (EC) decided to limit DON in cereals and derived products (flour, pasta or bran) depending on the exposed population (children or adults) or depending on the exposed species (farm and domestic animals) (European Commission 2007; FDA 2010).

The development of new analytical methods highlighted new fungal secondary metabolites and modified forms of well-known mycotoxins. The term of “masked mycotoxins” was introduced in 1990 by Gareis to describe a zearalenone glucoside not detected during routine analysis, but hydrolyzed during digestion (Gareis et al. 1990). Indeed, biological (by plant, fungus or animal), and chemical changes (induced during thermal food processing) can modify the structure of mycotoxins, which make them undetectable by conventional analytical techniques. The term of “masked mycotoxin” was used ambiguously, and recently, a precise terminology for the various forms

of mycotoxins has been proposed (Berthiller et al. 2013; Rychlik et al. 2014). The term “modified” mycotoxins includes the different forms of mycotoxins while the term “masked mycotoxins” only applies to toxins conjugated by plants. Several hierarchic levels of modification are described: (1) the highest level differentiates the native (free and unmodified forms) mycotoxins from the “modified” forms and matrix-associated mycotoxins; (2) the next level for example differentiates the “modified” forms further in detail: the biologically modified forms from the chemically modified forms; (3) the third lower level differentiates more precisely the single modified forms, discriminating the functionalized from conjugated forms in biologically modified forms and thermally from nonthermally forms in chemically modified forms; (Rychlik et al. 2014) (Fig. 1).

As far as DON is concerned, the main derivatives of DON are described in Fig. 1. They encompass, the acetylated forms produced by the fungi (3-acetyl-DON and 15-acetyl-DON); the biologically “modified” forms produced by the plant (deoxynivalenol-3- $\beta$ -D-glucopyranoside (D3G), or after bacteria transformation (de-epoxy DON (DOM-1), 3-epi-DON and 3-keto-DON) as well as chemically “modified” forms (norDON A-C and DON-sulfonates DONSSs).

This review summarizes the current knowledge regarding DON, its acetylated and modified forms especially (i) their toxicokinetics, (ii) their molecular effects, (iii) their cellular effects, (iv) their toxicity and (v) their impact on human and animal health.


## DON

### Toxicokinetics of DON

Toxicokinetics is frequently described by the processes of absorption (A), distribution (D), metabolism (M) and elimination (E), collectively termed as ADME.

### Absorption

Absorption of DON in animals depends on several parameters including the species, the age and even the sex. Although the appearance of DON in blood following oral ingestion is rapid, within 15–30 min in most mammal species, the extent of absorption varies largely from 7 % in ruminants (sheep and cow) to 25 % in rat, and up to 89 % after chronic exposure in pig (Lake et al. 1987; Prelusky et al. 1988; Goyarts and Danicke 2006). The oral bioavailability of DON in poultry is particularly poor, and the low absorption rate may explain the low sensitivity of poultry to



**Fig. 1** Deoxynivalenol and its acetylated and modified forms covered in this review. Three hierarchic levels of modification have been described: 1st level discriminates the highest level and differentiates the native mycotoxins (DON, 3-acetyl-DON and 15-acetyl-DON) from the “modified” forms. 2nd level differentiates the biologically modified forms from the chemically modified forms. Finally, the 3rd

level differentiates the functionalized-conjugated forms (D3G) from the “differently modified” forms (DOM-1, 3-epi-DON and 3-keto-DON) within the biologically modified forms and thermally (norDON A-C) from non-thermally (DONSs) formed forms within the chemically modified forms

DON (Prelusky et al. 1986; Osselaere et al. 2013a). Plasma and tissue DON concentrations in weanling mice were approximately twice that of adult mice given same doses of the toxin, indicating that DON may be more efficiently absorbed in young animals (Pestka and Amuzie 2008). Likewise, the comparison of DON bioavailability in male and female rats suggested that this mycotoxin may have higher absorption in female compared to male (Wan et al. 2013).

### Distribution

Studies using radio-labeled DON have shown that DON and radioactively labeled metabolites are rapidly and transiently distributed to all tissues (plasma, muscle, abdominal fat, stomach, intestines, liver, kidney, heart, brain, lung, skin, spleen, testes, ovary and adrenals) (Prelusky et al. 1988; Rotter et al. 1991). The transient tissue distribution is probably linked with the rapid elimination of DON. DON passage across porcine placenta has also been shown (Goyarts et al. 2007; Danicke et al. 2007).

### Metabolism

Both host tissue metabolism and microbiota transformations of DON are documented. Indeed, bacteria colonizing the intestinal tract can be considered as an additional organ system playing a significant role in the metabolism of orally ingested xenobiotics.

**Metabolic fate of DON in host tissue** Though a P450 catalytic system mediating the DON hydroxylation to 16-OH-DON has been identified in *Sphingomonas* bacteria (Ito et al. 2013), there is still no report of phase I metabolism of DON in human and animals, suggesting that DON unlike some other mycotoxins may not be bio-activated to more toxic compounds, nor oxidized to a less toxic compounds by the phase I metabolism.

By contrast, various pathways for phase II metabolism (conjugations to glucuronic acid, sulfate or sulfonate) have been reported for DON in animals and human. The glucuronidation appears to be the major metabolic pathway to reduce DON toxicity. Depending on the species, this glucuronidation involves different sites on the DON molecule

(Maul et al. 2012; Uhlig et al. 2016). The glucuronidation patterns of DON in carp, trout, porcine, bovine and rat differ from human, and none of these animal models is suitable for modeling DON metabolism in human (Nagl et al. 2012; Sarkanj et al. 2013).

Sulfonation in rat and both sulfonation and sulfation pathways in chicken have also recently been reported; the new compounds were tentatively identified as 10-deoxynivalenol-sulfonate and deoxynivalenol-3 $\alpha$ -sulfate (Wan et al. 2013; Schwartz-Zimmermann et al. 2014a).

Further DON-biotransformation products including two DON-glutathione conjugates and their processing products DON-S-cysteine and DON-S-cysteinyl-glycine have been recently revealed in wheat (Kluger et al. 2013). The significant inductions of GST  $\pi$  and  $\alpha$  subfamilies observed in livers of mice sub-chronically exposed to DON suggest that in animals also, the mycotoxin may be substrate for glutathione S-transferases for phase II metabolism (Gouze et al. 2006). However, another hypothesis may be that the increase in glutathione S-transferases activity results from the DON-induced oxidative stress (Wu et al. 2015).

**Microbial transformation of DON in the digestive tract** In vitro cultures of ruminal and intestinal bacteria from several species including cows, fish, poultry, rats and pigs can reduce the epoxide group of DON, generating a 9,12-diene form, the de-epoxy-DON (DOM-1) (Karlovsky 2011). Since the epoxide ring is the primary determinant of DON toxicity, its removal is considered a detoxification process (Sundstol Eriksen and Pettersson 2004). Microbial transformation of DON into DOM-1 has been observed in vivo in several animal species as well as in human (Danicke et al. 2004; Turner et al. 2010). Of note, the efficiency of the depoxidation ability may be ascribed to a preliminary adaptation in link with previous exposure to trichothecenes (Hedman and Pettersson 1997; Yu et al. 2010).

### Excretion

DON and its metabolites can be found in feces and urine, and to a very low extent, in expired air (Prelusky et al. 1986; Lake et al. 1987). Most animal species demonstrate a rapid clearance of DON. A plasma elimination half-time of 3.9 h was reported after intravenous administration of DON to pigs, while in male sheep this half-life of elimination from plasma was 100–125 min (Prelusky et al. 1985; Rotter et al. 1991). The toxin clearance from plasma of mice orally exposed to DON followed a two-compartment kinetics with an initial rapid disappearance ( $t_{1/2} = 0.36$  h) and a slower terminal elimination ( $t_{1/2} = 7.62$ ) (Azcona-Olivera et al. 1995). The carry-over rates expressing the ratio between the excretion of DON in the edible tissues of food animals as well as in milk and eggs and DON intake

are globally negligible, resulting in limited risk for the consumer regarding DON transfer from feed to food (Danicke and Brezina 2013).

### Molecular effects of DON

#### *Ribotoxic stress*

**Binding to ribosomes** DON targets ribosome and binds to its 60S subunit. A recent X-ray crystallography study has determined the high-resolution structures of 80S ribosomes from *Saccharomyces cerevisiae* in complexes with DON (Garreau de Loubresse et al. 2014). The 3-hydroxyl group of DON is associated with a magnesium atom and stabilized by other nucleotides in the A-site of the 60S subunit of the ribosome (Pierron et al. 2016b). Pierron et al., also found that DON is able to form three hydrogen bonds with nucleotide bases of the A-site of the peptidyl transferase center of the ribosome, one in C12, C15 and C3 allowing a high stability of the molecule in the A-site (Pierron et al. 2016c) (Fig. 3). The binding of DON to ribosome induces a process known as “ribotoxic stress response” and interferes the elongation step of protein (Pestka 2010).

**Ribotoxic signal transduction** Ribotoxic stress response involves damage to the 3' end of the large 28S ribosomal RNA molecule, which functions in aminoacyl-tRNA binding, peptidyl transferase activity and ribosomal translocation. Besides translation inhibition as a consequence of the ribotoxic stress response, DON activates mitogen-activated protein kinases (MAP Kinases). The activation of protein kinase R (PKR) and tyrosine protein kinase (Hck) by DON is the first initiating event before MAP kinase activation (Zhou et al. 2014). Indeed, DON strongly activates the double-stranded RNA-activated PKR which is a critical upstream sensor and transducer of ribotoxic stress response. After its activation PKR triggers several pathways: (1) it phosphorylates eukaryotic initiation factor 2 $\alpha$  thereby to inhibiting translation; (2) it activates a wide range of factors including the signal transducer and activator of transcription, interferon regulatory factor 1 that plays central roles in gene expression; (3) it activates several proteins as p53, Jun N-terminal kinase (JNK), p38 and nuclear factor- $\kappa$ B (NF- $\kappa$ B) involved in cell growth, tumor suppression and apoptosis (Zhou et al. 2014). Following DON interaction, the ribosome might function as signal transducer and/or scaffold leading to impact transcriptional and post-transcriptional regulation of stress- and immune-related genes (Pestka 2008).

The demonstration that DON activates MAP Kinases in vitro and in vivo suggests that the ribotoxic stress response mediates DON toxicity (Pestka 2007). Several MAP Kinase pathways are activated by DON, including

p38, JNK and extracellular signal-regulated kinase 1 and 2 (ERK1/2). MAP Kinases modulate physiological processes including cell growth, differentiation and apoptosis and are critical for signal transduction in the immune response (Cobb 1999; Yang et al. 2000). Moreover, the activation of p38 and ERK affects cytokine production and contributes to the up-regulation of TNF $\alpha$  gene expression (Chung et al. 2003) and high expression of the cyclooxygenase-2 gene induced by DON in macrophages (Moon and Pestka 2002). DON induces the expression of IL-6 gene in a PKR dependent pathway. However, JNK pathway does not mediate the increase in TNF $\alpha$  and cyclooxygenase-2 gene expression but plays an important role in increasing mRNA stability.

DON exposure can also inhibit cell survival pathway (ERK/AKT/Bad) and activate cell apoptotic pathway (p38/Mitochondria/Bax/caspase-3) in Raw264.7 macrophages and induce ribosomal RNA cleavage (He et al. 2012).

### *Oxidative stress*

Oxidative stress may occur either due to the overproduction of reactive oxygen species (ROS) such as superoxide anion, perhydroxyl radical and hydroxyl radical and by reactive nitrogen species (RNS) or due to a decrease in cellular antioxidant levels with a reduced activity of the antioxidant enzymes, superoxidase dismutase (SOD), catalase (CAT) and glutathione peroxidase. An increased production of ROS leads to DNA damage, protein oxidation and lipid peroxidation. DON induces production of ROS in several cell types leading to cellular and nuclear damages as apoptosis and DNA damage (Mishra et al. 2014). For example, exposure of human colon carcinoma cells (HT29) to DON for 24 h induced ROS and RNS generation and altered the antioxidant status, with a decrease in glutathione content leading to the induction of cox-2 and the phosphorylation of NF- $\kappa$ B and apoptosis (Krishnaswamy et al. 2010). Furthermore, DON potentiates pro-oxidative effects in the non-tumorigenic intestinal epithelial cell line (IEC-6), increasing the release of ROS and the production of superoxide by mitochondria. Mechanistic analyses showed that the oxidative stress was mediated by NADPH oxidase, calcium homeostasis alteration, nitric oxide synthase (iNOS) expression and the activation of NF- $\kappa$ B and nuclear factor (erythroid-derived 2)-like 2 (Nrf2) pathways (Del Regno et al. 2015). Moreover, in Caco-2 cells, DON dose-dependently up-regulated the gene expression of iNOS but failed to increase the iNOS protein. Low concentrations of DON stimulate the protein degradation pathway by the proteasome. Inhibiting the production of iNOS by epithelial cells at lower doses, DON potentially increased susceptibility of animal to intestinal infection (Graziani et al. 2015).

In vivo, a significant up-regulation of hypoxia-inducible factor 1, subunit alpha and heme oxygenase, two sensitive

markers of oxidative stress, has been observed in the gut of chickens fed with DON-contaminated diet (Osselaere et al. 2013b). Piglets exposed to DON-contaminated feed (4 mg kg<sup>-1</sup>, 5 weeks) demonstrated a significant induction of oxidative stress with an increased concentration of CAT, hydrogen peroxide, nitric oxide (NO) and malondialdehyde. A decrease in the total antioxidant capacity was also observed in the same animals (Wu et al. 2014a, b). DON was also found to down-regulate CPT1A mRNA, a key regulatory enzyme of beta-oxidation, required for transport of long chain fatty acids into mitochondria. The metallothioneins, MT1A, MT1M and MT2B, markers of oxidative stress were increased in presence of DON (Pierron et al. 2016c).

### *Lipid peroxidation*

Lipid peroxidation induces many changes in the cell membrane integrity. In vitro, several studies report that DON treatment increased significantly the production of malondialdehyde in different cell lines (Caco-2, Hek-293 and HepG2) in a time- and dose-dependent manner (Zhang et al. 2009; Dinu et al. 2011). DON treatment also decreased the level of the glutathione S transferase (Zhang et al. 2009).

In vivo, the effect of DON exposure on lipid peroxidation is more controversial. DON exposure induces lipid peroxidation with an increase in the production of malondialdehyde, free radical-mediated lipid peroxidation preferentially in the liver (Rizzo et al. 1994; Frankic et al. 2006). However, in feeding trial, treatment with DON did not significantly alter the malondialdehyde content in hepatopancreas of common carp juveniles (Pelyhe et al. 2016). In chicken fed with DON-contaminated diet, the presence of DNA damage in their lymphocytes was observed without increase in the concentration of mitochondrial thiobarbituric acid substance (TBARS), an indicator of lipid peroxidation (Awad et al. 2012). To conclude, more investigations are needed to determine the role of DON on the lipid peroxidation.

### **Cellular effects of DON**

As reported in several reviews, DON induces cell-cycle arrest and apoptosis in vitro and in vivo (Pestka 2010; Mishra et al. 2014; Wu et al. 2014b).

#### *Apoptosis and cell death*

The pro-apoptotic effect of DON has been characterized in hematopoietic and intestinal epithelial cell lines, as well as in macrophages, monocytes and hepatocytes.

MAP Kinase signaling precedes and correlates with apoptosis (Yang et al. 2000). Several studies report that

DON-induced apoptosis is mediated by the activation of PKR and Hck and the subsequent activation of caspase-3 dependent as well as the phosphorylation of MAP kinase p38, ERK and JNK (He et al. 2012). DON reduces intestinal cell proliferation in several species including human and pig (Pinton et al. 2012; Pierron et al. 2016b). Proliferating intestinal human Caco-2 cells are more sensitive than differentiated cells to DON; the greater sensitivity of dividing cells might be due to the capacity of DON to inhibit protein synthesis and consequently decrease cell survival (Bony et al. 2006).

After DON exposure, an increased apoptosis has been observed in pig jejunal tissue in correlation with the phosphorylation of MAP kinase ERK and p38 (Pinton et al. 2012; Lucioli et al. 2013). In this species, a decrease in the ratio proliferation/apoptosis has been observed in jejuna villi together with an increased number of apoptotic cells in the lamina propria (Wu et al. 2014a; Cheat et al. 2015).

Other studies demonstrated that apoptosis caused by DON occurred in a ROS-dependent manner. An over-expression of NF- $\kappa$ B p65 and cyclooxygenase 2 genes has been reported in human HT-29 cell line exposed to DON (Krishnaswamy et al. 2010; Kalaiselvi et al. 2013). It has also been suggested that DON-induced apoptosis is caused by mitochondrial dysfunction and release of cytochrome c into the cytoplasm (Ma et al. 2012). An up-regulation in the apoptosis-related gene AIFM1 (apoptosis inducing factor mitochondrion associated 1) has been reported in DF-1 cells exposed with DON (Li et al. 2014b).

#### Cell-cycle effects

In addition to induce apoptosis, DON also induces cell-cycle arrest. In the rat intestinal epithelial cell line IEC-6, DON produces a cell-cycle arrest primarily in the G0/G1 phase. By contrast, in human epithelial cell lines, G2/M phase arrest was observed. It was associated with an increase in p21 levels, a cyclin-dependent kinase (CDK) inhibitor known mainly to inhibit cyclin-CDK2 complex and negatively regulate cell-cycle progression (Yang et al. 2008). At lower doses (0.1–2.5  $\mu$ M), DON exerts a significant remodeling of the actin cytoskeleton which is associated with a decrease in cell motility (Bianco et al. 2012).

### Toxicity of DON

#### Intestinal toxicity

The intestinal epithelium serves as a selective barrier enabling the transport of various nutrients and ions while not allowing the uptake of luminal bacteria and toxins. The intestine has been identified as one of the first targeted tissue for the action of DON, and the impacts of DON on this

organ have received an increasing interest in the last years (Pinton and Oswald 2014; Ghareeb et al. 2015; Akbari et al. 2016).

*DON induces intestinal lesions* Histology is a sensitive end point to assess the deleterious effects of DON on the intestine. In several species including rodents, pigs and chicken, DON induces significant histological changes in the intestine. The intestinal lesions reveal mainly defaults in morphology of enterocytes, a decrease in the length of villi, an increase in dead cells in the lumen and at the top of villi, edema and fusion of villi (Pinton and Oswald 2014; Ghareeb et al. 2015).

The intestinal lesions were associated with shorter and thinner villi (Kolf-Clauw et al. 2009; Pinton and Oswald 2014). The decreased villi height suggests an effect of DON in the balance between epithelial cell proliferation and apoptosis although none difference in crypt depth was observed.

The effects of DON on the impairment of the intestinal morphology participate to alteration of absorption and nutrient intake by intestinal epithelial cells (IECs) and more precisely enterocytes. Consequently, this can affect energy and nutrient uptake and adversely lead to growth deficiency in animals (Yunus et al. 2012) (Fig. 4).

*DON affects intestinal cell proliferation and differentiation* To maintain the integrity and a functional intestinal barrier, the intestinal epithelium needs to keep a constant state of regeneration, established by at least two mechanisms: epithelial cell proliferation/apoptosis and maturation/differentiation (van der Flier and Clevers 2009). DON impacts the capacity of IECs to proliferate and potentially, the renewal and repair of intestinal epithelium. DON reduced the activation of proteins implicated in the mTOR (mammalian target of rapamycin) signaling pathway, a key mechanism in the regulation of IECs proliferation and renewal. This effect traduces a reduction in IECs proliferation in crypts and so a decrease in intestinal epithelial cell renewal (Wu et al. 2014a).

The effects of DON on cell proliferation and apoptosis have been reported in IECs (see part *apoptosis and cell death*) (Pinton and Oswald 2014; Ghareeb et al. 2015). Sub-toxic doses of DON (lower than 1  $\mu$ M) decreased mucin (MUC) production in human goblet cells HT29-16E as indicated by the specific reduction in the mucin gene expression. At the molecular level, this effect relied on the inhibition of resistin-like molecule  $\beta$  (RELM $\beta$ ) gene expression, a positive regulator of MUC gene transcription (Pinton et al. 2015). A reduction in the number of calciform cells in jejunum and ileum and an increase in IECs apoptosis were observed in piglets fed a lower DON-contaminated feed (Bracarense et al. 2012). Recently, DON

exposure in chickens altered the mucus layer by suppressing MUC2 gene expression in the duodenum and by modifying the mucin monosaccharide composition in these animals (Antonissen et al. 2015).

**DON alters intestinal barrier function** Intestinal epithelium forms a strong barrier between the organism and the external environment through the development of tight junctions between epithelial cells. The trans-epithelial electrical resistance (TEER) can be considered as a good indicator of the degree of integrity of the intestinal barrier. Determination of the paracellular (small molecules < 4KDa) and transcellular (larger molecules > 4KDa) permeability reports, respectively, the flux of molecules between IECs or through the enterocytes.

The effects of DON on the intestinal barrier dysfunction have been well characterized (Fig. 4). A reduction in TEER was inversely associated with an increase in the paracellular passage of FITC-dextran in the intestine of rodent, chickens, pigs and rodents exposed to DON (Pinton and Oswald 2014; Akbari et al. 2014; Ghareeb et al. 2015). The alteration of the intestinal barrier observed in response to DON induces higher trans-epithelial passage of commensal and pathogenic bacteria as described for pathogenic *Escherichia coli* and *Salmonella* across the porcine intestinal epithelial cells (Pinton et al. 2009; Vandenbroucke et al. 2011).

The increase in intestinal permeability was mediated by MAP Kinases and associated with a specific reduction in the expression of tight-junction proteins especially claudins (Diesing et al. 2011; Pinton and Oswald 2014; Akbari et al. 2014). Interestingly, the decreased expression of junction proteins was accompanied by an up-regulation of their mRNAs levels (Osselaere et al. 2013b; Akbari et al. 2014).

**Effects of DON on gut microbiota** The effect of DON on the microbiota composition and diversity has been poorly evaluated. Studies report that sub-chronic exposure of animals to DON induces a “transient dysbiosis” and influences the intestinal microbiota diversity in the colon content decreasing the amounts of *Clostridium perfringens* without change in the amounts of *Enterobacteria* (Saint-Cyr et al. 2013; Piotrowska et al. 2014). In pigs, a moderate effect on cultivable bacteria has been observed and showed a decrease in the richness index (Wache et al. 2009b). In rats, no effect on the composition and diversity of the microbiota was observed (Payros et al. 2016).

DON also modulates the bacterial expression of two regulators of *Salmonella* pathogenicity island *hilA* and *ssrA* (Vandenbroucke et al. 2009). Additional studies are required to conclude on the effect of DON on the microbiota.

### Immunotoxicity

Once the DON has crossed the intestinal barrier, the second target to this mycotoxin is the immune system. Depending on the dose and the duration of the exposure DON modulates immune functions in multiple ways. At high concentrations, DON induces immune cells apoptosis and strong immune-suppression. At low concentrations, DON can potentiate immune cells proliferation, production of cytokines or disrupt normal immune responses, increasing the risk to develop chronic immune diseases or the susceptibility to infections (Fig. 5).

**DON exacerbates immune intestinal inflammation** Cytokines are small peptides produced by immune cells (lymphocytes, macrophages, dendritic cells...) but also by intestinal epithelial cells (IECs). As described in intestinal tissues, DON modulates the production of these mediators involved in the regulation of the immune and inflammatory response. On human IECs (Caco2 cells), DON possesses a direct pro-inflammatory effect increasing the secretion of IL8 after the activation of MAP kinase and PKR activation leading to a NFκB-dependent transcription of the IL8 gene (Maresca et al. 2008). In porcine IPEC-1 cells, DON induced a pro-inflammatory response with a significant increase in the expression of IL8, IL1β, IL1α and TNFα gene (Cano et al. 2013) (Fig. 5).

In jejunal explants, DON increased the expression of inflammatory genes (IL1β, IL1α, IL8, IL17A, IL22, TNFα) as well as genes involved in the differentiation of Th17 cells (STAT3, IL17A, ...) while decreasing the ones involved in the T cells regulatory pathway (FoxP3, RALDH1) (Cano et al. 2013; Pierron et al. 2016b, c). Similarly, in mice, DON increased the expression of mRNA encoding TGFβ and IFNγ in the small intestine (Azcona-Olivera et al. 1995).

**DON increases Immunoglobulin (Ig) secretion in the systemic circulation** A dramatic increase in the level of serum IgA as well as an elevation of serum IgA-immune complex was observed in mice or farm animal fed with a DON-contaminated diet (Pestka 2003; Grenier et al. 2011). Moreover, a concomitant increase in IgE and a decrease in IgG and IgM levels were also reported (Pestka et al. 2004; Grenier et al. 2011). The ability of DON to increase production of IgA was due to the expansion of polyclonal IgA-secreting cells in Peyer’s patches and spleen in mice (Bondy and Pestka 1991). The increased production of IgA may be related to the capacity of macrophages and T cells to secrete helper cytokines IL-2, IL-5 and IL-6 (Pestka and Smolinski 2005) (Fig. 5).

**DON modulates the systemic immune response** DON also modulates the number of immune cells numbers and their

functions in primary and secondary lymphoid organs such as spleen and lymph nodes. DON treatment induces cell apoptosis in mouse thymic epithelial cell line MTEC1 contributing to exacerbate the default of immune responses (Li et al. 2014a). In vitro, DON inhibits proliferation of B- and T cell subsets stimulated by different mitogens but also increases the production of pro-inflammatory cytokines in Jurkat T cells and primary human lymphocytes (Forsell and Pestka 1985; Pestka et al. 2005; Severino et al. 2006). DON was also found to decrease the proliferation of human and porcine peripheral blood mononuclear cells and murine splenocytes in a dose-dependent manner (Taranu et al. 2010). An increase in apoptosis and lesions in spleen and mesenteric lymph nodes has been reported in rodents exposed to DON, indicating that ingestion of DON induces immunosuppression in these organs (Bracarense et al. 2016). DON impacts on the expression of cell surface markers altering macrophage activation by reducing the expression of CD54, CD14, CD119 and HLA-DP/DQ/DR molecules limiting the capacity of these macrophages to play their role as antigen-presenting cells (APCs) (Wache et al. 2009a). DON induces apoptosis on human immature dendritic cells and inhibits the dendritic cell maturation down-regulating the gene expression of CD86, HLA-DR and CCR7, markers of dendritic cells maturation (Hymery et al. 2006) (Fig. 5).

In human macrophage cell lines, DON up-regulates the production of pro-inflammatory cytokines, and co-stimulation of macrophage with LPS potentiates TNF $\alpha$  and IL8 production in DON-exposed macrophages (Sugita-Konishi and Pestka 2001). DON suppresses Th1 immune response and IFN $\gamma$  production in mice and in the same time exacerbates Th2 cytokine gene expression, enhancing an increase in the expression of IL4, IL6 and IL10. Alterations of the proliferation of lymphocytes and the secretion of particular cytokines such as IL22A, IL4, IL2 and IL23 could explain the imbalance between Th1/Th2/Th17 immune responses. Moreover, studies suggest that DON interferes with the regulatory T cells decreasing the transcription factors FoxP3 and STAT3 (Cano et al. 2013) (Fig. 5).

#### *Hematotoxicity and myelotoxicity*

Haematotoxicity and myelotoxicity are generally defined as toxic effects in blood and bone marrow cell lineages. In rodents and farm animals, long-term ingestion of DON causes a decrease in circulating blood cells frequently associated with bone marrow dysfunction (Parent-Massin 2004). The haematopoietic progenitors, as platelets, white and red blood cells progenitors, are the main target of DON. Exposure to DON also decreases peripheral and splenic blood leukocytes associated with a decrease in erythrocytes and platelets with a greater sensitivity in mice females (Wu et al. 2009).

#### *Action on the central nervous system*

DON is able to cross the blood brain barrier few minutes after exposure and directly acts on the neurons and glial cells. Pigs are the most sensitive animal species with more than 25 % of the plasmatic DON found in the cerebro-spinal fluid, whereas only 10 and 5 % of the plasmatic DON crosses the blood brain barrier in mice and sheep, respectively. However, no studies have reported the mechanism responsible for the transport, absorption and excretion, of DON across the blood brain barrier (Behrens et al. 2015).

At very low doses ( $\leq 100$  nM), DON potentiates the inflammation caused by LPS in the glial cells, inducing iNOS and TNF $\alpha$  secretion. By contrast, at higher doses ( $>300$  nM), DON inhibits the inflammation through its cytotoxicity on microglia cells. Interestingly, DON inhibits the absorption of glutamate by astrocytes, a neurotransmitter involved in the mechanisms of obesity and food intake (Razafimanjato et al. 2011). In vivo, DON affected the activity of brain neurons which could be implicated in anorexia and emesis. DON exposure activates POMC and nesfatin-1 neurons, specific neurons implicated in central anorexigenic neuronal pathways, and present in a specific area controlling the food intake and vomiting (Ossenkopp et al. 1994; Girardet et al. 2011a; Gaige et al. 2013). In mice, DON causes a neuro-inflammation with an increased expression of IL-1 $\beta$ , TNF- $\alpha$  and the prostaglandin PGE2 in brain. This inflammation is not sufficient to induce anorexia as anorexia is still observed in mPGES-1 knock-out mice (Girardet et al. 2011b).

#### *Carcinogenic, mutagenic and reprotoxic effects*

The effect of DON on DNA damages is poorly documented. However, a genotoxic effect of DON was shown by comet assay on intestinal cells lines, lymphocytes and hepatocytes (Bony et al. 2006; Bensassi et al. 2009; Awad and Zentek 2015). DON exacerbates apoptosis in the human intestinal cell line HT-29 in association with a fragmentation of DNA and the activation of the apoptotic molecules, p53 and caspase-3 (Bensassi et al. 2009). Long-term carcinogenic risk and genotoxicity with DON in the gastrointestinal tract have not yet been explored, and the first results are controversial (Ma and Guo 2008; Singh et al. 2015). However, a recent study suggested a synergistic role of DON associated with a specific microbiota in the appearance of DNA damage in jejunal epithelial cells in vivo on rodents (Payros et al. 2016).

According to these results, DON has been classified in group 3 by IARC (International Agency from Research on Cancer) as “not classifiable as to its carcinogenicity to humans” (IARC 1993).

The effects of DON on the fertility and the development of fetuses have been investigated in rodents and pigs. In rats, a decrease in fertility was observed upon DON exposure (50 vs 80 % of mating in DON and control groups, respectively) (Morrissey and Vesonder 1985). In male, DON exposure decreases the sperm count and increases the abnormalities in sperm with an alteration of the production of the reproductive hormones, testosterone, follicle-stimulating hormone and luteal-hormone (Sprando et al. 2005).

After oral force-feeding with DON, complete resorptions, a decrease in the number of live fetuses and an increase in newborns presenting skeletal malformations were observed in mice (Khera et al. 1986). Low doses of DON are sufficient to increase the postnatal mortality without concomitant effects on fetal development and malformations in mice but not in rats, suggesting that rat is more resistant (Khera et al. 1984). In utero, DON exposure induces fetal abnormalities at 2.5 and 5 mg/kg bw/day in rats (Collins et al. 2006). In pigs, DON exposure during gestation has no adverse effects on offspring number, survival or malformations (Eriksen and Alexander 1998) although the exposure of pig oocytes to DON causes aneuploidy and abnormal embryo development in vitro (Malekinejad et al. 2007).

## Effects of DON on human and animal health

### *Emesis, anorexia and growth effects*

Acute exposure to DON induces abdominal pain, increased salivation, diarrhea and emesis in human and several animal species. Originally called “vomitoxin,” DON has a critical emesis effect in human, pigs, dogs and mink but not in mice and rats, animals unable to vomit. In rodents, abnormal feed intake is considered as indicative of anorexia.

Exposure of farm animals to diet contaminated with low-dose of DON results in a reduced feed intake and a reduced weight gain. Anorexia induced by DON can be explained by various mechanisms, including an effect on growth and satiety hormones, immune response, neuro-endocrine pathway or a central neuronal signaling. As described above (part *action on the central nervous system*), through the action of DON on the central nervous system, these changes contribute in part to impaired food intake and weight gain. The impact of DON on the intestinal immune response can indirectly affect feed consumption through the growth hormone axis. Pro-inflammatory cytokines induced by DON increase expression of suppressors of cytokine signaling which impaired growth hormone signaling (Amuzie et al. 2009). Produced by entero-chromaffin cells in the gastro-intestinal tract, neuro-endocrine factors, as serotonin acts as a paracrine fraction on the enteric nervous system and can impact the secretion of

both anorexigenic orexigenic hormones. In turkeys, the consumption of DON increases serotonin level in the brain which correlates with a decreased body weight gain (Girish et al. 2008). An adverse effect of DON on gastric emptying through a peripheral action on serotonin-3 receptors has also been described in rodents (Fioramonti et al. 1993).

Recent advances report that reduced feed intake and feed refusal might be partly due to effect of DON on the release of satiety hormones as peptide YY and cholecystokinin, critical mediators related to DON-induced anorexia and growth suppression (Flannery et al. 2012). A link between this effect of DON and the activation of receptors implicated in growth control, calcium-sensing receptor and the transient receptor potential ankyrin-1, has been shown in entero-endocrine cells. Treatment with antagonists of these receptors inhibits both DON-induced food refusal and the increase in plasma peptide YY and cholecystokinin suggesting that the activation of these two receptors contributes to food refusal induced by DON (Wu et al. 2016).

### *Effects on intestinal absorption of nutrients*

The selective permeability of the intestine for dietary nutrients, electrolytes and water is mediated through a transcellular and a paracellular ways. DON affects the absorption of amino acids and sugars through (1) the inhibition of nutrients transporters and (2) the intestinal lesions which decrease the absorption surface.

Both in vitro and in vivo data indicate that DON exposure inhibits the activities of several intestinal transporters such as the D-glucose/D-galactose sodium-dependent transporter (SGLT1), the D-fructose Transporter Glucose Transporter-5 (GLUT5), the active and passive L-serine transporters (reviewed in Maresca et al. 2002; Ghareeb et al. 2015). As SGLT-1 is the main apical transporter for active glucose uptake and absorption of water in the gut (Meinild et al. 1998), its inhibition could explain the diarrhea associated with DON exposure. In chickens exposed to DON, the electrophysiological properties of the intestinal mucosa and more precisely short-circuit current were decreased after addition of D-glucose using Ussing chambers in jejunum tissue and decreased the uptake of glucose through the inhibition of the intestinal SGLT1 (Awad et al. 2008, 2014).

### *Susceptibility to infections*

The susceptibility for the host to develop bacterial, viral or parasitic infections is increased through the disruption of the intestinal barrier function and the alteration of the immune response induced by DON exposure.

DON increases susceptibility to *Salmonella* infection. Indeed, ingestion of DON-contaminated feed potentiates oral infection with *S. enteritidis* in mice (Hara-Kudo et al.

1996). In a porcine ileal loop model, exposure to low concentrations of DON exacerbates the early immune response initiated by *S. typhimurium* infection and contributes to *Salmonella* invasion and translocation (Vandenbroucke et al. 2011).

In chickens, necrotic enteritis is an infectious disease caused by the bacteria *Clostridium perfringens* (*C. perfringens*). DON acts as a predisposing factor for the development of necrotic enteritis due to its negative effect on the intestinal barrier increasing cellular debris, so supplying the necessary nutrients for the proliferation and toxin production of *C. perfringens* (Antonissen et al. 2014).

Few studies report the effect of DON on parasitic diseases and viral infection. Nevertheless, Girgis et al., showed that the combination of DON affects the Eimeria-induced immune response in chickens, reduces the efficacy of the anti-Eimeria treatment in these animals and delays intestinal recovery from Eimeria-induced lesions in adult birds (Girgis et al. 2008, 2010).

Similarly, in mice, DON negatively affects the intestinal clearance of reovirus, a non-enveloped double-stranded RNA virus that causes arthritis in poultry, potentiating the intestinal viral load and suppress the host immune response against the virus by increased its fecal shedding (Li et al. 2007). The underlying mechanism involves a suppression of the Th1 response, limiting IFN $\gamma$  gene expression with a subsequent enhancement of the Th2 response with an increased IL4 gene expression (Li et al. 2005).

In farm animals, ingestion of feed naturally contaminated with DON was found to increase the susceptibility to two economically important viral diseases. It negatively affects the specific humoral response against the porcine reproductive and respiratory syndrome virus (PRRSV) and increases the effect of this infection on weight gain, lung lesions and mortality (Pierron et al. 2016a). DON also increases the viral replication of the porcine circovirus type 2 (PCV-2) (Savard et al. 2015).

#### Vaccine responses

In pigs, chronic intoxication with DON resulted in an impairment of immune response to a vaccinal protocol with ovalbumin in the terms of alteration of the antibody response, lymphocyte proliferation and the production of cytokine in the lymph nodes (Pinton et al. 2008). Vaccination of animals with ovalbumin results in the activation of specific lymphocytes, through the priming with ovalbumin-presenting cells. Ingestion of DON results in a decrease in lymphocyte proliferation, suggesting that the cells were unable to answer upon antigenic stimulation in animals fed with DON-contaminated diet. Moreover, the level of

anti-ovalbumin IgG and anti-ovalbumin IgA was affected by DON consumption and expression of several cytokines (IL-1 $\beta$ , IL-6, IL-8 and TGF- $\beta$ ) was reduced in spleen of animals exposed to DON (Grenier et al. 2011). In chickens, dietary DON impairs the humoral immune response against viral vaccine and affects the success of virus vaccine as infectious bronchitis virus or Newcastle virus (Yunus et al. 2012; Ghareeb et al. 2016).

These studies suggest implications of DON-contaminated food consumption for human and animal health as breakdown in vaccinal immunity.

#### Chronic diseases

Because DON generates an intestinal inflammation, it has been suggested that this toxin could predispose to chronic intestinal bowel diseases, such as celiac and Crohn's diseases or ulcerative colitis (Maresca and Fantini 2010); however, this hypothesis has never been investigated.


More recently, in a mouse model, DON was found to exacerbate allergic sensitization to milk protein (Bol-Schoenmakers et al. 2016). The underlying mechanism suggests that DON modulates intestinal barrier function to exacerbate allergic response to milk protein by impairing the epithelial permeability and modulating tight-junction mRNA and proteins expression. This allows allergen translocation and the activation of pro-inflammatory immune response by increasing IL-33 levels and enhancing local innate lymphoid cells type 2 cell numbers (Bol-Schoenmakers et al. 2016).

#### Acetylated forms of DON

Besides DON, *Fusarium graminearum* and *F. culmorum* also produce acetylated forms, 3-acetyl deoxynivalenol (3-acetyl-DON) and 15-acetyl deoxynivalenol (15-acetyl-DON).

Moreover, different *Fusarium* strains can produce different patterns of mycotoxins (chemotypes). As an example, 3-acetyl-DON and 15-acetyl-DON chemotypes of *F. graminearum* produce DON together with 3-acetyl-DON, and DON together with 15-acetyl-DON, respectively, and their distribution depends on the geographical location and years of sampling (Gilbert et al. 2014).

The evaluation of both acetyl-DONs has been poorly addressed probably because they represent only a small amount of DON and, in addition, it is admitted that their de-acetylation contributes to decrease their toxicity. However, a better knowledge of their toxic characteristics in comparison with DON, is useful to better assess the risk associated with their production by fungi.



**Fig. 2** Metabolism of DON and its acetylated and modified forms in human and animals. Microbiota transformation, intestinal absorption, host-tissue metabolism pathways and excretion of DON and illustrative acetylated (15-acetyl-DON), “masked” (D3G), bacterial modified (DOM-1) and chemically modified (DONS) forms. Depending on the species, significant amounts of DON and its acetylated and modified

forms are transformed by the microbiota before uptake by the host tissue. The main phase II biotransformation pathways take place in the liver and include glucuronidation, sulfatation and sulfonation. The resulting hydrophilic forms are mainly excreted in urine, though significant parts may undergo entero-hepatic cycling, extending the organism exposure to the deleterious effects of DON

## Toxicokinetics of acetylated forms of DON

### Absorption and distribution

The absorbed fractions of orally administrated 3-acetyl-DON and 15-acetyl-DON in chickens correspond to approximately double and quadruple that of DON, respectively (Broekaert et al. 2015a). Likewise, the administration of the acetylated forms leads to a more rapid absorption, reflected by a shorter  $t_{max}$  ( $5.0 \pm 0$ ;  $9.0 \pm 6.5$  and  $32 \pm 11$  min for 3-acetyl-DON, 15-acetyl-DON and DON, respectively). This was expected since 3-acetyl-DON and 15-acetyl-DON are less polar and have a more favorable log D value for absorption by passive nonionic diffusion compared to DON (estimated log D for acetylated forms of DON and log D for native DON at physiological pH are  $-0.5$  and  $-1$ , respectively) (Fig. 2).


There are no specific data on the distribution of the acetylated forms of DON. However, because of the conversion

of the absorbed acetylated forms into DON, we can anticipate that their distribution pattern do not differ significantly from the one of DON.

### Metabolism

**Microbial transformation of DON and derivatives in the digestive tract** De-acetylation but not de-epoxidation has been demonstrated in human feces incubated under anaerobic conditions with 3-acetyl-DON (Sundstol Eriksen and Pettersson 2003; Ajandouz et al. 2016). However, it could be possible that, just like DON, the naive intestinal bacteria naturally do not possess the ability to de-epoxydate 3-acetyl-DON and that pre-exposure of the microbiota to 3-acetyl-DON induces the appearance of the bacterial transformation activity (Turner et al. 2010; Maresca 2013; Gratz et al. 2013).

Only two metabolites, DOM-1 and DON, have been reported from the microbial transformation of 3-acetyl-DON in the chicken digestive tract. By contrast,



**Fig. 3** Interaction between the Ribosome 60S subunit binding site and DON or its acetylated and modified forms. In each panel, surface of ribosome is colored in *red* and the referent binding molecule DON is colored in *white*. Each acetylated or modified form of DON is represented by a different color: 3-acetyl-DON in *orange*; 15-acetyl-DON in *green*; D3G in *yellow*; DOM-1 in *purple*; 3-epi-DON in *cyan*; and 3-keto-DON in *gray*. Hydrogen bonds between DON or

modified forms and the A-site of the peptidyl transferase center of the ribosome are indicated by a *dashed line* in the same color than the molecule. (A) Front view. (B) Top view. DOM-1, 3-epi-DON and 3-keto-DON being totally aligned with the DON structure, the picture only highlights the part of the DON structure that differs with the modified forms

15-acetyl-DON showed besides DOM-1 and DON an additional de-epoxidation product with the acetyl group still intact (de-epoxy-15-acetyl-DON) (Young et al. 2007).

De-acetylation of 3-acetyl-DON seems also required prior to de-epoxidation by the cow ruminal flora and microorganisms in pig ileal digesta or faeces (King et al. 1984; Eriksen et al. 2002). Whether microbiota from other species than chicken could de-epoxydate 15-acetyl-DON keeping its acetyl group intact has not been investigated yet.


*Metabolic fate of the acetylated derivatives of DON in host tissue* Upon oral administration of 3-acetyl-DON or 15-acetyl-DON, only DON, DON-glucuronide, DOM-1 and trace amounts of the acetylated forms are detected in pig blood (Eriksen et al. 2003; Maresca 2013; Broekaert et al. 2015b). On the other hand, extensive de-acetylation of both 3-acetyl-DON and 15-acetyl-DON was reported following

their intra-venous administration to pig and poultry (Broekaert et al. 2015a).

By contrast to pig and poultry, in rats the phase II biotransformations can directly glucuronidate the acetylated forms of DON without preliminary de-acetylation (Versilovskis and Bartkevics 2012). Rats fed 3- and 15-acetyl-DON, accumulated glucuronidated 3-acetyl-DON in their small intestine, together with DON-3GlcA; low level of 15-acetyl-DON-3-glucuronide was found in the bladder.

#### Excretion

There is a lack of data on the excretion of acetylated forms of DON in rodents and human. A higher clearance of 3-acetyl-DON compared to that of DON and 15-acetyl-DON has been reported in chickens and pigs (Broekaert et al. 2015a). In contrast, compared to DON,



**Fig. 4** Intestinal toxicity of DON and its acetylated and modified forms. Schematic illustration of effect of DON, its acetylated and modified forms on (1) intestinal lesions, (2) epithelial cells differentiation, (3) barrier function and (4) gut microbiota. Few data described

the effects of the modified forms on the immune responses. For each toxin, adverse effects are indicated in *red* (strong) or *orange*. The absence of effects of a given compound is indicated in *green*. In *black* are mentioned toxins for which no information is available

15-acetyl-DON showed a faster clearance in pigs and a slower clearance in chickens.

### Molecular and cellular effects of acetylated forms of DON


In contrast to DON, 3-acetyl-DON and 15-acetyl-DON bind the ribosome with only 2 hydrogen bonds: the first one involved the epoxy group and the second one the non-acetylated remaining hydroxyl group in position 15 and 3 for the 3-acetyl-DON and the 15-acetyl-DON, respectively (Fig. 3).

The hydrophobic acetyl group in position 15 induces additional hydrophobic interaction which may stabilize 15-acetyl-DON in the ribosome. On the opposite, the hydrophobic acetyl group in position 3 induces a VanDer Waals interaction that may stabilize the complex but also

induce some steric hindrance deepest in the A pocket of the ribosome.

The ability of 3-acetyl-DON and 15-acetyl-DON to induce ribotoxic stress response was examined through the activation of MAP Kinases in differentiated intestinal epithelial cell line, intestinal explants and in the intestine of exposed animals. In these three different models, 15-acetyl-DON activated the MAP Kinases ERK1/2, p38, and JNK at lower dose than DON and 3-acetyl-DON. As a consequence, 15-acetyl-DON displays a higher toxicity as measured by histological intestinal lesions and by impaired barrier function (Pinton et al. 2012).

3-acetyl-DON and 15-acetyl-DON were found to induce apoptosis of Jurkat T-cell at the same concentration than DON. The underlying mechanism of apoptosis has not been investigated for these two metabolites (Pestka and Smolinski 2005).



**Fig. 5** Immunotoxicity of DON and its acetylated and modified forms. **a** Effects on primary lymphoid organs, bone marrow and thymus. **b** Effects on secondary lymphoid organs, spleen and lymph nodes. **c** Effects on gut-associated lymphoid tissue. At these different levels, DON is known to induce a default in the production of functional immune cells; to induce apoptosis of immune cells and a default in immune cell response; to exacerbate the inflammatory responses in the intestinal mucosa increasing the production of

pro-inflammatory cytokines, neutrophil infiltration, over-reaction against self-antigens and microbiota, alteration of immunoglobulin production and an imbalance of T cell response. Few data described the effects of the modified forms on the immune responses. Adverse effect are in red (strong) and orange (mild). The absence of effects is colored in green. In black are mentioned toxins for which no information is available

## Toxicity of acetylated forms of DON

### Intestinal Function toxicity

Gastro-intestinal lesions induced by acetyl-DONs are similar to those observed with DON (Fig. 4). In mice given a single exposure of 100 mg/kg bw 15-acetyl-DON, severe necrosis of the crypt epithelial cells in the small and large intestine was observed. Lumen of the crypts was dilated and filled with necrotic cellular debris. Lamina propria of the villi was edematous and congested (Forsell et al. 1987). In mice gavaged with 40 mg/kg bw of 3-acetyl-DON reduction in mitotic activity and presence of necrotic cells were

observed in duodenal crypts (Kasali et al. 1985). In pig, jejunal explants short term exposed to DON, 3-acetyl-DON and 15-acetyl-DON the main observed effects were the presence of flattened and coalescent villi, the lyses of enterocytes and interstitial edema and apoptosis. DON and its modified forms could be ranked in the following order of deleterious potency 15-acetyl-DON  $\gg$  DON = 3-acetyl-DON (Pinton et al. 2012) (Fig. 4).

3-Acetyl-DON and 15-acetyl-DON differentially impact the capacity of intestinal epithelial cells to proliferate and so could affect the renewal and repair of intestinal epithelium. They were cytotoxic in a dose-dependent manner for human intestinal Caco-2 cells. However, 3-acetyl-DON was

two fold less toxic for proliferating cells than 15-acetyl-DON in both human and porcine intestinal cells (Alassane-Kpembi et al. 2013, 2015).

Transport studies were conducted on Caco-2 cells exposed to DON, 3- or 15-acetyl-DON, which reduce TEER with no cell damage. The transport rate of 15-acetyl-DON was higher compared to DON and 3-acetyl-DON in apical-to-basolateral direction, suggesting that 15-acetyl-DON was absorbed more efficiently across the monolayer. Similar values observed for apical-basolateral and basolateral-apical transport rates, suggested that the transport mechanisms are similar for DON and acetylated forms. Furthermore, 3 and 15-acetyl-DON were absorbed without deacetylation to form DON (Kadota et al. 2013). A decrease in the TEER of differentiated IPEC-1 cells and an increase in the paracellular permeability were observed after a 24-h exposure to the toxins that were ranked in the following order of toxicity: 15-acetyl-DON  $\gg$  DON > 3-acetyl-DON (Pinton et al. 2012). Similar data were obtained using human Caco-2 cells (Kadota et al. 2013) (Fig. 4).

In isolated chicken jejunum fragments mounted in Ussing chamber, the exposure to 15-acetyl-DON led to a decrease in the short-circuit current, a measure of ion transmembrane flux previously induced by addition of D-glucose. 15-acetyl-DON disrupted the Na(+)-D-glucose co-transport decreasing the jejunal glucose uptake (Awad et al. 2008).

The capacity of 15-acetyl-DON to activate the MAP Kinases ERK1/2, p38 and JNK, both in intestinal cell line, explants and jejunum from exposed pigs at a lower dose than DON and 3-acetyl-DON were correlated with their differential effects on intestinal barrier function. Moreover, 15-acetyl-DON demonstrated a higher capacity to decrease the expression of the tight junction proteins, claudin 3 and 4 (Pinton et al. 2012).

### Immunotoxicity

Very few studies have investigated the in vivo effect of 3-acetyl-DON and 15-acetyl-DON on the immune response (Fig. 5).

In mice orally exposed to 3-acetyl-DON, an increased dose-dependent humoral immune response was observed; but the Mitogen-induced lymphocyte proliferation and T cell-independent antibody responses were not altered (Kasali et al. 1985; Tomar et al. 1987). Exposure to acetyl-DONs also induced a transient upregulation of inflammatory cytokines in the spleen but to a lower extent than DON (Wu et al. 2014c).

In vitro exposure of human peripheral blood lymphocytes to 3-acetyl-DON inhibits their proliferative response and their ability to produce antibodies without altering their viability (Tomar et al. 1986). Exposure of human U937

macrophages to acetylated forms of DON led to induction of inflammatory cytokines in similar way than DON (Sugita-Konishi and Pestka 2001). Exposure of human intestinal cells Caco-2 to acetylated forms of DON induced the production of IL-8; regarding the production of this cytokine IL-8 the toxins could be ranked in the following order 3-acetyl-DON < DON < 15-acetyl-DON (Kadota et al. 2013).

A toxicity assessment aimed to compare the effects of DON and 3-acetyl-DON on virus hemagglutinin biosynthesis in Newcastle disease virus-infected BHK cells. The concentration required to reduce virus hemagglutinin was lowered by one-sixth between DON (300 ng/ml) and 3-acetyl-DON (2000 ng/ml) (Kimura et al. 1998).

### Effects of acetylated forms of DON on human and animal health

In mice, 15-acetyl-DON and DON demonstrated similar effects on induction of feed refusal and growth retardation (Forsell et al. 1987). In rats exposed to 0.15 mg/kg bw of 3-acetyl-DON, a 50 % of feed refusal was observed (Abbas et al. 1986). In swine, the doses inducing the lowest observed adverse effect level (LOAEL) for emesis following oral exposure to DON or 15-acetyl-DON were very similar (50–75  $\mu$ g/kg bw) (Pestka et al. 1987). Using a bioassay for anorexia induction, mice were administered DON, 3 and 15-acetyl-DON; both acetyl-DONs induced feed refusal comparable to DON; this was followed by a dose-dependent compensatory food intake as the cumulative food intakes were similar after 16 h (Wu et al. 2012). Mice are useful for studying food intake but lack an emetic reflex, and mink appears as a suitable animal model for investigating DON-induced emesis. In mink, the LOAELs following oral exposure were 50, 100 and 250  $\mu$ g/kg bw for DON, 15-acetyl-DON and 3-acetyl-DON, respectively. Moreover, 3-acetyl-DON revealed a longer latency for emetic effects that may be in link with the metabolic conversion of 3-acetyl-DON in DON in the digestive tract (Wu et al. 2013a, b).

DON-induced anorexia is linked with a plasma elevation of the satiety hormones cholecystokinin and peptide YY3–36. When mice were orally gavaged with 2.5 mg/kg bw DON, 3-acetyl-DON and 15-acetyl-DON, a similar elevation of plasmatic cholecystokinin was observed, consistent with the induction of food refusal. By contrast, 3 and 15-acetyl-DON exposure had only a moderate effect on plasma peptide YY3–36 (Wu et al. 2014d).

### Deoxynivalenol-3- $\beta$ -D-glucopyranoside

In natural conditions, some plants are able with efficacy to manage xenobiotics including mycotoxins. Plant

detoxification mechanisms include several chemical modifications, phase I transformation (hydrolysis, reduction or oxidation), phase II solubilization (conjugation) and phase III compartmentalization (Berthiller et al. 2013). DON-3- $\beta$ -D-glucopyranoside (D3G) is issued from phase II metabolism, glucosylation, of DON by the action of UDP-glycosyl-transferase enzymes naturally presents in the plant (Poppenberger et al. 2003; Berthiller et al. 2013). By convention, D3G is called “masked” mycotoxin *sensu stricto* as it results from a conjugation reaction in plant (Berthiller et al. 2013; Rychlik et al. 2014). D3G is found in cereals and cereal products all over the world at concentrations that can reach nearly half that of DON (Berthiller et al. 2009; Broekaert et al. 2015a).

### Toxicokinetics of D3G

The absolute oral bioavailability of D3G is approximately 2–5 times lower than the one of DON in rats, chicken and pigs (Nagl et al. 2012, 2014; Broekaert et al. 2016). In vitro, there is no evidence of transfer from the apical to baso-lateral side in human intestinal epithelial cells placed in transwell suggesting also low bioavailability of D3G in humans (De Nijs et al. 2012).

The distribution of D3G is in straight line with its poor absorption. Orally administrated D3G to Wistar rats was mainly recovered in the gastrointestinal tract, and less than 0.1 % of the administered dose was found as metabolite in the systemic compartment (Versilovskis and Bartkevics 2012). Similarly in pigs, the absorbed fraction of D3G is confined to intravascular fluids, as suggested by its lower volume of distribution (Broekaert et al. 2016).

Hydrolysis of D3G to DON is well established following incubation of the toxin with intestinal microbiota from human and pig (Dall’Erta et al. 2013; Gratz et al. 2013; Broekaert et al. 2016). Microbiota composition and abundance in the intestinal tract determine the efficacy of conversion of D3G into DON (Berthiller et al. 2013).

By contrast, the host has a negligible role in the hydrolysis of D3G to DON. In vitro, D3G is stable toward sequential chemical treatments mimicking the gastrointestinal digestive process of mammals (De Nijs et al. 2012; De Angelis et al. 2014). No hydrolysis could be observed in vivo after intravenous administration of D3G to chickens and pigs (Nagl et al. 2014; Broekaert et al. 2016). All together, these results underline the negligible role of host tissue in a hypothetical hydrolysis of the D3G conjugate back to its toxic precursor DON.

D3G showed a markedly shorter elimination half-life ( $54.1 \pm 3$  min) compared to DON ( $163 \pm 59$  min) in pig, which may be explained by higher polarity and lower distribution volume (Broekaert et al. 2016) (Fig. 2).

### Molecular effects and toxicity of D3G

Due to the sterical hindrance of the glucoside moiety, D3G, in contrast to DON, is not able to fit inside the A-site pocket of the peptidyl transferase center of the ribosome (Fig. 3). As a consequence, D3G does not induce a ribotoxic stress as demonstrated in human intestinal epithelial cells and in porcine jejunal explants. D3G did not activate JNK and p38 MAP kinases, did not alter viability and barrier function of intestinal cells, did not induced morphological lesions and pro-inflammatory gene expression in explants. A pan-genomic microarray analysis further demonstrated that the gene expression profile of D3G-treated explants was similar to that of control ones (Pierron et al. 2016b). Similarly, D3G did not elicit inflammatory response in the spleen of orally exposed mice (Wu et al. 2014c).

Using both a mouse and a mink model, Wu et al., demonstrated that D3G rapidly elicited an anorectic response that mimicked DON over a narrow dose range, but D3G was markedly less effective than DON-inducing emesis (Wu et al. 2014d).

### Bacterial modified forms of DON

Several aerobic and anaerobic bacterial strains are able to transform DON in “bacterial modified” forms. In many cases, these transformations lead to less toxic metabolites and they are being used as detoxification strategies (He et al. 2010; Karlovsky 2011). The main modifications are, reducing de-epoxidation to obtain the de-epoxy DON (DOM-1), epimerization for 3-epi-deoxynivalenol (3-epi-DON) and oxidation for 3-keto-deoxynivalenol (3-keto-DON) (Table 1).

#### DOM-1

The major bacterial modified form of DON is the de-epoxy-DON or DOM-1 issued from a de-epoxidation of the molecule (Yoshizawa et al. 1983). Several microbial strains isolated from rumen, chicken, pig, fish or soil are able to de-epoxidize DON (Table 1).

#### Toxicokinetics of DOM-1

No studies have investigated the intestinal absorption of the bacterial modified forms of DON. However, the fact that DOM-1 is found in blood and urine confirms that this modified form resulting solely from DON transformation by the microbiota is first absorbed from gut lumen (Maresca 2013).

**Table 1** Biological transformation of DON by microorganisms

Condition	Transformation reaction	Transformation product	Microorganism	Origin	References
<i>Anaerobic</i>	Reduction by deepoxidation	DOM-1	Rumen microorganisms (mixture)	(Dairy) Cow	King et al. (1984), Cote et al. (1986), Swanson et al. (1987), He et al. (1992)
			Chicken gut microorganisms (mixture)	Chicken	He et al. (1992)
			Pig gut microorganisms (mixture)	Pig	Kollarezik et al. (1994)
			A catfish digesta microbial culture (mixture)	Fish	Guan et al. (2009)
			Bacterial strain BBSH 797	Rumen fluid	Fuchs et al. (2002)
			Ten isolates belonging to <i>Clostridiales</i> , <i>Anaerofilum</i> , <i>Collinsella</i> , and <i>Bacillus</i> genera	Chicken digesta	Young et al. (2007), Yu et al. (2010)
			A culture from a mixture of 165 agricultural soils, containing six bacterial genera ( <i>Serratia</i> , <i>Clostridium</i> , <i>Citrobacter</i> , <i>Enterococcus</i> , <i>Stenotrophomonas</i> and <i>Streptomyces</i> ) (mixture)	Soil	Islam et al. (2012)
			<i>Agrobacterium Rhizobium</i> sp. E3-39	Soil	Shima et al. (1997)
			A culture D107 (mixture)	Agricultural commodity	Volkl et al. 2004
			<i>Aerobic</i>	Oxidation	3-keto-DON
<i>Deviosa</i> sp. 17-2-E-8	Soil	Zhou and He (2009)			
<i>Nocardiotodes</i> sp. WSN05-2	Soil	Ikunaga et al. (2011)			
Nine <i>Nocardiotodes</i> spp. and five <i>Deviosa</i> spp.	Soil	Sato et al. (2012)			
Epimerization	3-epi-DON				

The carry-over factors of DOM-1 in different edible tissues of food animals are globally low and suggest that this modified form of DON may be poorly distributed (Danicke and Brezina 2013). However, intra-uterine exposure of developing pig fetus to DOM-1 has been reported in pregnant sow fed a DON-contaminated diet (Goyarts et al. 2007).

Most publications only indirectly addressed the fate of DOM-1, as the microbiota-driven metabolite of DON. Very few focused on the intrinsic metabolism of orally supplied DOM-1. Regarding phase I metabolism, DOM-1 is not activated in vitro by the mixed-function oxidase of rat hepatic microsomal preparations (Cote et al. 1987). The preponderant formation of DOM-3-sulfate after oral administration of DOM-1 to both turkeys and chickens and the detection of DOM-1 glucuronide in the urine of rats fed a DON-contaminated diet suggest that phase II metabolism may follow the same species-oriented pathways reported for DON (Lattanzio et al. 2011; Schwartz-Zimmermann et al. 2015).

DOM-1 is found in feces and urine following oral or IV dosing of DON in many species (Prelusky et al. 1986; Erikson et al. 2002; Turner et al. 2010). DOM-1 is the only form for DON excretion in the milk of lactating cows (Cote et al. 1986; Seeling et al. 2006; Keese et al. 2008). Substantial amounts of the systemic DOM-1 can be excreted by bile route, joining the non-absorbed fraction of the microbiota-transformed DON and feeding an entero-hepatic cycling (Danicke and Brezina 2013) (Fig. 2).

#### *Molecular and cellular effects of DOM-1*

As DON, DOM-1 is also able to fit into the A-site of the peptidyl transferase center of the ribosome (Fig. 3). However, due to loss of the epoxy group, DOM-1 only forms two hydrogen bonds, resulting in less stable binding to the ribosome (Pierron et al. 2016c). The hydrogen bond between the DON epoxy group on C12 and one hydrogen from the sugar of the uracil U2873 is missing. As a consequence, DOM-1 do not activate JNK and p38 MAP kinases and do not induce a ribotoxic stress.

Using different cytotoxicity tests, DOM-1 was found not toxic to different cell types such as intestinal epithelial cells, lymphocytes or fibroblasts as compared to DON (Sundstol Eriksen et al. 2004; Schatzmayr et al. 2006; Pierron et al. 2016c). Commonly considered as a non-toxic metabolite of DON, DOM-1 did not impair the cell viability of proliferating or differentiated Caco-2 cells. Moreover, in mouse fibroblast and chicken lymphocytes, DOM-1 was respectively, 54 and 500 less toxic than DON.

Pierron et al., also demonstrated that DOM-1 did not change the oxygen consumption nor impair the barrier function of human intestinal cells. In intestinal explants, exposure to DOM-1 did not induce intestinal

lesions nor modified the transcriptional pattern as determined with a pan-genomic analysis. A special attention was given to the inflammatory response and as anticipated, DOM-1 was unable to elicit the expression of pro-inflammatory cytokines (Pierron et al. 2016c).

#### *Effects of DOM-1 on human and animal health*

Concerning the in vivo toxicity of DOM-1, there are only indirect studies on in vivo pigs experiments using contaminated feed supplemented with bacteria or digestive content able to transform DON into DOM-1. DOM-1 effects on zootechnical parameters were assessed based on the fact that the only major product of bacterial transformation of DON by these bacteria is the DOM-1 (He et al. 1992). For the two studies, animals receiving the contaminated feed, during 2 weeks or 9 days, with DON 5 mg/kg feed and the bacterial strain capable of de-epoxydise DON, showed no decrease in food intake or weight gain (He et al. 1993; Xi et al. 2011). However, the toxicity of the pure DOM-1 to intestine and on the all pig organism has not been evaluated.

#### **3-epi-DON and 3-keto-DON**

Several bacterial strains, all isolated from soil, have been described to epimerize DON into 3-epi-DON (Table 1). The only difference between DON and its epimer 3-epi-DON is the stereochemistry at the 3-OH group (He et al. 2015).

3-keto-DON can be obtained from the metabolization of DON by *Agrobacterium rhizobium* strain E3-39 (Table 1). It seems that, once formed, the molecule is stable and the transformation irreversible (Karlovsky 2011).

#### *Toxicokinetics of 3-epi-DON and 3-keto-DON*

There is still no report for the toxicokinetic of 3-epiDON and 3-ketoDON.

#### *Toxicity of 3-epi-DON and 3-keto-DON*

As described for DOM-1, 3-epi-DON and 3-keto-DON fit into the A-site of the peptidyl transferase center of the ribosome. In contrast to DON, which forms three hydrogen bonds, these modified forms only form two hydrogen bonds with the ribosome, leading to a less stable binding to the ribosome (Fig. 3). As a consequence, 3-epi DON does not activate JNK and p38 MAP Kinases and does not induce a ribotoxic stress (Pierron et al. 2016c). Based on the data presented in Fig. 3, we can also anticipate that 3-keto-DON will not induce a ribotoxic stress.

In vitro, 3-epi DON did not impair the viability of the human intestinal cell line Caco-2 and the murine fibroblast cell line 3T3. Similarly, 3-epi-DON did not change the oxygen consumption and the barrier function, as measured by the trans-epithelial resistance of human Caco-2 cells. As far as 3-keto-DON is concerned, one study investigated its cytotoxicity and demonstrated that this modified form of DON was about four times less toxic than DON on human intestinal Caco-2 cells and swiss mouse fibroblasts NIH/3T3 cells (He et al. 2015). Another study demonstrated that 3-keto-DON, in contrast to DON, did not decrease the proliferation of splenic murine (Shima et al. 1997; Pierron et al. 2016c).

*Ex vivo*, in porcine intestinal explant, Pierron et al. observed that 3-epi DON did not change the profile of gene expression on pan-genomic microarray and did not impact genes involved in oxidative pathway and inflammatory response (Pierron et al. 2016c).

In vivo, one study evaluated the toxicity of 3-epi-DON on mice (Wu et al. 2015). No organ (thymus, adrenal, stomach, colon, spleen, kidney, heart and liver tissues) was impacted by the 3-epi-DON treatment. Hematological and biochemical parameters were similar to control mice (Wu et al. 2015).

### Chemically modified forms of DON

Chemically modified forms of DON can be split in “thermally formed” and “nonthermally formed” (Rychlik et al. 2014). The thermal degradation products of DON have been identified as norDON A-F and 9-hydroxymethyl DON lactone in model heating experiments (Young et al. 1986; Bretz et al. 2006). However, some of these compounds, especially the norDON A–C degradation products, are also formed under alkaline conditions without thermal treatment (Young et al. 1986; Humpf and Voss 2004). Typical “non-thermally formed” forms of DON are the DON-sulfonates generated by treatment of contaminated material with sodium bisulfite (Young et al. 1986; Beyer et al. 2010). Of note, DON-sulfonates (DONS) have also been identified as DON metabolites in Wistar and Sprague–Dawley rats (Wan et al. 2013; Schwartz-Zimmermann et al. 2014a). Depending on the pH value, three DON-10-sulfonates termed DONS 1, 2 and 3 and differing in structure and stability are formed upon treatment of DON with sulfur reagents (Schwartz et al. 2013; Schwartz-Zimmermann et al. 2014b). DONS 1 is formed under neutral to alkaline conditions and characterized by loss of the epoxide group. It shows high stability in a broad pH range. DONS 3 is the initial reaction product of DON in cereals treated with sulfur reagents that is formed at slightly acidic to neutral pH. It occurs as a mixture of a ketone and a hemiketal form

and decomposes back to DON and to DONS 2 at neutral to alkaline pH. Besides the DONS 3 degradation process, DONS 2 can be directly formed under the neutral to alkaline conditions. It features a hemiketal between C-8 and C-15 and shows slight instability at pH > 8.

### Toxicokinetics of chemically modified forms of DON

No studies investigated the absorption and the metabolism of thermally modified forms of DON. There are conflicting results regarding the intestinal absorption of the chemically modified form DONS. In one study, similar blood concentrations of DONS and DON were reported in pigs fed equivalent amounts of either a DONS- or DON-contaminated feed, indicating that DONS are probably absorbed in the same extent as DON (Danicke et al. 2010). On the other hand, no DONS could be detected in blood following administration of an oral bolus of a DONS mixture (Paulick et al. 2015b). Further studies are thus required to elucidate the intestinal absorption of the DONS. With respect to their metabolism, preliminary observations suggest that DONS are converted back to DON, but it needs to be confirmed (Paulick et al. 2015b). The fate of the chemically formed DONS in feed is still unknown (Fig. 2).

### Toxicity of chemically modified forms of DON

The few existing data suggest that the thermal transformation may significantly reduce the toxicity of DON. Incubation of immortalized human kidney epithelial cells with up to 100  $\mu$ M norDON A, B and C did not decrease their viability compared to DON (Bretz et al. 2006).

Regarding the nonthermal chemical modifications of DON, few in vivo and in vitro data investigated their toxicity, though the exact profile of the DONS has rarely been specified. In three different toxicity tests, DONS 1 that is characterized by loss of the epoxide group revealed reduction in toxicity at least by a factor of 330 compared to DON, while DONS 2 and DONS 3 that both keep their epoxide group presented a toxicity only lowered by a factor 30 (Schwartz-Zimmermann et al. 2014a). In vitro unspecified DONS at concentrations up to 17  $\mu$ M did not alter the viability of unstimulated or concanavalin A-stimulated pig peripheral blood mononuclear cells, or pig intestinal epithelial cell lines (Danicke et al. 2010).

Pigs exposed to DONS orally or by intravenous route did not show clinical sign (Paulick et al. 2015a). Marginal effects were observed on the feed intake, live weight gain, liver function and clinical plasma parameters of weaned piglets orally exposed to DONS for 28 days (Danicke et al. 2010). This suggests that sulfonate-forms of DON are not toxic.

## Conclusion

New analytical techniques have increased our knowledge on emerging food contaminants and highlighted the presence of acetylated and modified forms of mycotoxins. As far as DON is concerned, the recent and ongoing unveiling of its acetylated and modified forms adds to the traditional concerns about the food and feed safety issues of this mycotoxin and the existing knowledge gaps in many research aspects. The structural modifications of DON may result from chemical transformations catalyzed by plant enzymes, especially those involved in detoxification processes, bacterial or fungal enzymes. The modifications may also result from food/feed processing or detoxification treatments.

Several aspects of the toxicity of DON are well characterized. At the molecular level, the binding of DON to ribosomes inhibits the elongation step of the protein synthesis concurrently with the onset of a ribotoxic stress which leads to the activation of critical cellular kinases involved in signal transduction. The implications at the cellular level include oxidative stress response, disrupted cell signaling, impaired cell growth and differentiation and apoptosis. Regarding organs especially toward the intestine, DON impairs the barrier function and the nutrient uptake. It also interferes with local immune response. Thus it cannot be excluded that this mycotoxin plays a role in the etiology of some chronic intestinal inflammatory diseases in adult, and participate to food allergy reactions in childhood. With respect to systemic toxicity, the primary physiological response to acute DON exposure in many species is emesis. DON also affects hematopoiesis and induces neuro-endocrine effects. Several animal studies also pointed out the potential for chronic effects of DON on growth, reproduction and immune function. Chronic exposure to DON has been linked to increased susceptibility to infections and chronic diseases.

As highlighted in this review; much less data are available regarding the metabolism and toxicity of the acetylated and modified forms of DON. Nevertheless, the toxicokinetics of these acetylated and modified forms of DON are gaining more attention as these compounds may be converted back to DON following cleavage by gut microbiota and host tissue metabolism. Also, the changes in physico-chemical properties associated with the different structural modifications of DON may have impact on the gastro-intestinal absorption and subsequent plasma concentrations of the acetylated and modified forms. Moreover, these forms of DON can be involved in complex metabolism processes including entero-hepatic cycling. As of now, toxicokinetics data indicate that the reduced polarity of the acetylated forms obviously enhances their oral absorption.

If the acetylated forms 3-acetyl-DON and 15-acetyl-DON induce, as DON, adverse effects on numerous parameters, data concerning the modified forms, D3G, DOM-1, 3-epi-DON, 3-keto-DON and DONS, demonstrate the significant decreases in the intrinsic toxic effects.

Overall considering that modified mycotoxins could be released, hydrolyzed, biotransformed and absorbed in the gastrointestinal tract primarily as their parent compound, the EFSA Panel on Contaminants in the Food Chain (CONTAM Panel) recently delivered a scientific opinion on the risk to human and animal health related to the presence of modified forms of major *Fusarium* toxins excepted DON (EFSA 2014). The same critical evaluation is welcomed specifically for the DON modified forms, owing to the narrow margin between exposure to DON of some consumer groups and tolerable daily intake. In that regard, it is also expected that this opinion will not ignore the specific issue of the cumulative risk assessment of DON, its acetylated and modified forms, since convergent indications suggest at least for DON and co-occurring acetylated forms, that their combined toxicity is higher than predicted from individual effects (Alassane-Kpembé et al. 2016).

**Acknowledgments** AP was supported by a CIFRE fellowship (2012/0572, jointly financed by the BIOMIN holding GmbH, ANRT and INRA). This work was supported in part by the ANR (Agence Nationale de la Recherche) projects ImBio (ANR-13-CESA-0003-03) and CaDON (ANR-15-CE21-0001-02) and the Partenariat Hubert Curien project (PHC 35813XM).

## References

- Abbas HK, Mirocha CJ, Tuite J (1986) Natural occurrence of deoxynivalenol, 15-acetyl-deoxynivalenol, and zearalenone in refusal factor corn stored since 1972. *Appl Environ Microbiol* 51:841–843
- Ajandouz EH, Berdah S, Moutardier V et al (2016) Hydrolytic Fate of 3/15-acetyldeoxynivalenol in Humans: specific Deacetylation by the Small Intestine and Liver Revealed Using in Vitro and ex Vivo Approaches. *Toxins*. doi:10.3390/toxins8080232
- Akbari P, Braber S, Gremmels H et al (2014) Deoxynivalenol: a trigger for intestinal integrity breakdown. *FASEB J* 28:2414–2429. doi:10.1096/fj.13-238717
- Akbari P, Braber S, Varasteh S et al (2016) The intestinal barrier as an emerging target in the toxicological assessment of mycotoxins. *Arch Toxicol*. doi:10.1007/s00204-016-1794-8
- Alassane-Kpembé I, Kolf-Clauw M, Gauthier T et al (2013) New insights into mycotoxin mixtures: the toxicity of low doses of Type B trichothecenes on intestinal epithelial cells is synergistic. *Toxicol Appl Pharmacol* 272:191–198. doi:10.1016/j.taap.2013.05.023
- Alassane-Kpembé I, Puel O, Oswald IP (2015) Toxicological interactions between the mycotoxins deoxynivalenol, nivalenol and their acetylated derivatives in intestinal epithelial cells. *Arch Toxicol* 89:1337–1346. doi:10.1007/s00204-014-1309-4
- Alassane-Kpembé I, Schatzmayr G, Taranu I et al (2016) Mycotoxins co-contamination: methodological aspects and biological

- relevance of combined toxicity studies. *Crit Rev Food Sci Nutr*. doi:10.1080/10408398.2016.1140632
- Amuzie CJ, Shinozuka J, Pestka JJ (2009) Induction of suppressors of cytokine signaling by the trichothecene deoxynivalenol in the mouse. *Toxicol Sci* 111:277–287. doi:10.1093/toxsci/kfp150
- Antonissen G, Van Immerseel F, Pasmans F et al (2014) The mycotoxin deoxynivalenol predisposes for the development of *Clostridium perfringens*-induced necrotic enteritis in broiler chickens. *PLoS One* 9:e108775. doi:10.1371/journal.pone.0108775
- Antonissen G, Van Immerseel F, Pasmans F et al (2015) Mycotoxins deoxynivalenol and fumonisins alter the extrinsic component of intestinal barrier in broiler chickens. *J Agric Food Chem* 63:10846–10855. doi:10.1021/acs.jafc.5b04119
- Awad WA, Zentek J (2015) The feed contaminant deoxynivalenol affects the intestinal barrier permeability through inhibition of protein synthesis. *Arch Toxicol*. 89:961–965. doi:10.1007/s00204-014-1284-9
- Awad WA, Razzazi-Fazeli E, Bohm J, Zentek J (2008) Effects of B-trichothecenes on luminal glucose transport across the isolated jejunal epithelium of broiler chickens. *J Anim Physiol Anim Nutr* 92:225–230. doi:10.1111/j.1439-0396.2007.00709.x
- Awad WA, Ghareeb K, Dadak A et al (2012) Genotoxic effects of deoxynivalenol in broiler chickens fed low-protein feeds. *Poult Sci* 91:550–555. doi:10.3382/ps.2011-01742
- Awad WA, Ghareeb K, Zentek J (2014) Mechanisms underlying the inhibitory effect of the feed contaminant deoxynivalenol on glucose absorption in broiler chickens. *Vet J* 202:188–190. doi:10.1016/j.tvjl.2014.06.012
- Azcona-Olivera JI, Ouyang Y, Murtha J et al (1995) Induction of cytokine mRNAs in mice after oral exposure to the trichothecene vomitoxin (deoxynivalenol): relationship to toxin distribution and protein synthesis inhibition. *Toxicol Appl Pharmacol* 133:109–120. doi:10.1006/taap.1995.1132
- Behrens M, Huwel S, Galla H-J, Humpf H-U (2015) Blood-brain barrier effects of the fusarium mycotoxins deoxynivalenol, 3 acetyldeoxynivalenol, and moniliformin and their transfer to the brain. *PLoS One* 10:e0143640. doi:10.1371/journal.pone.0143640
- Bensassi F, El Golli-Bennour E, Abid-Essefi S et al (2009) Pathway of deoxynivalenol-induced apoptosis in human colon carcinoma cells. *Toxicology* 264:104–109. doi:10.1016/j.tox.2009.07.020
- Berthiller F, Dall'asta C, Corradini R et al (2009) Occurrence of deoxynivalenol and its 3-beta-D-glucoside in wheat and maize. *Food Addit Contam Part A Chem Anal Control Expo Risk Assess* 26:507–511. doi:10.1080/02652030802555668
- Berthiller F, Crews C, Dall'asta C et al (2013) Masked mycotoxins: a review. *Mol Nutr Food Res* 57:165–186. doi:10.1002/mnfr.201100764
- Beyer M, Danicke S, Rohweder D, Humpf H-U (2010) Determination of deoxynivalenol-sulfonate (DONS) in cereals by hydrophilic interaction chromatography coupled to tandem mass spectrometry. *Mycotoxin Res* 26:109–117. doi:10.1007/s12550-010-0046-7
- Bianco G, Fontanella B, Severino L et al (2012) Nivalenol and deoxynivalenol affect rat intestinal epithelial cells: a concentration related study. *PLoS One* 7:e52051. doi:10.1371/journal.pone.0052051
- Bol-Schoenmakers M, Braber S, Akbari P et al (2016) The mycotoxin deoxynivalenol facilitates allergic sensitization to whey in mice. *Mucosal Immunol*. doi:10.1038/mi.2016.13
- Bondy GS, Pestka JJ (1991) Dietary exposure to the trichothecene vomitoxin (deoxynivalenol) stimulates terminal differentiation of Peyer's patch B cells to IgA secreting plasma cells. *Toxicol Appl Pharmacol* 108:520–530. doi:10.1016/0041-008x(91)90098-y
- Bony S, Carcelen M, Olivier L, Devaux A (2006) Genotoxicity assessment of deoxynivalenol in the Caco-2 cell line model using the Comet assay. *Toxicol Lett* 166:67–76. doi:10.1016/j.toxlet.2006.04.010
- Bracarense APFL, Luciola J, Grenier B et al (2012) Chronic ingestion of deoxynivalenol and fumonisin, alone or in interaction, induces morphological and immunological changes in the intestine of piglets. *Br J Nutr* 107:1776–1786. doi:10.1017/S0007114511004946
- Bracarense APFL, Basso KM, Silva EO, Oswald IP (2016) Deoxynivalenol in liver and lymphoid organs of rats effects of dose and duration on immunohistological changes. *World Mycotoxins J*. doi:10.3920/wmj2016.2094
- Bretz M, Beyer M, Cramer B et al (2006) Thermal degradation of the Fusarium mycotoxin deoxynivalenol. *J Agric Food Chem* 54:6445–6451. doi:10.1021/jf061008g
- Broekaert N, Devreese M, De Baere S et al (2015a) Modified Fusarium mycotoxins unmasked: from occurrence in cereals to animal and human excretion. *Food Chem Toxicol* 80:17–31. doi:10.1016/j.fct.2015.02.015
- Broekaert N, Devreese M, De Mil T et al (2015b) Oral bioavailability, hydrolysis, and comparative toxicokinetics of 3-acetyldeoxynivalenol and 15-acetyldeoxynivalenol in broiler chickens and pigs. *J Agric Food Chem* 63:8734–8742. doi:10.1021/acs.jafc.5b03270
- Broekaert N, Devreese M, van Bergen T et al (2016) In vivo contribution of deoxynivalenol-3-beta-D-glucoside to deoxynivalenol exposure in broiler chickens and pigs: oral bioavailability, hydrolysis and toxicokinetics. *Arch Toxicol*. doi:10.1007/s00204-016-1710-2
- Canady RA, Coker RD, Rgan SK et al (2001) Deoxynivalenol. Safety evaluation of certain mycotoxins in food. *WHO Food Addit Ser* 47:420–555
- Cano PM, Seeboth J, Meurens F et al (2013) Deoxynivalenol as a new factor in the persistence of intestinal inflammatory diseases: an emerging hypothesis through possible modulation of Th17-mediated response. *PLoS One* 8:e53647. doi:10.1371/journal.pone.0053647
- Cheat S, Gerez JR, Cognie J et al (2015) Nivalenol has a greater impact than deoxynivalenol on pig jejunum mucosa in vitro on explants and in vivo on intestinal loops. *Toxins* 7:1945–1961. doi:10.3390/toxins7061945
- Chung YJ, Zhou HR, Pestka JJ (2003) Transcriptional and posttranscriptional roles for p38 mitogen-activated protein kinase in upregulation of TNF-alpha expression by deoxynivalenol (vomitoxin). *Toxicol Appl Pharmacol* 193:188–201. doi:10.1016/S0041-008X(03)00299-0
- Cobb MH (1999) MAP kinase pathways. *Prog Biophys Mol Biol* 71:479–500. doi:10.1016/S0079-6107(98)00056-X
- Collins TFX, Sprando RL, Black TN et al (2006) Effects of deoxynivalenol (DON, vomitoxin) on in utero development in rats. *Food Chem Toxicol* 44:747–757. doi:10.1016/j.fct.2005.10.007
- European Commission (2007) Commission Regulation (EC) No 1126/2007 of 28 September 2007 amending Regulation (EC) No 1881/2006 setting maximum levels for certain contaminants in foodstuffs as regards Fusarium toxins in maize and maize products. *Off J Eur Union L* 255(2007):14–17
- Cote LM, Dahlem AM, Yoshizawa T et al (1986) Excretion of deoxynivalenol and its metabolite in milk, urine, and feces of lactating dairy cows. *J Dairy Sci* 69:2416–2423. doi:10.3168/jds.S0022-0302(86)80681-6
- Cote LM, Buck W, Jeffery E (1987) Lack of hepatic microsomal metabolism of deoxynivalenol and its metabolite, DOM-1. *Food Chem Toxicol* 25:291–295. doi:10.1016/0278-6915(87)90125-6
- Dall'Erta A, Cirlini M, Dall'Asta M et al (2013) Masked mycotoxins are efficiently hydrolyzed by human colonic microbiota releasing their aglycones. *Chem Res Toxicol* 26:305–312. doi:10.1021/tx300438c

- Danicke S, Brezina U (2013) Kinetics and metabolism of the Fusarium toxin deoxynivalenol in farm animals: consequences for diagnosis of exposure and intoxication and carry over. *Food Chem Toxicol* 60:58–75. doi:[10.1016/j.fct.2013.07.017](https://doi.org/10.1016/j.fct.2013.07.017)
- Danicke S, Valenta H, Doll S (2004) On the toxicokinetics and the metabolism of deoxynivalenol (DON) in the pig. *Arch Anim Nutr* 58:169–180. doi:[10.1080/00039420410001667548](https://doi.org/10.1080/00039420410001667548)
- Danicke S, Brussow K-P, Goyarts T et al (2007) On the transfer of the Fusarium toxins deoxynivalenol (DON) and zearalenone (ZON) from the sow to the full-term piglet during the last third of gestation. *Food Chem Toxicol* 45:1565–1574. doi:[10.1016/j.fct.2007.02.016](https://doi.org/10.1016/j.fct.2007.02.016)
- Danicke S, Hegewald AK, Kahler S et al (2010) Studies on the toxicity of deoxynivalenol (DON), sodium metabisulfite, DON-sulfonate (DONS) and de-epoxy-DON for porcine peripheral blood mononuclear cells and the Intestinal Porcine Epithelial Cell lines IPEC-1 and IPEC-J2, and on effects of DON and DONS. *Food Chem Toxicol* 48:2154–2162. doi:[10.1016/j.fct.2010.05.022](https://doi.org/10.1016/j.fct.2010.05.022)
- De Angelis E, Monaci L, Visconti A (2014) Investigation on the stability of deoxynivalenol and DON-3 glucoside during gastro-duodenal in vitro digestion of a naturally contaminated bread model food. *Food Control* 43:270–275. doi:[10.1016/j.foodcont.2014.03.032](https://doi.org/10.1016/j.foodcont.2014.03.032)
- De Nijs M, Van den Top HJ, Portier L et al (2012) Digestibility and absorption of deoxynivalenol-3- $\beta$ -glucoside in in vitro models. *World Mycotoxin J* 5:319–324. doi:[10.3920/WMJ2012.1430](https://doi.org/10.3920/WMJ2012.1430)
- Del Regno M, Adesso S, Popolo A et al (2015) Nivalenol induces oxidative stress and increases deoxynivalenol pro-oxidant effect in intestinal epithelial cells. *Toxicol Appl Pharmacol* 285:118–127. doi:[10.1016/j.taap.2015.04.002](https://doi.org/10.1016/j.taap.2015.04.002)
- Diesing AK, Nossol C, Danicke S et al (2011) Vulnerability of polarised intestinal porcine epithelial cells to mycotoxin deoxynivalenol depends on the route of application. *PLoS One* 6:e17472. doi:[10.1371/journal.pone.0017472](https://doi.org/10.1371/journal.pone.0017472)
- Dinu D, Bodea GO, Ceapa CD et al (2011) Adapted response of the antioxidant defense system to oxidative stress induced by deoxynivalenol in Hek-293 cells. *Toxicol* 57:1023–1032. doi:[10.1016/j.toxicol.2011.04.006](https://doi.org/10.1016/j.toxicol.2011.04.006)
- EFSA (2014) Scientific opinion on the risks for human and animal health related to the presence of modified forms of certain mycotoxins in food and feed. *EFSA J* 12:3916. doi:[10.2903/j.efsa.2014.3916](https://doi.org/10.2903/j.efsa.2014.3916)
- Eriksen GS, Alexander J (1998) Fusarium toxins in cereals—a risk assessment. Nordic Council of Ministers, Copenhagen, Denmark.
- Eriksen GS, Pettersson H (2004) Toxicological evaluation of trichothecenes in animal feed. *Anim Feed Sci Technol* 114:205–239. doi:[10.1016/j.anifeeds.2003.08.008](https://doi.org/10.1016/j.anifeeds.2003.08.008)
- Eriksen GS, Pettersson H, Johnsen K, Lindberg JE (2002) Transformation of trichothecenes in ileal digesta and faeces from pigs. *Arch Tierernahr* 56:263–274. doi:[10.1080/00039420214343](https://doi.org/10.1080/00039420214343)
- Eriksen GS, Pettersson H, Lindberg JE (2003) Absorption, metabolism and excretion of 3-acetyl DON in pigs. *Arch Tierernahr* 57:335–345. doi:[10.1080/00039420310001607699](https://doi.org/10.1080/00039420310001607699)
- Eriksen GS, Pettersson H, Lundh T (2004) Comparative cytotoxicity of deoxynivalenol, nivalenol, their acetylated derivatives and de-epoxy metabolites. *Food Chem Toxicol* 42:619–624. doi:[10.1016/j.fct.2003.11.006](https://doi.org/10.1016/j.fct.2003.11.006)
- FDA (2010) Guidance for industry and FDAi advisory levels for deoxynivalenol (DON) in finished wheat products for human consumption and grains and grain by-products used for animal feed. <http://www.fda.gov/downloads/Food/GuidanceRegulation/UCM217558.pdf>
- Fioramonti J, Dupuy C, Dupuy J, Bueno L (1993) The mycotoxin, deoxynivalenol, delays gastric emptying through serotonin-3 receptors in rodents. *J Pharmacol Exp Ther* 266:1255–1260
- Flannery BM, Clark ES, Pestka JJ (2012) Anorexia induction by the trichothecene deoxynivalenol (vomitoxin) is mediated by the release of the gut satiety hormone peptide YY. *Toxicol Sci* 130:289–297. doi:[10.1093/toxsci/kfs255](https://doi.org/10.1093/toxsci/kfs255)
- Forsell JH, Pestka JJ (1985) Relation of 8-ketotrichothecene and zearalenone analog structure to inhibition of mitogen-induced human lymphocyte blastogenesis. *Appl Environ Microbiol* 50:1304–1307
- Forsell JH, Jensen R, Tai JH et al (1987) Comparison of acute toxicities of deoxynivalenol (vomitoxin) and 15-acetyldeoxynivalenol in the B6C3F1 mouse. *Food Chem Toxicol* 25:155–162. doi:[10.1016/0278-6915\(87\)90149-9](https://doi.org/10.1016/0278-6915(87)90149-9)
- Frankic T, Pajk T, Rezar V et al (2006) The role of dietary nucleotides in reduction of DNA damage induced by T-2 toxin and deoxynivalenol in chicken leukocytes. *Food Chem Toxicol* 44:1838–1844. doi:[10.1016/j.fct.2006.06.002](https://doi.org/10.1016/j.fct.2006.06.002)
- Fuchs E, Binder EM, Heidler D, Krska R (2002) Structural characterization of metabolites after the microbial degradation of type A trichothecenes by the bacterial strain BBSH 797. *Food Addit Contam* 19:379–386. doi:[10.1080/02652030110091154](https://doi.org/10.1080/02652030110091154)
- Gaige S, Bonnet MS, Tardivel C et al (2013) c-Fos immunoreactivity in the pig brain following deoxynivalenol intoxication: focus on NUCB2/nesfatin-1 expressing neurons. *Neurotoxicology* 34:135–149. doi:[10.1016/j.neuro.2012.10.020](https://doi.org/10.1016/j.neuro.2012.10.020)
- Gareis M, Bauer J, Thiem J et al (1990) Cleavage of zearalenone-glycoside, a “masked” mycotoxin, during digestion in swine. *Zentralbl Veterinarmed B* 37:236–240. doi:[10.1111/j.1439-0450.1990.tb01052.x](https://doi.org/10.1111/j.1439-0450.1990.tb01052.x)
- Garreau de Loubresse N, Prokhorova I, Holtkamp W et al (2014) Structural basis for the inhibition of the eukaryotic ribosome. *Nature* 513:517–522. doi:[10.1038/nature13737](https://doi.org/10.1038/nature13737)
- Ghareeb K, Awad WA, Böhm J, Zebeli Q (2015) Impacts of the feed contaminant deoxynivalenol on the intestine of monogastric animals: poultry and swine. *J Appl Toxicol* 35:327–337. doi:[10.1002/jat.3083](https://doi.org/10.1002/jat.3083)
- Ghareeb K, Awad WA, Zebeli Q, Böhm J (2016) Deoxynivalenol in chicken feed alters the vaccinal immune response and clinical biochemical serum parameters but not the intestinal and carcass characteristics. *J Anim Physiol Anim Nutr* 100:53–60. doi:[10.1111/jpn.12328](https://doi.org/10.1111/jpn.12328)
- Gilbert J, Brûlé-Babel A, Guerrieri AT et al (2014) Ratio of 3-ADON and 15-ADON isolates of *Fusarium graminearum* recovered from wheat kernels in Manitoba from 2008 to 2012. *Can J Plant Pathol* 36:54–63. doi:[10.1080/07060661.2014.887033](https://doi.org/10.1080/07060661.2014.887033)
- Girardet C, Bonnet MS, Jdir R et al (2011a) The food-contaminant deoxynivalenol modifies eating by targeting anorexigenic neurocircuitry. *PLoS One* 6:e26134. doi:[10.1371/journal.pone.0026134](https://doi.org/10.1371/journal.pone.0026134)
- Girardet C, Bonnet MS, Jdir R et al (2011b) Central inflammation and sickness-like behavior induced by the food contaminant deoxynivalenol: a PGE2-independent mechanism. *Toxicol Sci* 124:179–191. doi:[10.1093/toxsci/kfr219](https://doi.org/10.1093/toxsci/kfr219)
- Girgis GN, Sharif S, Barta JR et al (2008) Immunomodulatory effects of feed-borne *Fusarium* mycotoxins in chickens infected with coccidia. *Exp Biol Med* 233:1411–1420. doi:[10.3181/0805-RM-173](https://doi.org/10.3181/0805-RM-173)
- Girgis GN, Barta JR, Brash M, Smith TK (2010) Morphologic changes in the intestine of broiler breeder pullets fed diets naturally contaminated with *Fusarium* mycotoxins with or without coccidial challenge. *Avian Dis* 54:67–73. doi:[10.1637/8945-052809-Reg.1](https://doi.org/10.1637/8945-052809-Reg.1)
- Girish CK, Smith TK, Boermans HJ, Karrow NA (2008) Effects of feeding blends of grains naturally contaminated with *Fusarium* mycotoxins on performance, hematology, metabolism, and immunocompetence of turkeys. *Poult Sci* 87:421–432. doi:[10.3382/ps.2007-00181](https://doi.org/10.3382/ps.2007-00181)

- Gouze ME, Laffitte J, Rouimi P et al (2006) Effect of various doses of deoxynivalenol on liver xenobiotic metabolizing enzymes in mice. *Food Chem Toxicol* 44:476–483. doi:10.1016/j.fct.2005.08.020
- Goyarts T, Danicke S (2006) Bioavailability of the Fusarium toxin deoxynivalenol (DON) from naturally contaminated wheat for the pig. *Toxicol Lett* 163:171–182. doi:10.1016/j.toxlet.2005.10.007
- Goyarts T, Danicke S, Brussow K-P et al (2007) On the transfer of the Fusarium toxins deoxynivalenol (DON) and zearalenone (ZON) from sows to their fetuses during days 35–70 of gestation. *Toxicol Lett* 171:38–49. doi:10.1016/j.toxlet.2007.04.003
- Gratz SW, Duncan G, Richardson AJ (2013) The human fecal microbiota metabolizes deoxynivalenol and deoxynivalenol-3-glucoside and may be responsible for urinary de-epoxy-deoxynivalenol. *Appl Environ Microbiol* 79:1821–1825. doi:10.1128/AEM.02987-12
- Graziani F, Pujol A, Nicoletti C et al (2015) The food-associated ribotoxin deoxynivalenol modulates inducible NO synthase in human intestinal cell model. *Toxicol Sci* 145:372–382. doi:10.1093/toxsci/kfv058
- Grenier B, Loureiro-Bracarense AP, Luciola J et al (2011) Individual and combined effects of subclinical doses of deoxynivalenol and fumonisins in piglets. *Mol Nutr Food Res* 55:761–771. doi:10.1002/mnfr.201000402
- Guan S, He J, Young JC, Zhu H, Li XZ, Ji C, Zhou T (2009) Transformation of trichothecene mycotoxins by microorganisms from fish digesta. *Aquaculture* 290:290–295. doi:10.1016/j.aquaculture.2009.02.037
- Hara-Kudo Y, Sugita-Konoshi Y, Kasuga F, Kumagai S (1996) Effects of deoxynivalenol on Salmonella enteritidis infection. *Mycotoxins* 42:51–55. doi:10.2520/myco1975.1996.51
- He P, Young LG, Forsberg C (1992) Microbial transformation of deoxynivalenol (vomitoxin). *Appl Environ Microbiol* 58:3857–3863
- He P, Young LG, Forsberg C (1993) Microbially detoxified vomitoxin-contaminated corn for young pigs. *J Anim Sci* 71:963–967. doi:10.2527/1993.714963x
- He J, Zhou T, Young JC et al (2010) Chemical and biological transformations for detoxification of trichothecene mycotoxins in human and animal food chains: a review. *Trends Food Sci Technol* 21:67–76. doi:10.1016/j.tifs.2009.08.002
- He K, Zhou HR, Pestka JJ (2012) Targets and intracellular signaling mechanisms for deoxynivalenol-induced ribosomal RNA cleavage. *Toxicol Sci* 127:382–390. doi:10.1093/toxsci/kfs134
- He JW, Bondy GS, Zhou T et al (2015) Toxicology of 3-epi-deoxynivalenol, a deoxynivalenol-transformation product by *Devosia* mutants 17-2-E-8. *Food Chem Toxicol* 84:250–259. doi:10.1016/j.fct.2015.09.003
- Hedman R, Pettersson H (1997) Transformation of nivalenol by gastrointestinal microbes. *Arch Tierernahr* 50:321–329. doi:10.1080/17450399709386142
- Humpf H-U, Voss KA (2004) Effects of thermal food processing on the chemical structure and toxicity of fumonisin mycotoxins. *Mol Nutr Food Res* 48:255–269. doi:10.1002/mnfr.200400033
- Hymery N, Sibiril Y, Parent-Massin D (2006) In vitro effects of trichothecenes on human dendritic cells. *Toxicol In Vitro* 20:899–909. doi:10.1016/j.tiv.2006.01.015
- IARC (1993) Some naturally occurring substances: food items and constituents, heterocyclic aromatic amines and mycotoxins. In: *Monographs on the evaluation of carcinogenic risks to humans*, Vol 56. International Agency for Research on Cancer, World Health Organization, Lyon, France, pp 397–444
- Ikunaga Y, Sato I, Grond S et al (2011) *Nocardioides* sp. strain WSN05-2, isolated from a wheat field, degrades deoxynivalenol, producing the novel intermediate 3-epi-deoxynivalenol. *Appl Microbiol Biotechnol* 89:419–427. doi:10.1007/s00253-010-2857-z
- Islam R, Zhou T, Young JC et al (2012) Aerobic and anaerobic de-epoxydation of mycotoxin deoxynivalenol by bacteria originating from agricultural soil. *World J Microbiol Biotechnol* 28:7–13. doi:10.1007/s11274-011-0785-4
- Ito M, Sato I, Ishizaka M et al (2013) Bacterial cytochrome P450 system catabolizing the Fusarium toxin deoxynivalenol. *Appl Environ Microbiol* 79:1619–1628. doi:10.1128/AEM.03227-12
- Kadota T, Furusawa H, Hirano S et al (2013) Comparative study of deoxynivalenol, 3-acetyldeoxynivalenol, and 15-acetyldeoxynivalenol on intestinal transport and IL-8 secretion in the human cell line Caco-2. *Toxicol Vitro* 27:1888–1895. doi:10.1016/j.tiv.2013.06.003
- Kalaiselvi P, Rajashree K, Priya LB, Padma VV (2013) Cytoprotective effect of epigallocatechin-3-gallate against deoxynivalenol-induced toxicity through anti-oxidative and anti-inflammatory mechanisms in HT-29 cells. *Food Chem Toxicol* 56:110–118. doi:10.1016/j.fct.2013.01.042
- Karlovsky P (2011) Biological detoxification of the mycotoxin deoxynivalenol and its use in genetically engineered crops and feed additives. *Appl Microbiol Biotechnol* 91:491–504. doi:10.1007/s00253-011-3401-5
- Kasali OB, Schiefer HB, Hancock DS et al (1985) Subacute toxicity of dietary 3-acetyldeoxynivalenol in mice. *Can J Comp Med* 49:319–322
- Keese C, Meyer U, Valenta H et al (2008) No carry over of unmetabolised deoxynivalenol in milk of dairy cows fed high concentrate proportions. *Mol Nutr Food Res* 52:1514–1529. doi:10.1002/mnfr.200800077
- Khera KS, Arnold DL, Whalen C et al (1984) Vomitoxin (4-deoxynivalenol): effects on reproduction of mice and rats. *Toxicol Appl Pharmacol* 74:345–356. doi:10.1016/0041-008X(84)90288-6
- Khera KS, Whalen C, Angers G (1986) A teratology study on vomitoxin (4-deoxynivalenol) in rabbits. *Food Chem Toxicol* 24:421–424. doi:10.1016/0278-6915(86)90207-3
- Kimura M, Kaneko I, Komiyama M et al (1998) Trichothecene 3-O-acetyltransferase protects both the producing organism and transformed yeast from related mycotoxins. Cloning and characterization of Tri101. *J Biol Chem* 273:1654–1666. doi:10.1074/jbc.273.3.1654
- King RR, Mcqueen RE, Levesque D, Greenhalgh R (1984) Transformation of deoxynivalenol (vomitoxin) by rumen microorganisms. *J Agri Food Chem* 32:1181–1183. doi:10.1021/jf00125a061
- Kluger B, Bueschl C, Lemmens M et al (2013) Stable isotopic labeling-assisted untargeted metabolic profiling reveals novel conjugates of the mycotoxin deoxynivalenol in wheat. *Anal Bioanal Chem* 405:5031–5036. doi:10.1007/s00216-012-6483-8
- Kolf-Claw M, Castellote J, Joly B et al (2009) Development of a pig jejunal explant culture for studying the gastrointestinal toxicity of the mycotoxin deoxynivalenol: histopathological analysis. *Toxicol In Vitro* 23:1580–1584. doi:10.1016/j.tiv.2009.07.015
- Kollarczik B, Gareis M, Hanelt M (1994) In vitro transformation of the Fusarium mycotoxins deoxynivalenol and zearalenone by the normal gut microflora of pigs. *Nat Toxins* 2:105–110. doi:10.1002/mt.2620020303
- Krishnaswamy R, Devaraj SN, Padma VV (2010) Lutein protects HT-29 cells against Deoxynivalenol-induced oxidative stress and apoptosis: prevention of NF-kappaB nuclear localization and down regulation of NF-kappaB and Cyclo-Oxygenase-2 expression. *Free Radic Biol Med* 49:50–60. doi:10.1016/j.freeradbiomed.2010.03.016
- Lake BG, Phillips JC, Walters DG et al (1987) Studies on the metabolism of deoxynivalenol in the rat. *Food Chem Toxicol* 25:589–592. doi:10.1016/0278-6915(87)90019-6

- Larsen JC, Hunt J, Perrin I, Ruckebauer P (2004) Workshop on trichothecenes with a focus on DON: summary report. *Toxicol Lett* 153:1–22. doi:[10.1016/j.toxlet.2004.04.020](https://doi.org/10.1016/j.toxlet.2004.04.020)
- Lattanzio VMT, Solfrizzo M, De Girolamo A et al (2011) LC-MS/MS characterization of the urinary excretion profile of the mycotoxin deoxynivalenol in human and rat. *J Chromatogr B Anal Technol Biomed Life Sci* 879:707–715. doi:[10.1016/j.jchromb.2011.01.029](https://doi.org/10.1016/j.jchromb.2011.01.029)
- Li M, Cuff CF, Pestka JJ (2005) Modulation of murine host response to enteric reovirus infection by the trichothecene deoxynivalenol. *Toxicol Sci* 87:134–145. doi:[10.1093/toxsci/kfi225](https://doi.org/10.1093/toxsci/kfi225)
- Li M, Harkema JR, Cuff CF, Pestka JJ (2007) Deoxynivalenol exacerbates viral bronchopneumonia induced by respiratory reovirus infection. *Toxicol Sci* 95:412–426. doi:[10.1093/toxsci/kfl153](https://doi.org/10.1093/toxsci/kfl153)
- Li D, Ma H, Ye Y et al (2014a) Deoxynivalenol induces apoptosis in mouse thymic epithelial cells through mitochondria-mediated pathway. *Environ Toxicol Pharmacol* 38:163–171. doi:[10.1016/j.etap.2014.05.015](https://doi.org/10.1016/j.etap.2014.05.015)
- Li D, Ye Y, Lin S et al (2014b) Evaluation of deoxynivalenol-induced toxic effects on DF-1 cells in vitro: cell-cycle arrest, oxidative stress, and apoptosis. *Environ Toxicol Pharmacol* 37:141–149. doi:[10.1016/j.etap.2013.11.015](https://doi.org/10.1016/j.etap.2013.11.015)
- Lucioli J, Pinton P, Callu P et al (2013) The food contaminant deoxynivalenol activates the mitogen activated protein kinases in the intestine: interest of ex vivo models as an alternative to in vivo experiments. *Toxicol* 66:31–36. doi:[10.1016/j.toxicol.2013.01.024](https://doi.org/10.1016/j.toxicol.2013.01.024)
- Ma YY, Guo HW (2008) Mini-review of studies on the carcinogenicity of deoxynivalenol. *Environ Toxicol Pharmacol* 25:1–9. doi:[10.1016/j.etap.2007.09.007](https://doi.org/10.1016/j.etap.2007.09.007)
- Ma Y, Zhang A, Shi Z et al (2012) A mitochondria-mediated apoptotic pathway induced by deoxynivalenol in human colon cancer cells. *Toxicol Vitr* 26:414–420. doi:[10.1016/j.tiv.2012.01.010](https://doi.org/10.1016/j.tiv.2012.01.010)
- Malekinejad H, Schoevers EJ, Daemen IJMM et al (2007) Exposure of oocytes to the Fusarium toxins zearalenone and deoxynivalenol causes aneuploidy and abnormal embryo development in pigs. *Biol Reprod* 77:840–847. doi:[10.1095/biolreprod.107.062711](https://doi.org/10.1095/biolreprod.107.062711)
- Maresca M (2013) From the gut to the brain: journey and pathophysiological effects of the food-associated trichothecene mycotoxin deoxynivalenol. *Toxins* 5:784–820. doi:[10.3390/toxins5040784](https://doi.org/10.3390/toxins5040784)
- Maresca M, Fantini J (2010) Some food-associated mycotoxins as potential risk factors in humans predisposed to chronic intestinal inflammatory diseases. *Toxicol* 56:282–294. doi:[10.1016/j.toxicol.2010.04.016](https://doi.org/10.1016/j.toxicol.2010.04.016)
- Maresca M, Mahfoud R, Garmy N, Fantini J (2002) The mycotoxin deoxynivalenol affects nutrient absorption in human intestinal epithelial cells. *J Nutr* 132:2723–2731
- Maresca M, Yahi N, Younes-Sakr L et al (2008) Both direct and indirect effects account for the pro-inflammatory activity of enteropathogenic mycotoxins on the human intestinal epithelium: stimulation of interleukin-8 secretion, potentiation of interleukin-1beta effect and increase in the trans-epithelial passage of commensal bacteria. *Toxicol Appl Pharmacol* 228:84–92. doi:[10.1016/j.taap.2007.11.013](https://doi.org/10.1016/j.taap.2007.11.013)
- Maul R, Warth B, Kant J-S et al (2012) Investigation of the hepatic glucuronidation pattern of the Fusarium mycotoxin deoxynivalenol in various species. *Chem Res Toxicol* 25:2715–2717. doi:[10.1021/tx300348x](https://doi.org/10.1021/tx300348x)
- Meinild A, Klaerke DA, Loo DD et al (1998) The human Na<sup>+</sup>-glucose cotransporter is a molecular water pump. *J Physiol*. doi:[10.1111/j.1469-7793.1998.015br.x](https://doi.org/10.1111/j.1469-7793.1998.015br.x)
- Mishra S, Dwivedi PD, Pandey HP, Das M (2014) Role of oxidative stress in Deoxynivalenol induced toxicity. *Food Chem Toxicol*. 72:20–29. doi:[10.1016/j.fct.2014.06.027](https://doi.org/10.1016/j.fct.2014.06.027)
- Moon Y, Pestka JJ (2002) Vomitoxin-induced cyclooxygenase-2 gene expression in macrophages mediated by activation of ERK and p38 but not JNK mitogen-activated protein kinases. *Toxicol Sci* 69:373–382. doi:[10.1093/toxsci/69.2.373](https://doi.org/10.1093/toxsci/69.2.373)
- Morrissey RE, Vesonder RF (1985) Effect of deoxynivalenol (vomitoxin) on fertility, pregnancy, and postnatal development of Sprague-Dawley rats. *Appl Environ Microbiol* 49:1062–1066
- Nagl V, Schwartz H, Krska R et al (2012) Metabolism of the masked mycotoxin deoxynivalenol-3-glucoside in rats. *Toxicol Lett* 213:367–373. doi:[10.1016/j.toxlet.2012.07.024](https://doi.org/10.1016/j.toxlet.2012.07.024)
- Nagl V, Woechtl B, Schwartz-Zimmermann HE et al (2014) Metabolism of the masked mycotoxin deoxynivalenol-3-glucoside in pigs. *Toxicol Lett* 229:190–197. doi:[10.1016/j.toxlet.2014.06.032](https://doi.org/10.1016/j.toxlet.2014.06.032)
- Osselaere A, Devreese M, Goossens J et al (2013a) Toxicokinetic study and absolute oral bioavailability of deoxynivalenol, T-2 toxin and zearalenone in broiler chickens. *Food Chem Toxicol* 51:350–355. doi:[10.1016/j.fct.2012.10.006](https://doi.org/10.1016/j.fct.2012.10.006)
- Osselaere A, Santos R, Hautekiet V et al (2013b) Deoxynivalenol impairs hepatic and intestinal gene expression of selected oxidative stress, tight junction and inflammation proteins in broiler chickens, but addition of an adsorbing agent shifts the effects to the distal parts of the small intestine. *PLoS One* 8:e69014. doi:[10.1371/journal.pone.0069014](https://doi.org/10.1371/journal.pone.0069014)
- Ossenkopp KP, Hirst M, Rapley WA (1994) Deoxynivalenol (vomitoxin)-induced conditioned taste aversions in rats are mediated by the chemosensitive area postrema. *Pharmacol Biochem Behav* 47:363–367. doi:[10.1016/0091-3057\(94\)90024-8](https://doi.org/10.1016/0091-3057(94)90024-8)
- Parent-Massin D (2004) Haematotoxicity of trichothecenes. *Toxicol Lett* 153:75–81. doi:[10.1016/j.toxlet.2004.04.024](https://doi.org/10.1016/j.toxlet.2004.04.024)
- Paulick M, Rempe I, Kersten S et al (2015a) Effects of increasing concentrations of sodium sulfite on deoxynivalenol and deoxynivalenol sulfonate concentrations of maize kernels and maize meal preserved at various moisture content. *Toxins* 7:791–811. doi:[10.3390/toxins7030791](https://doi.org/10.3390/toxins7030791)
- Paulick M, Winkler J, Kersten S et al (2015b) Studies on the bioavailability of deoxynivalenol (DON) and DON sulfonate (DONS) 1, 2, and 3 in pigs fed with sodium sulfite-treated DON-contaminated maize. *Toxins* 7:4622–4644. doi:[10.3390/toxins7114622](https://doi.org/10.3390/toxins7114622)
- Payros D, Oswald E, Oswald IP (2016) Food contaminant exacerbates the genotoxicity of gut microbiota. (**submitted**)
- Pelyhe C, Kovesi B, Zandoki E et al (2016) Effect of 4-week feeding of deoxynivalenol- or T-2-toxin-contaminated diet on lipid peroxidation and glutathione redox system in the hepatopancreas of common carp (*Cyprinus carpio* L.). *Mycotoxin Res* 32:77–83. doi:[10.1007/s12550-016-0242-1](https://doi.org/10.1007/s12550-016-0242-1)
- Pestka JJ (2003) Deoxynivalenol-induced IgA production and IgA nephropathy-aberrant mucosal immune response with systemic repercussions. *Toxicol Lett* 140–141:287–295. doi:[10.1016/S0378-4274\(03\)00024-9](https://doi.org/10.1016/S0378-4274(03)00024-9)
- Pestka JJ (2007) Deoxynivalenol: toxicity, mechanisms and animal health risks. *Anim Feed Sci Technol* 137:283–298. doi:[10.1016/j.anifeedsci.2007.06.006](https://doi.org/10.1016/j.anifeedsci.2007.06.006)
- Pestka JJ (2008) Mechanisms of deoxynivalenol-induced gene expression and apoptosis. *Food Addit Contam* 25:1128–1140. doi:[10.1080/02652030802056626](https://doi.org/10.1080/02652030802056626)
- Pestka JJ (2010) Deoxynivalenol: mechanisms of action, human exposure, and toxicological relevance. *Arch Toxicol* 84:663–679. doi:[10.1007/s00204-010-0579-8](https://doi.org/10.1007/s00204-010-0579-8)
- Pestka JJ, Amuzie CJ (2008) Tissue distribution and proinflammatory cytokine gene expression following acute oral exposure to deoxynivalenol: comparison of weanling and adult mice. *Food Chem Toxicol* 46:2826–2831. doi:[10.1016/j.fct.2008.05.016](https://doi.org/10.1016/j.fct.2008.05.016)
- Pestka JJ, Smolinski AT (2005) Deoxynivalenol: toxicology and potential effects on humans. *J Toxicol Environ Health B Crit Rev* 8:39–69. doi:[10.1080/10937400590889458](https://doi.org/10.1080/10937400590889458)
- Pestka JJ, Lin WS, Miller ER (1987) Emetic activity of the trichothecene 15-acetyldeoxynivalenol in swine. *Food Chem Toxicol* 25:855–858. doi:[10.1016/0278-6915\(87\)90264-X](https://doi.org/10.1016/0278-6915(87)90264-X)

- Pestka JJ, Zhou HR, Moon Y, Chung YJ (2004) Cellular and molecular mechanisms for immune modulation by deoxynivalenol and other trichothecenes: unraveling a paradox. *Toxicol Lett* 153:61–73. doi:[10.1016/j.toxlet.2004.04.023](https://doi.org/10.1016/j.toxlet.2004.04.023)
- Pestka JJ, Uzarski RL, Islam Z (2005) Induction of apoptosis and cytokine production in the Jurkat human T cells by deoxynivalenol: role of mitogen-activated protein kinases and comparison to other 8-ketotrichothecenes. *Toxicology* 206:207–219. doi:[10.1016/j.tox.2004.08.020](https://doi.org/10.1016/j.tox.2004.08.020)
- Pierron A, Alassane-Kpembi I, Oswald IP (2016a) Impact of mycotoxin on immune response and consequences for pig health. *Anim Nutr* 2:63–68. doi:[10.1016/j.aninu.2016.03.001](https://doi.org/10.1016/j.aninu.2016.03.001)
- Pierron A, Mimoun S, Murate LS et al (2016b) Intestinal toxicity of the masked mycotoxin deoxynivalenol-3-beta-D-glucoside. *Arch Toxicol* 90:2037–2046. doi:[10.1007/s00204-015-1592-8](https://doi.org/10.1007/s00204-015-1592-8)
- Pierron A, Mimoun S, Murate LS et al (2016c) Microbial biotransformation of DON: molecular basis for reduced toxicity. *Sci Rep* 6:29105. doi:[10.1038/srep29105](https://doi.org/10.1038/srep29105)
- Pinton P, Oswald IP (2014) Effect of deoxynivalenol and other Type B trichothecenes on the intestine: a review. *Toxins* 6:1615–1643. doi:[10.3390/toxins6051615](https://doi.org/10.3390/toxins6051615)
- Pinton P, Accensi F, Beauchamp E et al (2008) Ingestion of deoxynivalenol (DON) contaminated feed alters the pig vaccinal immune responses. *Toxicol Lett* 177:215–222. doi:[10.1016/j.toxlet.2008.01.015](https://doi.org/10.1016/j.toxlet.2008.01.015)
- Pinton P, Nougayrede JP, Del Rio JC et al (2009) The food contaminant deoxynivalenol, decreases intestinal barrier permeability and reduces claudin expression. *Toxicol Appl Pharmacol* 237:41–48. doi:[10.1016/j.taap.2009.03.003](https://doi.org/10.1016/j.taap.2009.03.003)
- Pinton P, Tsybul'skyy D, Lucieli J et al (2012) Toxicity of deoxynivalenol and its acetylated derivatives on the intestine: differential effects on morphology, barrier function, tight junction proteins, and mitogen-activated protein kinases. *Toxicol Sci* 130:180–190. doi:[10.1093/toxsci/kfs239](https://doi.org/10.1093/toxsci/kfs239)
- Pinton P, Graziani F, Pujol A et al (2015) Deoxynivalenol inhibits the expression by goblet cells of intestinal mucins through a PKR and MAP kinase dependent repression of the resistin-like molecule beta. *Mol Nutr Food Res* 59:1076–1087. doi:[10.1002/mnfr.201500005](https://doi.org/10.1002/mnfr.201500005)
- Piotrowska M, Slizewska K, Nowak A et al (2014) The effect of experimental fusarium mycotoxicosis on microbiota diversity in porcine ascending colon contents. *Toxins* 6:2064–2081. doi:[10.3390/toxins6072064](https://doi.org/10.3390/toxins6072064)
- Poppenberger B, Berthiller F, Lucyshyn D et al (2003) Detoxification of the Fusarium mycotoxin deoxynivalenol by a UDP-glucosyltransferase from *Arabidopsis thaliana*. *J Biol Chem* 278:47905–47914. doi:[10.1074/jbc.M307552200M307552200](https://doi.org/10.1074/jbc.M307552200M307552200)
- Prelusky DB, Veira DM, Trenholm HL (1985) Plasma pharmacokinetics of the mycotoxin deoxynivalenol following oral and intravenous administration to sheep. *J Environ Sci Health B* 20:603–624. doi:[10.1080/03601238509372499](https://doi.org/10.1080/03601238509372499)
- Prelusky DB, Veira DM, Trenholm HL, Hartin KE (1986) Excretion profiles of the mycotoxin deoxynivalenol, following oral and intravenous administration to sheep. *Fundam Appl Toxicol* 6:356–363. doi:[10.1016/0272-0590\(86\)90251-4](https://doi.org/10.1016/0272-0590(86)90251-4)
- Prelusky DB, Hartin KE, Trenholm HL, Miller JD (1988) Pharmacokinetic fate of <sup>14</sup>C-labeled deoxynivalenol in swine. *Fundam Appl Toxicol* 10:276–286
- Razafimanjato H, Benzaria A, Taieb N et al (2011) The ribotoxin deoxynivalenol affects the viability and functions of glial cells. *Glia* 59:1672–1683. doi:[10.1002/glia.21214](https://doi.org/10.1002/glia.21214)
- Rizzo AF, Atroshi F, Ahotupa M et al (1994) Protective effect of antioxidants against free radical-mediated lipid peroxidation induced by DON or T-2 toxin. *Zentralbl Veterinarmed A* 41:81–90. doi:[10.1111/j.1439-0442.1994.tb00070.x](https://doi.org/10.1111/j.1439-0442.1994.tb00070.x)
- Rodriguez-Carrasco Y, Molto JC, Berrada H, Manes J (2014) A survey of trichothecenes, zearalenone and patulin in milled grain-based products using GC-MS/MS. *Food Chem* 146:212–219. doi:[10.1016/j.foodchem.2013.09.053](https://doi.org/10.1016/j.foodchem.2013.09.053)
- Rotter BA, Thompson BK, Prelusky DB, Trenholm HL (1991) Evaluation of potential interactions involving trichothecene mycotoxins using the chick embryotoxicity bioassay. *Arch Environ Contam Toxicol* 21:621–624
- Rychlik M, Humpf H-U, Marko D et al (2014) Proposal of a comprehensive definition of modified and other forms of mycotoxins including “masked” mycotoxins. *Mycotoxin Res* 30:197–205. doi:[10.1007/s12550-014-0203-5](https://doi.org/10.1007/s12550-014-0203-5)
- Saint-Cyr MJ, Perrin-Guyomard A, Houée P et al (2013) Evaluation of an oral subchronic exposure of deoxynivalenol on the composition of human gut microbiota in a model of human microbiota-associated rats. *PLoS One* 8:e80578. doi:[10.1371/journal.pone.0080578](https://doi.org/10.1371/journal.pone.0080578)
- Sarkanj B, Warth B, Uhlig S et al (2013) Urinary analysis reveals high deoxynivalenol exposure in pregnant women from Croatia. *Food Chem Toxicol* 62:231–237. doi:[10.1016/j.fct.2013.08.043](https://doi.org/10.1016/j.fct.2013.08.043)
- Sato I, Ito M, Ishizaka M et al (2012) Thirteen novel deoxynivalenol-degrading bacteria are classified within two genera with distinct degradation mechanisms. *FEMS Microbiol Lett* 327:110–117. doi:[10.1111/j.1574-6968.2011.02461.x](https://doi.org/10.1111/j.1574-6968.2011.02461.x)
- Savard C, Provost C, Alvarez F et al (2015) Effect of deoxynivalenol (DON) mycotoxin on in vivo and in vitro porcine circovirus type 2 infections. *Vet Microbiol* 176:257–267. doi:[10.1016/j.vetmic.2015.02.004](https://doi.org/10.1016/j.vetmic.2015.02.004)
- Schatzmayr G, Zehner F, Taubel M et al (2006) Microbiologicals for deactivating mycotoxins. *Mol Nutr Food Res* 50:543–551. doi:[10.1002/mnfr.200500181](https://doi.org/10.1002/mnfr.200500181)
- Schwartz HE, Hametner C, Slavik V et al (2013) Characterization of three deoxynivalenol sulfonates formed by reaction of deoxynivalenol with sulfur reagents. *J Agric Food Chem* 61:8941–8948. doi:[10.1021/jf403438b](https://doi.org/10.1021/jf403438b)
- Schwartz-Zimmermann HE, Hametner C, Nagl V et al (2014a) Deoxynivalenol (DON) sulfonates as major DON metabolites in rats: from identification to biomarker method development, validation and application. *Anal Bioanal Chem* 406:7911–7924. doi:[10.1007/s00216-014-8252-3](https://doi.org/10.1007/s00216-014-8252-3)
- Schwartz-Zimmermann HE, Wiesenberger G, Unbekannt C et al (2014b) Reaction of (conjugated) deoxynivalenol with sulphur reagents—novel metabolites, toxicity and application. *World Mycotoxin J* 7:187–197. doi:[10.3920/WMJ2013.1632](https://doi.org/10.3920/WMJ2013.1632)
- Schwartz-Zimmermann HE, Fruhmann P, Danicke S et al (2015) Metabolism of deoxynivalenol and de-epoxy-deoxynivalenol in broiler chickens, pullets, roosters and turkeys. *Toxins* 7:4706–4729. doi:[10.3390/toxins7114706](https://doi.org/10.3390/toxins7114706)
- Seeling K, Danicke S, Valenta H et al (2006) Effects of Fusarium toxin-contaminated wheat and feed intake level on the biotransformation and carry-over of deoxynivalenol in dairy cows. *Food Addit Contam* 23:1008–1020. doi:[10.1080/02652030600723245](https://doi.org/10.1080/02652030600723245)
- Severino L, Luongo D, Bergamo P et al (2006) Mycotoxins nivale-nol and deoxynivalenol differentially modulate cytokine mRNA expression in Jurkat T cells. *Cytokine* 36:75–82. doi:[10.1016/j.cyto.2006.11.006](https://doi.org/10.1016/j.cyto.2006.11.006)
- Shima J, Takase S, Takahashi Y et al (1997) Novel detoxification of the trichothecene mycotoxin deoxynivalenol by a soil bacterium isolated by enrichment culture. *Appl Environ Microbiol* 63:3825–3830
- Singh S, Banerjee S, Chattopadhyay P et al (2015) Deoxynivalenol induces cytotoxicity and genotoxicity in animal primary cell culture. *Toxicol Mech Methods* 6516:1–8. doi:[10.3109/15376516.2015.1006743](https://doi.org/10.3109/15376516.2015.1006743)

- Sprando RL, Collins TF, Black TN et al (2005) Characterization of the effect of deoxynivalenol on selected male reproductive endpoints. *Food Chem Toxicol* 43:623–635. doi:10.1016/j.fct.2004.12.017
- Streit E, Naehrer K, Rodrigues I, Schatzmayr G (2013) Mycotoxin occurrence in feed and feed raw materials worldwide: long-term analysis with special focus on Europe and Asia. *J Sci Food Agric* 93:2892–2899. doi:10.1002/jsfa.6225
- Sugita-Konishi Y, Pestka JJ (2001) Differential upregulation of TNF- $\alpha$ , IL-6, and IL-8 production by deoxynivalenol (vomitoxin) and other 8-ketotrichothecenes in a human macrophage model. *J Toxicol Environ Health A* 64:619–636. doi:10.1080/152873901753246223
- Sugita-Konishi Y, Park BJ, Kobayashi-Hattori K et al (2006) Effect of cooking process on the deoxynivalenol content and its subsequent cytotoxicity in wheat products. *Biosci Biotechnol Biochem* 70:1764–1768. doi:10.1271/bbb.50571
- Sundstol Eriksen G, Pettersson H (2003) Lack of de-epoxidation of type B trichothecenes in incubates with human faeces. *Food Addit Contam* 20:579–582. doi:10.1080/0265203031000102573
- Swanson SP, Nicoletti J, Rood HDJ et al (1987) Metabolism of three trichothecene mycotoxins, T-2 toxin, diacetoxyscirpenol and deoxynivalenol, by bovine rumen microorganisms. *J Chromatogr* 414:335–342
- Taranu I, Marina DE, Burlacu R et al (2010) Comparative aspects of in vitro proliferation of human and porcine lymphocytes exposed to mycotoxins. *Arch Anim Nutr* 64:383–393. doi:10.1080/1745039X.2010.492140
- Tomar RS, Blakley BR, Schiefer HB, DeCoteau WE (1986) In vitro effects of 3-acetyl-deoxynivalenol on the immune response of human peripheral blood lymphocytes. *Int J Immunopharmacol* 8:125–130. doi:10.1016/0192-0561(86)90051-2
- Tomar RS, Blakley BR, DeCoteau WE (1987) Immunological responsiveness of mouse spleen cells after in vivo or in vitro exposure to 3-acetyldeoxynivalenol. *Food Chem Toxicol* 25:393–8. doi:10.1016/0278-6915(87)90175-X
- Turner PC, Hopton RP, Lecluse Y et al (2010) Determinants of urinary deoxynivalenol and de-epoxy deoxynivalenol in male farmers from Normandy, France. *J Agric Food Chem* 58:5206–5212. doi:10.1021/jf100892v
- Uhlig S, Ivanova L, Faeste CK (2016) Correction to enzyme-assisted synthesis and structural characterization of the 3-, 8-, and 15-glucuronides of deoxynivalenol. *J Agric Food Chem* 64:3732. doi:10.1021/acs.jafc.6b01413
- van der Flier LG, Clevers H (2009) Stem cells, self-renewal, and differentiation in the intestinal epithelium. *Annu Rev Physiol* 71:241–260. doi:10.1146/annurev.physiol.010908.163145
- Vandenbroucke V, Croubels S, Verbrugghe E et al (2009) The mycotoxin deoxynivalenol promotes uptake of *Salmonella Typhimurium* in porcine macrophages, associated with ERK1/2 induced cytoskeleton reorganization. *Vet Res* 40:64. doi:10.1051/vetres/2009045
- Vandenbroucke V, Croubels S, Martel A et al (2011) The mycotoxin deoxynivalenol potentiates intestinal inflammation by *Salmonella typhimurium* in porcine ileal loops. *PLoS One* 6:e23871. doi:10.1371/journal.pone.0023871
- Versilovskis A, Bartkevics V (2012) Stability of sterigmatocystin during the bread making process and its occurrence in bread from the Latvian market. *Mycotoxin Res* 28:123–129. doi:10.1007/s12550-012-0124-0
- Volkl A, Vogler B, Schollenberger M, Karlovsky P (2004) Microbial detoxification of mycotoxin deoxynivalenol. *J Basic Microbiol* 44:147–156. doi:10.1002/jobm.200310353
- Wache YJ, Hbabi-Haddioui L, Guzylack-Piriou L et al (2009a) The mycotoxin Deoxynivalenol inhibits the cell surface expression of activation markers in human macrophages. *Toxicology* 262:239–244. doi:10.1016/j.tox.2009.06.014
- Wache YJ, Valat C, Postollec G et al (2009b) Impact of deoxynivalenol on the intestinal microflora of pigs. *Int J Mol Sci* 10:1–17. doi:10.3390/ijms10010001
- Wan D, Huang L, Pan Y et al (2013) Metabolism, distribution, and excretion of deoxynivalenol with combined techniques of radiotracing, high-performance liquid chromatography ion trap time-of-flight mass spectrometry, and online radiometric detection. *J Agric Food Chem* 62:288–296. doi:10.1021/jf4047946
- Wu X, Kohut M, Cunnick J et al (2009) Deoxynivalenol suppresses circulating and splenic leukocyte subpopulations in BALB/c mice: dose response, time course and sex differences. *Food Addit Contam Part A Chem Anal Control Expo Risk Assess* 26:1070–1080. doi:10.1080/02652030902832959
- Wu W, Flannery BM, Sugita-Konishi Y et al (2012) Comparison of murine anorectic responses to the 8-ketotrichothecenes 3-acetyldeoxynivalenol, 15-acetyldeoxynivalenol, fusarenon X and nivalenol. *Food Chem Toxicol* 50:2056–2061. doi:10.1016/j.fct.2012.03.055
- Wu W, Bates MA, Bursian SJ et al (2013a) Peptide YY3-36 and 5-hydroxytryptamine mediate emesis induction by trichothecene deoxynivalenol (vomitoxin). *Toxicol Sci* 133:186–195. doi:10.1093/toxsci/kf033
- Wu W, Bates MA, Bursian SJ et al (2013b) Comparison of emetic potencies of the 8-ketotrichothecenes deoxynivalenol, 15-acetyldeoxynivalenol, 3-acetyldeoxynivalenol, fusarenon X, and nivalenol. *Toxicol Sci* 131:279–291. doi:10.1093/toxsci/kfs286
- Wu M, Xiao H, Ren W et al (2014a) Therapeutic effects of glutamic acid in piglets challenged with deoxynivalenol. *PLoS One* 9:e100591. doi:10.1371/journal.pone.0100591
- Wu QH, Wang X, Yang W et al (2014b) Oxidative stress-mediated cytotoxicity and metabolism of T-2 toxin and deoxynivalenol in animals and humans: an update. *Arch Toxicol* 88:1309–1326. doi:10.1007/s00204-014-1280-0
- Wu W, He K, Zhou HR et al (2014c) Effects of oral exposure to naturally-occurring and synthetic deoxynivalenol congeners on proinflammatory cytokine and chemokine mRNA expression in the mouse. *Toxicol Appl Pharmacol*. doi:10.1016/j.taap.2014.04.016
- Wu W, Zhou H-R, He K et al (2014d) Role of cholecystokinin in anorexia induction following oral exposure to the 8-ketotrichothecenes deoxynivalenol, 15-acetyldeoxynivalenol, 3-acetyldeoxynivalenol, fusarenon X, and nivalenol. *Toxicol Sci* 138:278–289. doi:10.1093/toxsci/kft335
- Wu L, Liao P, He L et al (2015) Growth performance, serum biochemical profile, jejunal morphology, and the expression of nutrients transporter genes in deoxynivalenol (DON)-challenged growing pigs. *BMC Vet Res* 11:144. doi:10.1186/s12917-015-0449-y
- Wu W, Zhou H-R, Pestka JJ (2016) Potential roles for calcium-sensing receptor (CaSR) and transient receptor potential ankyrin-1 (TRPA1) in murine anorectic response to deoxynivalenol (vomitoxin). *Arch Toxicol*. doi:10.1007/s00204-016-1687-x
- Xi P, Jiang Z, Dai Z et al (2011) Regulation of protein turnover by L-glutamine in porcine intestinal epithelial cells. *J Nutr Biochem*. doi:10.1016/j.jnutbio.2011.05.009
- Yang GH, Jarvis BB, Chung YJ, Pestka JJ (2000) Apoptosis induction by the satratoxins and other trichothecene mycotoxins: relationship to ERK, p38 MAPK, and SAPK/JNK activation. *Toxicol Appl Pharmacol* 164:149–160. doi:10.1006/taap.1999.8888
- Yang H, Chung DH, Kim YB et al (2008) Ribotoxic mycotoxin deoxynivalenol induces G(2)/M cell cycle arrest via p21(Cip/WAF1) mRNA stabilization in human epithelial cells. *Toxicology* 243:145–154. doi:10.1016/j.tox.2007.10.002

- Yoshizawa T, Takeda H, Ohi T (1983) Structure of a novel metabolite from Deoxynivalenol, a trichothecene mycotoxin, in animals. *Agric Biol Chem* 47:2133–2135. doi:[10.1080/00021369.1983.10865926](https://doi.org/10.1080/00021369.1983.10865926)
- Young JC, Blackwell BA, ApSimon JW (1986) Alkaline degradation of the mycotoxin 4-deoxynivalenol. *Tetrahedron Lett* 27:1019–1022. doi:[10.1016/S0040-4039\(86\)80037-5](https://doi.org/10.1016/S0040-4039(86)80037-5)
- Young JC, Zhou T, Yu H et al (2007) Degradation of trichothecene mycotoxins by chicken intestinal microbes. *Food Chem Toxicol* 45:136–143. doi:[10.1016/j.fct.2006.07.028](https://doi.org/10.1016/j.fct.2006.07.028)
- Yu H, Zhou T, Gong J et al (2010) Isolation of deoxynivalenol-transforming bacteria from the chicken intestines using the approach of PCR-DGGE guided microbial selection. *BMC Microbiol* 10:182. doi:[10.1186/1471-2180-10-182](https://doi.org/10.1186/1471-2180-10-182)
- Yunus AW, Ghareeb K, Twaruzek M et al (2012) Deoxynivalenol as a contaminant of broiler feed: effects on bird performance and response to common vaccines. *Poult Sci* 91:844–851. doi:[10.3382/ps.2011-01873](https://doi.org/10.3382/ps.2011-01873)
- Zhang X, Jiang L, Geng C et al (2009) The role of oxidative stress in deoxynivalenol-induced DNA damage in HepG2 cells. *Toxicon* 54:513–518. doi:[10.1016/j.toxicon.2009.05.021](https://doi.org/10.1016/j.toxicon.2009.05.021)
- Zhou T, He J (2009) Bacterial isolate, methods of isolating bacterial isolates and Methods for detoxification of trichothecene mycotoxins. US Patent 61/249,023, 6 Oct 2009
- Zhou HR, He K, Landgraf J et al (2014) Direct activation of ribosome-associated double-stranded RNA-dependent protein kinase (PKR) by deoxynivalenol, anisomycin and ricin: a new model for ribotoxic stress response induction. *Toxins* 6:3406–3425. doi:[10.3390/toxins6123406](https://doi.org/10.3390/toxins6123406)